

Regina Rodrigo

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/419247/regina-rodrigo-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

60
papers

2,187
citations

26
h-index

46
g-index

66
ext. papers

2,444
ext. citations

5.1
avg. IF

4.37
L-index

#	Paper	IF	Citations
60	NUTRARET: Effect of 2-Year Nutraceutical Supplementation on Redox Status and Visual Function of Patients With Retinitis Pigmentosa: A Randomized, Double-Blind, Placebo-Controlled Trial.. <i>Frontiers in Nutrition</i> , 2022 , 9, 847910	6.2	0
59	Nutraceutical Supplementation Ameliorates Visual Function, Retinal Degeneration, and Redox Status in rd10 Mice. <i>Antioxidants</i> , 2021 , 10,	7.1	5
58	[Decreased DNA hydroxymethylation and increased DNA demethylation are associated with high antioxidant response in systemic lupus erythematosus patients]. <i>Medicina Clínica</i> , 2021 , 157, 575-579	1	
57	Retinal Inflammation, Cell Death and Inherited Retinal Dystrophies. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	14
56	Oxidative Stress, a Crossroad Between Rare Diseases and Neurodegeneration. <i>Antioxidants</i> , 2020 , 9,	7.1	20
55	Intravitreal administration of adalimumab delays retinal degeneration in rd10 mice. <i>FASEB Journal</i> , 2020 , 34, 13839-13861	0.9	7
54	Redox Status, Dose and Antioxidant Intake in Healthcare Workers Occupationally Exposed to Ionizing Radiation. <i>Antioxidants</i> , 2020 , 9,	7.1	1
53	HIF-1 β stabilization reduces retinal degeneration in a mouse model of retinitis pigmentosa. <i>FASEB Journal</i> , 2018 , 32, 2438-2451	0.9	16
52	Bioelectrochemical monitoring of soluble guanylate cyclase inhibition by the natural β -carboline canthin-6-one. <i>Journal of Molecular Structure</i> , 2017 , 1134, 661-667	3.4	3
51	cGMP-Phosphodiesterase Inhibition Prevents Hypoxia-Induced Cell Death Activation in Porcine Retinal Explants. <i>PLoS ONE</i> , 2016 , 11, e0166717	3.7	10
50	UV protection from cotton fabrics dyed with different tea extracts. <i>Dyes and Pigments</i> , 2016 , 134, 448-452	4.26	41
49	Adalimumab Reduces Photoreceptor Cell Death in A Mouse Model of Retinal Degeneration. <i>Scientific Reports</i> , 2015 , 5, 11764	4.9	35
48	Targeted next generation sequencing for molecular diagnosis of Usher syndrome. <i>Orphanet Journal of Rare Diseases</i> , 2014 , 9, 168	4.2	52
47	Infliximab reduces Zaprinst-induced retinal degeneration in cultures of porcine retina. <i>Journal of Neuroinflammation</i> , 2014 , 11, 172	10.1	14
46	Cerebral oedema is not responsible for motor or cognitive deficits in rats with hepatic encephalopathy. <i>Liver International</i> , 2014 , 34, 379-87	7.9	23
45	Phosphodiesterase inhibition induces retinal degeneration, oxidative stress and inflammation in cone-enriched cultures of porcine retina. <i>Experimental Eye Research</i> , 2013 , 111, 122-33	3.7	20
44	Study of USH1 splicing variants through minigenes and transcript analysis from nasal epithelial cells. <i>PLoS ONE</i> , 2013 , 8, e57506	3.7	20

43	Altered antioxidant-oxidant status in the aqueous humor and peripheral blood of patients with retinitis pigmentosa. <i>PLoS ONE</i> , 2013 , 8, e74223	3.7	52
42	Two novel disease-causing mutations in the CLRN1 gene in patients with Usher syndrome type 3. <i>Molecular Vision</i> , 2012 , 18, 3070-8	2.3	9
41	Brain region-selective mechanisms contribute to the progression of cerebral alterations in acute liver failure in rats. <i>Gastroenterology</i> , 2011 , 140, 638-45	13.3	50
40	Haemodynamic effects of long-term administration of sildenafil in normotensive pregnant and non-pregnant rats. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2011 , 118, 615-23	3.7	12
39	cGMP modulates stem cells differentiation to neurons in brain in vivo pathological implications. <i>BMC Pharmacology</i> , 2011 , 11,		1
38	Mutational screening of the USH2A gene in Spanish USH patients reveals 23 novel pathogenic mutations. <i>Orphanet Journal of Rare Diseases</i> , 2011 , 6, 65	4.2	41
37	p38 MAP kinase is a therapeutic target for hepatic encephalopathy in rats with portacaval shunts. <i>Gut</i> , 2011 , 60, 1572-9	19.2	49
36	Chronic hyperammonemia induces tonic activation of NMDA receptors in cerebellum. <i>Journal of Neurochemistry</i> , 2010 , 112, 1005-14	6	25
35	Hyperammonemia induces neuroinflammation that contributes to cognitive impairment in rats with hepatic encephalopathy. <i>Gastroenterology</i> , 2010 , 139, 675-84	13.3	224
34	cGMP modulates stem cells differentiation to neurons in brain in vivo. <i>Neuroscience</i> , 2010 , 165, 1275-83	3.9	29
33	Cyclic GMP pathways in hepatic encephalopathy. Neurological and therapeutic implications. <i>Metabolic Brain Disease</i> , 2010 , 25, 39-48	3.9	32
32	Novel mutations in the USH1C gene in Usher syndrome patients. <i>Molecular Vision</i> , 2010 , 16, 2948-54	2.3	4
31	Neuroinflammation contributes to hypokinesia in rats with hepatic encephalopathy: ibuprofen restores its motor activity. <i>Journal of Neuroscience Research</i> , 2009 , 87, 1369-74	4.4	55
30	Glutamatergic and gabaergic neurotransmission and neuronal circuits in hepatic encephalopathy. <i>Metabolic Brain Disease</i> , 2009 , 24, 69-80	3.9	103
29	Polychlorinated biphenyls PCB 153 and PCB 126 impair the glutamate-nitric oxide-cGMP pathway in cerebellar neurons in culture by different mechanisms. <i>Neurotoxicity Research</i> , 2009 , 16, 97-105	4.3	15
28	Role of NMDA receptors in acute liver failure and ammonia toxicity: therapeutical implications. <i>Neurochemistry International</i> , 2009 , 55, 113-8	4.4	48
27	Mechanisms of cognitive alterations in hyperammonemia and hepatic encephalopathy: therapeutical implications. <i>Neurochemistry International</i> , 2009 , 55, 106-12	4.4	49
26	Chronic hyperammonemia reduces the activity of neuronal nitric oxide synthase in cerebellum by altering its localization and increasing its phosphorylation by calcium-calmodulin kinase II. <i>Journal of Neurochemistry</i> , 2008 , 106, 1440-9	6	36

25	Brain cholinergic impairment in liver failure. <i>Brain</i> , 2008 , 131, 2946-56	11.2	74
24	Acute liver failure-induced death of rats is delayed or prevented by blocking NMDA receptors in brain. <i>American Journal of Physiology - Renal Physiology</i> , 2008 , 295, G503-11	5.1	26
23	Inflammation and hepatic encephalopathy: ibuprofen restores learning ability in rats with portacaval shunts. <i>Hepatology</i> , 2007 , 46, 514-9	11.2	162
22	Glutamate-induced activation of nitric oxide synthase is impaired in cerebral cortex in vivo in rats with chronic liver failure. <i>Journal of Neurochemistry</i> , 2007 , 102, 51-64	6	31
21	Hyperammonaemia alters the mechanisms by which metabotropic glutamate receptors in nucleus accumbens modulate motor function. <i>Journal of Neurochemistry</i> , 2007 , 103, 38-46	6	27
20	NMDA receptors in hyperammonemia and hepatic encephalopathy. <i>Metabolic Brain Disease</i> , 2007 , 22, 321-35	3.9	52
19	Control of brain glutamine synthesis by NMDA receptors. <i>Frontiers in Bioscience - Landmark</i> , 2007 , 12, 883-90	2.8	24
18	Brain edema and inflammatory activation in bile duct ligated rats with diet-induced hyperammonemia: A model of hepatic encephalopathy in cirrhosis. <i>Hepatology</i> , 2006 , 43, 1257-66	11.2	122
17	Brain regional alterations in the modulation of the glutamate-nitric oxide-cGMP pathway in liver cirrhosis. Role of hyperammonemia and cell types involved. <i>Neurochemistry International</i> , 2006 , 48, 472-484	4.4	18
16	Mass spectrometrical analysis of galectin proteins in primary rat cerebellar astrocytes. <i>Neurochemical Research</i> , 2006 , 31, 945-55	4.6	7
15	Pharmacological manipulation of cyclic GMP levels in brain restores learning ability in animal models of hepatic encephalopathy: therapeutic implications. <i>Neuropsychiatric Disease and Treatment</i> , 2006 , 2, 53-63	3.1	5
14	Phosphate-activated glutaminase activity is enhanced in brain, intestine and kidneys of rats following portacaval anastomosis. <i>World Journal of Gastroenterology</i> , 2006 , 12, 2406-11	5.6	34
13	Neurons exposed to ammonia reproduce the differential alteration in nitric oxide modulation of guanylate cyclase in the cerebellum and cortex of patients with liver cirrhosis. <i>Neurobiology of Disease</i> , 2005 , 19, 150-61	7.5	23
12	Bile duct ligation plus hyperammonemia in rats reproduces the alterations in the modulation of soluble guanylate cyclase by nitric oxide in brain of cirrhotic patients. <i>Neuroscience</i> , 2005 , 130, 435-43	3.9	20
11	Proteome analysis of primary neurons and astrocytes from rat cerebellum. <i>Journal of Proteome Research</i> , 2005 , 4, 768-88	5.6	37
10	Oral administration of sildenafil restores learning ability in rats with hyperammonemia and with portacaval shunts. <i>Hepatology</i> , 2005 , 41, 299-306	11.2	131
9	Altered modulation of motor activity by group I metabotropic glutamate receptors in the nucleus accumbens in hyperammonemic rats. <i>Metabolic Brain Disease</i> , 2005 , 20, 347-58	3.9	8
8	Animal models in the study of episodic hepatic encephalopathy in cirrhosis. <i>Metabolic Brain Disease</i> , 2005 , 20, 399-408	3.9	22

7	Alterations in soluble guanylate cyclase content and modulation by nitric oxide in liver disease. <i>Neurochemistry International</i> , 2004 , 45, 947-53	4.4	21
6	Glutamine synthetase activity and glutamine content in brain: modulation by NMDA receptors and nitric oxide. <i>Neurochemistry International</i> , 2003 , 43, 493-9	4.4	121
5	Altered modulation of soluble guanylate cyclase in lymphocytes from patients with liver disease. <i>Journal of Molecular Medicine</i> , 2002 , 80, 117-23	5.5	14
4	Region selective alterations of soluble guanylate cyclase content and modulation in brain of cirrhotic patients. <i>Hepatology</i> , 2002 , 36, 1155-62	11.2	65
3	Selective regional alterations in the content or distribution of neuronal and glial cytoskeletal proteins in brain of rats chronically exposed to 2,5-hexanedione. <i>Toxicology and Industrial Health</i> , 2002 , 18, 333-41	1.8	2
2	Chronic hyperammonemia alters protein phosphorylation and glutamate receptor-associated signal transduction in brain. <i>Neurochemistry International</i> , 2002 , 41, 103-8	4.4	13
1	Chronic hyperammonemia in rats impairs activation of soluble guanylate cyclase in neurons and in lymphocytes: a putative peripheral marker for neurological alterations. <i>Biochemical and Biophysical Research Communications</i> , 1999 , 257, 405-9	3.4	13