

Milorad B Tesic

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4195367/publications.pdf>

Version: 2024-02-01

54
papers

625
citations

623188

14
h-index

676716

22
g-index

54
all docs

54
docs citations

54
times ranked

872
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | The retrograde technique for recanalization of chronically occluded coronary arteries: Case series report. <i>Vojnosanitetski Pregled</i> , 2022, 79, 503-509. | 0.1 | 1 |
| 2 | Left atrial volume changes during exercise stress echocardiography in heart failure and hypertrophic cardiomyopathy. <i>Hellenic Journal of Cardiology</i> , 2022, 67, 9-18. | 0.4 | 6 |
| 3 | Coronary Flow Velocity Reserve Using Dobutamine Test for Noninvasive Functional Assessment of Myocardial Bridging. <i>Journal of Clinical Medicine</i> , 2022, 11, 204. | 1.0 | 2 |
| 4 | The Coronary ARteriogenesis with combined Heparin and EXercise therapy in chronic refractory Angina (CARHEXA) trial: A double-blind, randomized, placebo-controlled stress echocardiographic study. <i>European Journal of Preventive Cardiology</i> , 2021, 28, 1452-1459. | 0.8 | 7 |
| 5 | Feasibility and functional correlates of left atrial volume changes during stress echocardiography in chronic coronary syndromes. <i>International Journal of Cardiovascular Imaging</i> , 2021, 37, 953-964. | 0.7 | 9 |
| 6 | Randomized Controlled Comparison of Optimal Medical Therapy with Percutaneous Recanalization of Chronic Total Occlusion (COMET-CTO). <i>International Heart Journal</i> , 2021, 62, 16-22. | 0.5 | 29 |
| 7 | Alpha-melanocyte-stimulating hormone during exercise recovery has prognostic value for coronary artery disease. <i>Hormones</i> , 2021, 20, 381-387. | 0.9 | 0 |
| 8 | Prognostic Value of Reduced Heart Rate Reserve during Exercise in Hypertrophic Cardiomyopathy. <i>Journal of Clinical Medicine</i> , 2021, 10, 1347. | 1.0 | 6 |
| 9 | Hemodynamic Heterogeneity of Reduced Cardiac Reserve Unmasked by Volumetric Exercise Echocardiography. <i>Journal of Clinical Medicine</i> , 2021, 10, 2906. | 1.0 | 6 |
| 10 | Physical activity and exercise as an essential medical strategy for the COVID-19 pandemic and beyond. <i>Experimental Biology and Medicine</i> , 2021, 246, 2324-2331. | 1.1 | 2 |
| 11 | Functional Assessment of Myocardial Bridging With Conventional and Diastolic Fractional Flow Reserve: Vasodilator Versus Inotropic Provocation. <i>Journal of the American Heart Association</i> , 2021, 10, e020597. | 1.6 | 21 |
| 12 | Contrast-induced nephropathy in a patient with type 2 diabetes and coronary artery disease: a case report. <i>Journal of International Medical Research</i> , 2021, 49, 030006052110331. | 0.4 | 1 |
| 13 | Stress Echo 2030: The Novel ABCDE-(FGLPR) Protocol to Define the Future of Imaging. <i>Journal of Clinical Medicine</i> , 2021, 10, 3641. | 1.0 | 33 |
| 14 | The Cardiology Society of Serbia. <i>European Heart Journal</i> , 2021, 42, 294-296. | 1.0 | 0 |
| 15 | Prognostic Value of Transthoracic Doppler Echocardiography Coronary Flow Velocity Reserve in Patients With Asymmetric Hypertrophic Cardiomyopathy. <i>Journal of the American Heart Association</i> , 2021, 10, e021936. | 1.6 | 12 |
| 16 | Lung Ultrasound and Pulmonary Congestion During Stress Echocardiography. <i>JACC: Cardiovascular Imaging</i> , 2020, 13, 2085-2095. | 2.3 | 53 |
| 17 | Prompt and consistent improvement of coronary flow velocity reserve following successful recanalization of the coronary chronic total occlusion in patients with viable myocardium. <i>Cardiovascular Ultrasound</i> , 2020, 18, 29. | 0.5 | 0 |
| 18 | Genetic determinants of clinical phenotype in hypertrophic cardiomyopathy. <i>BMC Cardiovascular Disorders</i> , 2020, 20, 516. | 0.7 | 33 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Impairment of coronary flow velocity reserve and global longitudinal strain in women with cardiac syndrome X and slow coronary flow. <i>Journal of Cardiology</i> , 2020, 76, 1-8. | 0.8 | 14 |
| 20 | Feasibility and value of two-dimensional volumetric stress echocardiography. <i>Minerva Cardiology and Angiology</i> , 2020, , . | 0.4 | 4 |
| 21 | Prognostic Value of Preserved Coronary Flow Velocity Reserve by Noninvasive Transthoracic Doppler Echocardiography in Patients With Angiographically Intermediate Left Main Stenosis. <i>Journal of the American Society of Echocardiography</i> , 2019, 32, 74-80. | 1.2 | 11 |
| 22 | Prognostic Value of Transthoracic Doppler Echocardiography Coronary Flow Velocity Reserve in Patients with Nonculprit Stenosis of Intermediate Severity Early after Primary Percutaneous Coronary Intervention. <i>Journal of the American Society of Echocardiography</i> , 2018, 31, 880-887. | 1.2 | 13 |
| 23 | Increased left ventricular mass index is present in patients with type 2 diabetes without ischemic heart disease. <i>Scientific Reports</i> , 2018, 8, 926. | 1.6 | 23 |
| 24 | Improved Propensity-Score Matched Long-Term Clinical Outcomes in Patients with Successful Percutaneous Coronary Interventions of Coronary Chronic Total Occlusion. <i>International Heart Journal</i> , 2018, 59, 719-726. | 0.5 | 13 |
| 25 | N-terminal pro-brain natriuretic peptide is related with coronary flow velocity reserve and diastolic dysfunction in patients with asymmetric hypertrophic cardiomyopathy. <i>Journal of Cardiology</i> , 2017, 70, 323-328. | 0.8 | 25 |
| 26 | Significance of relative coronary flow reserve in patient with microvascular dysfunction to differentiate significant coronary artery stenosis. <i>Srce I Krvni Sudovi</i> , 2017, 36, 102-104. | 0.1 | 0 |
| 27 | Left atrial appendage closure with Watchman device in prevention of thromboembolic complications in patients with atrial fibrillation: First experience in Serbia. <i>Vojnosanitetski Pregled</i> , 2017, 74, 378-385. | 0.1 | 0 |
| 28 | Prognostic role of stress echocardiography in hypertrophic cardiomyopathy: The International Stress Echo Registry. <i>International Journal of Cardiology</i> , 2016, 219, 331-338. | 0.8 | 38 |
| 29 | Manual versus target-controlled infusion of balanced propofol during diagnostic colonoscopy: A prospective randomized controlled trial. <i>Srpski Arhiv Za Celokupno Lekarstvo</i> , 2016, 144, 514-520. | 0.1 | 2 |
| 30 | Acute renal failure and hepatocellular damage as presenting symptoms of type II aortic dissection. <i>Srpski Arhiv Za Celokupno Lekarstvo</i> , 2016, 144, 320-324. | 0.1 | 0 |
| 31 | Percutaneous implantation of self-expandable aortic valve in high risk patients with severe aortic stenosis: The first experiences in Serbia. <i>Vojnosanitetski Pregled</i> , 2016, 73, 192-197. | 0.1 | 2 |
| 32 | The effects of nicorandil on microvascular function in patients with ST segment elevation myocardial infarction undergoing primary PCI. <i>Cardiovascular Ultrasound</i> , 2015, 13, 26. | 0.5 | 29 |
| 33 | Prognostic value of calcium score and coronary flow velocity reserve in asymptomatic diabetic patients. <i>Cardiovascular Ultrasound</i> , 2015, 13, 41. | 0.5 | 15 |
| 34 | Oxidized Low Density Lipoprotein and High Sensitive C-Reactive Protein in Non-Diabetic, Pre-Diabetic and Diabetic Patients in the Acute Phase of the First Myocardial Infarction Treated by Primary Percutaneous Coronary Intervention / Oksidovani Lipoprotein Niske Gustine I Visokosenzitivni C-Reaktivni Protein Kod Nedijabetičara, Predijabetičara I Dijabetičara U Akutnoj Fazi Prvog Infarkta Miokarda LeĀenog Primarnom Perkutanom Koronarom Intervencijom. <i>Journal of Medical Biochemistry</i> , 2015, 34, 160-169. | 0.7 | 5 |
| 35 | Time-dependent changes of plasma adiponectin concentration in relation to coronary microcirculatory function in patients with acute myocardial infarction treated by primary percutaneous coronary intervention. <i>Journal of Cardiology</i> , 2015, 65, 208-215. | 0.8 | 14 |
| 36 | Ophthalmic Manifestations in Children and Young Adults with Down Syndrome and Congenital Heart Defects. <i>Ophthalmic Epidemiology</i> , 2015, 22, 123-129. | 0.8 | 15 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Is there enough evidence for routine use of drug-eluting stents in acute myocardial infarction with ST segment elevation?. <i>Vojnosanitetski Pregled</i> , 2014, 71, 870-874. | 0.1 | 1 |
| 38 | The Randomized Physiologic Assessment of Thrombus Aspiration in Patients with Acute STâ€Segment Elevation Myocardial Infarction Trial (PATA STEMI): Study Rationale and Design. <i>Journal of Interventional Cardiology</i> , 2014, 27, 341-347. | 0.5 | 5 |
| 39 | Prognostic role of coronary flow reserve for left ventricular functional improvement after cardiac resynchronization therapy in patients with dilated cardiomyopathy. <i>European Heart Journal Cardiovascular Imaging</i> , 2014, 15, 1344-1349. | 0.5 | 16 |
| 40 | Coronary flow of the infarct artery assessed by transthoracic Doppler after primary percutaneous coronary intervention predicts final infarct size. <i>International Journal of Cardiovascular Imaging</i> , 2014, 30, 1509-1518. | 0.7 | 6 |
| 41 | The role of glycemia in acute heart failure patients. <i>Clinical Chemistry and Laboratory Medicine</i> , 2014, 52, 1437-46. | 1.4 | 4 |
| 42 | Coronary thrombi neovascularization in patients with ST-elevation myocardial infarction - clinical and angiographic implications. <i>Thrombosis Research</i> , 2014, 134, 1038-1045. | 0.8 | 4 |
| 43 | Structural myocardial alterations in diabetes and hypertension: the role of galectin-3. <i>Clinical Chemistry and Laboratory Medicine</i> , 2014, 52, 1499-505. | 1.4 | 16 |
| 44 | Acute insulin resistance in ST-segment elevation myocardial infarction in non-diabetic patients is associated with incomplete myocardial reperfusion and impaired coronary microcirculatory function. <i>Cardiovascular Diabetology</i> , 2014, 13, 73. | 2.7 | 37 |
| 45 | Regional Difference of Microcirculation in Patients withÂAsymmetric Hypertrophic Cardiomyopathy: Transthoracic Doppler Coronary Flow Velocity Reserve Analysis. <i>Journal of the American Society of Echocardiography</i> , 2013, 26, 775-782. | 1.2 | 26 |
| 46 | Two rare conditions in an Eisenmenger patient: Left main coronary artery compression and Ortner's syndrome due to pulmonary artery dilatation. <i>Heart and Lung: Journal of Acute and Critical Care</i> , 2013, 42, 382-386. | 0.8 | 11 |
| 47 | Estimation of infarct size using transthoracic Doppler echocardiographic measurement of coronary flow reserve in infarct related and reference coronary artery. <i>International Journal of Cardiology</i> , 2013, 168, 169-175. | 0.8 | 8 |
| 48 | Glycogen phosphorylase isoenzyme BB plasma kinetics is not related to myocardial ischemia induced by exercise stress echo test. <i>Clinical Chemistry and Laboratory Medicine</i> , 2013, 51, 2029-2035. | 1.4 | 3 |
| 49 | Predictors of diabetic cardiomyopathy in asymptomatic patients with type 2 diabetes. <i>International Journal of Cardiology</i> , 2012, 156, 219-221. | 0.8 | 11 |
| 50 | Prediction of a Good Response to Cardiac Resynchronization Therapy in Patients with Severe Dilated Cardiomyopathy: Could Conventional Echocardiography Be the Answer after All?. <i>Echocardiography</i> , 2012, 29, 267-275. | 0.3 | 5 |
| 51 | Prediction of Myocardial Functional Recovery by Noninvasive Evaluation of Basal and Hyperemic Coronary Flow in Patients with Previous Myocardial Infarction. <i>Journal of the American Society of Echocardiography</i> , 2011, 24, 573-581. | 1.2 | 15 |
| 52 | Echocardiography in sports cardiology: LV remodeling in athletes' heart â€” Questions to be answered. <i>Interventional Medicine & Applied Science</i> , 2011, 3, 120-123. | 0.2 | 0 |
| 53 | Asymptomatic cardiovascular manifestations in diabetes mellitus: Left ventricular diastolic dysfunction and silent myocardial ischemia. <i>Srpski Arhiv Za Celokupno Lekarstvo</i> , 2011, 139, 599-604. | 0.1 | 4 |
| 54 | The Use of Intracoronary Sodium Nitroprusside to Treat No-Reflow after Primary Percutaneous Coronary Intervention in Acute Myocardial Infarction. <i>Herz</i> , 2010, 35, 114-118. | 0.4 | 9 |