

Ji-Eun Lee

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4200261/publications.pdf>

Version: 2024-02-01

41
papers

6,799
citations

172207

29
h-index

264894

42
g-index

47
all docs

47
docs citations

47
times ranked

13517
citing authors

#	ARTICLE	IF	CITATIONS
1	Interplay of BAF and MLL4 promotes cell type-specific enhancer activation. <i>Nature Communications</i> , 2021, 12, 1630.	5.8	38
2	MED1 is a lipogenesis coactivator required for postnatal adipose expansion. <i>Genes and Development</i> , 2021, 35, 713-728.	2.7	9
3	Refining cell-based assay to detect MOG-IgG in patients with central nervous system inflammatory diseases. <i>Multiple Sclerosis and Related Disorders</i> , 2020, 40, 101939.	0.9	24
4	Loss of Function of the Gene Encoding the Histone Methyltransferase KMT2D Leads to Deregulation of Mitochondrial Respiration. <i>Cells</i> , 2020, 9, 1685.	1.8	10
5	Molecular basis for histone H3 H3K4me3-K9me3/2H3K4me3-methylation pattern readout by Spindlin1. <i>Journal of Biological Chemistry</i> , 2020, 295, 16877-16887.	1.6	15
6	MLL3/MLL4-Associated PAGR1 Regulates Adipogenesis by Controlling Induction of C/EBP β and C/EBP δ . <i>Molecular and Cellular Biology</i> , 2020, 40, .	1.1	15
7	A Mouse Homolog of a Human TP53 Germline Mutation Reveals a Lipolytic Activity of p53. <i>Cell Reports</i> , 2020, 30, 783-792.e5.	2.9	12
8	Opposing Functions of BRD4 Isoforms in Breast Cancer. <i>Molecular Cell</i> , 2020, 78, 1114-1132.e10.	4.5	95
9	Histone methyltransferase MLL4 controls myofiber identity and muscle performance through MEF2 interaction. <i>Journal of Clinical Investigation</i> , 2020, 130, 4710-4725.	3.9	24
10	Selective binding of the PHD6 finger of MLL4 to histone H4K16ac links MLL4 and MOF. <i>Nature Communications</i> , 2019, 10, 2314.	5.8	40
11	Transcriptional and Epigenomic Regulation of Adipogenesis. <i>Molecular and Cellular Biology</i> , 2019, 39, .	1.1	178
12	H3.3K4M destabilizes enhancer H3K4 methyltransferases MLL3/MLL4 and impairs adipose tissue development. <i>Nucleic Acids Research</i> , 2019, 47, 607-620.	6.5	1,326
13	Depletion of Nsd2-mediated histone H3K36 methylation impairs adipose tissue development and function. <i>Nature Communications</i> , 2018, 9, 1796.	5.8	58
14	MLL3/MLL4 are required for CBP/p300 binding on enhancers and super-enhancer formation in brown adipogenesis. <i>Nucleic Acids Research</i> , 2017, 45, 6388-6403.	6.5	131
15	MLL4 prepares the enhancer landscape for Foxp3 induction via chromatin looping. <i>Nature Immunology</i> , 2017, 18, 1035-1045.	7.0	63
16	Distinct Roles of Transcription Factors KLF4, Krox20, and Peroxisome Proliferator-Activated Receptor γ in Adipogenesis. <i>Molecular and Cellular Biology</i> , 2017, 37, .	1.1	44
17	Brd4 binds to active enhancers to control cell identity gene induction in adipogenesis and myogenesis. <i>Nature Communications</i> , 2017, 8, 2217.	5.8	161
18	Enhancer priming by H3K4 methyltransferase MLL4 controls cell fate transition. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016, 113, 11871-11876.	3.3	172

#	ARTICLE	IF	CITATIONS
19	Replication fork stability confers chemoresistance in BRCA-deficient cells. <i>Nature</i> , 2016, 535, 382-387.	13.7	685
20	KMT2D regulates specific programs in heart development via histone H3 lysine 4 di-methylation. <i>Development (Cambridge)</i> , 2016, 143, 810-821.	1.2	100
21	A PTIP-PA1 subcomplex promotes transcription for IgH class switching independently from the associated MLL3/MLL4 methyltransferase complex. <i>Genes and Development</i> , 2016, 30, 149-163.	2.7	27
22	Disruption of KMT2D perturbs germinal center B cell development and promotes lymphomagenesis. <i>Nature Medicine</i> , 2015, 21, 1190-1198.	15.2	372
23	The histone lysine methyltransferase KMT2D sustains a gene expression program that represses B cell lymphoma development. <i>Nature Medicine</i> , 2015, 21, 1199-1208.	15.2	359
24	ATP-citrate lyase regulates cellular senescence via an AMPK- and p53-dependent pathway. <i>FEBS Journal</i> , 2015, 282, 361-371.	2.2	53
25	Loss of function of mouse Pax-interacting Protein 1-associated glutamate rich protein 1a (<i>Pagr1a</i>) leads to reduced <i>Bmp2</i> expression and defects in chorion and amnion development. <i>Developmental Dynamics</i> , 2014, 243, 937-947.	0.8	19
26	DNA-damage-induced differentiation of leukaemic cells as an anti-cancer barrier. <i>Nature</i> , 2014, 514, 107-111.	13.7	174
27	Transcriptional and epigenetic regulation of PPAR β expression during adipogenesis. <i>Cell and Bioscience</i> , 2014, 4, 29.	2.1	182
28	A Multifunctional Protein, EWS, Is Essential for Early Brown Fat Lineage Determination. <i>Developmental Cell</i> , 2013, 26, 393-404.	3.1	70
29	p53 regulates glucose metabolism by miR-34a. <i>Biochemical and Biophysical Research Communications</i> , 2013, 437, 225-231.	1.0	90
30	53BP1 Mediates Productive and Mutagenic DNA Repair through Distinct Phosphoprotein Interactions. <i>Cell</i> , 2013, 153, 1266-1280.	13.5	292
31	H3K4 mono- and di-methyltransferase MLL4 is required for enhancer activation during cell differentiation. <i>ELife</i> , 2013, 2, e01503.	2.8	369
32	UTX regulates mesoderm differentiation of embryonic stem cells independent of H3K27 demethylase activity. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012, 109, 15324-15329.	3.3	183
33	A p53-inducible microRNA-34a downregulates Ras signaling by targeting IMPDH. <i>Biochemical and Biophysical Research Communications</i> , 2012, 418, 682-688.	1.0	34
34	Histone H3K9 methyltransferase G9a represses PPAR β expression and adipogenesis. <i>EMBO Journal</i> , 2012, 32, 45-59.	3.5	162
35	Distinct roles of GCN5/PCAF-mediated H3K9ac and CBP/p300-mediated H3K18/27ac in nuclear receptor transactivation. <i>EMBO Journal</i> , 2011, 30, 249-262.	3.5	655
36	Histone H3K27 methyltransferase Ezh2 represses <i>Wnt</i> genes to facilitate adipogenesis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010, 107, 7317-7322.	3.3	258

#	ARTICLE	IF	CITATIONS
37	Down syndrome critical region 1 enhances the proteolytic cleavage of calcineurin. <i>Experimental and Molecular Medicine</i> , 2009, 41, 471.	3.2	13
38	Calcineurin dephosphorylates glycogen synthase kinase-3 beta at serine-9 in neuroblastâ-derived cells. <i>Journal of Neurochemistry</i> , 2009, 111, 344-354.	2.1	62
39	Histone Methylation Regulator PTIP Is Required for PPAR β and C/EBP β Expression and Adipogenesis. <i>Cell Metabolism</i> , 2009, 10, 27-39.	7.2	117
40	Hydrogen peroxide triggers the proteolytic cleavage and the inactivation of calcineurin. <i>Journal of Neurochemistry</i> , 2007, 100, 070209222715097-???	2.1	21
41	Menin represses JunD transcriptional activity in protein kinase C δ -mediated Nur77 expression. <i>Experimental and Molecular Medicine</i> , 2005, 37, 466-475.	3.2	37