

Rania G Abdel-Latif

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4201228/publications.pdf>

Version: 2024-02-01

13
papers

326
citations

932766

10
h-index

1125271

13
g-index

13
all docs

13
docs citations

13
times ranked

404
citing authors

#	ARTICLE	IF	CITATIONS
1	Preparation and Evaluation of Atorvastatin-Loaded Nanoemulgel on Wound-Healing Efficacy. <i>Pharmaceutics</i> , 2019, 11, 609.	2.0	67
2	Empagliflozin attenuates transient cerebral ischemia/reperfusion injury in hyperglycemic rats via repressing oxidative-inflammatory-apoptotic pathway. <i>Fundamental and Clinical Pharmacology</i> , 2020, 34, 548-558.	1.0	59
3	Empagliflozin alleviates neuronal apoptosis induced by cerebral ischemia/reperfusion injury through HIF-1 α /VEGF signaling pathway. <i>Archives of Pharmacol Research</i> , 2020, 43, 514-525.	2.7	47
4	Nephroprotective effect of exogenous hydrogen sulfide donor against cyclophosphamide-induced toxicity is mediated by Nrf2/HO-1/NF- κ B signaling pathway. <i>Life Sciences</i> , 2021, 264, 118630.	2.0	28
5	Sildenafil protects against nitric oxide deficiency-related nephrotoxicity in cyclosporine A treated rats. <i>European Journal of Pharmacology</i> , 2013, 705, 126-134.	1.7	26
6	Lixisenatide, a novel GLP-1 analog, protects against cerebral ischemia/reperfusion injury in diabetic rats. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 2018, 391, 705-717.	1.4	25
7	TLRs-JNK/ NF- κ B Pathway Underlies the Protective Effect of the Sulfide Salt Against Liver Toxicity. <i>Frontiers in Pharmacology</i> , 2022, 13, 850066.	1.6	23
8	Lixisenatide ameliorates cerebral ischemia-reperfusion injury via GLP-1 receptor dependent/independent pathways. <i>European Journal of Pharmacology</i> , 2018, 833, 145-154.	1.7	13
9	Modulation of eNOS/iNOS by nebivolol protects against cyclosporine A-mediated nephrotoxicity through targeting inflammatory and apoptotic pathways. <i>Environmental Toxicology and Pharmacology</i> , 2019, 69, 26-35.	2.0	12
10	Low-dose lixisenatide protects against early-onset nephropathy induced in diabetic rats. <i>Life Sciences</i> , 2020, 263, 118592.	2.0	12
11	Comparative effectiveness of phosphodiesterase 3, 4, and 5 inhibitors in amelioration of high-fat diet-induced nonalcoholic fatty liver in rats. <i>Fundamental and Clinical Pharmacology</i> , 2020, 34, 353-364.	1.0	6
12	Statin therapy and SAR-COV-2: an available and potential therapy?. <i>European Heart Journal - Cardiovascular Pharmacotherapy</i> , 2020, 6, 333-334.	1.4	5
13	Combined treatments with metformin and phosphodiesterase inhibitors alleviate nonalcoholic fatty liver disease in high-fat diet fed rats: a comparative study. <i>Canadian Journal of Physiology and Pharmacology</i> , 2020, 98, 498-505.	0.7	3