

Harry S Paris

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4209669/publications.pdf>

Version: 2024-02-01

33
papers

1,069
citations

394421

19
h-index

434195

31
g-index

33
all docs

33
docs citations

33
times ranked

841
citing authors

#	ARTICLE	IF	CITATIONS
1	The multiple-flowering trait conferred by gene <i>mf</i> increases yield of field-grown Cocomelle and Zucchini squash. <i>Euphytica</i> , 2022, 218, 1.	1.2	0
2	Chilling sensitivity of four near-isogenic fruit-color genotypes of summer squash (<i>Cucurbita pepo</i>), Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 2020, 168, 111279.	6.0	7
3	Dark-stem-dependent reverse fruit striping in <i>Cucurbita pepo</i> (pumpkin, squash, gourd: Cucurbitaceae): Genes <i>l-2</i> and <i>W</i> are at the same locus. <i>Euphytica</i> , 2020, 216, 1.	1.2	1
4	Whole-genome resequencing of <i>Cucurbita pepo</i> morphotypes to discover genomic variants associated with morphology and horticulturally valuable traits. <i>Horticulture Research</i> , 2019, 6, 94.	6.3	34
5	Italian horticultural and culinary records of summer squash (<i>Cucurbita pepo</i> , Cucurbitaceae) and emergence of the zucchini in 19th-century Milan. <i>Annals of Botany</i> , 2016, 118, 53-69.	2.9	23
6	Germplasm enhancement of <i>Cucurbita pepo</i> (pumpkin, squash, gourd: Cucurbitaceae): progress and challenges. <i>Euphytica</i> , 2016, 208, 415-438.	1.2	45
7	Overview of the origins and history of the five major cucurbit crops: issues for ancient DNA analysis of archaeological specimens. <i>Vegetation History and Archaeobotany</i> , 2016, 25, 405-414.	2.1	45
8	Origin and emergence of the sweet dessert watermelon, <i>Citrullus lanatus</i> . <i>Annals of Botany</i> , 2015, 116, 133-148.	2.9	125
9	Cucurbits depicted in Byzantine mosaics from Israel, 350â€“600 ce. <i>Annals of Botany</i> , 2014, 114, 203-222.	2.9	6
10	Another gene affecting fruit and stem color in squash, <i>Cucurbita pepo</i> . <i>Euphytica</i> , 2013, 191, 99-107.	1.2	5
11	Medieval iconography of watermelons in Mediterranean Europe. <i>Annals of Botany</i> , 2013, 112, 867-879.	2.9	24
12	Medieval emergence of sweet melons, <i>Cucumis melo</i> (Cucurbitaceae). <i>Annals of Botany</i> , 2012, 110, 23-33.	2.9	56
13	Occidental diffusion of cucumber (<i>Cucumis sativus</i>) 500â€“1300 CE: two routes to Europe. <i>Annals of Botany</i> , 2012, 109, 117-126.	2.9	42
14	Medieval History of the Dudaâ€™im Melon (<i>Cucumis melo</i> , Cucurbitaceae). <i>Economic Botany</i> , 2012, 66, 276-284.	1.7	9
15	Parallel Evolution Under Domestication and Phenotypic Differentiation of the Cultivated Subspecies of <i>Cucurbita pepo</i> (Cucurbitaceae). <i>Economic Botany</i> , 2012, 66, 71-90.	1.7	36
16	Genetic relationships and evolution in <i>Cucurbita pepo</i> (pumpkin, squash, gourd) as revealed by simple sequence repeat polymorphisms. <i>Theoretical and Applied Genetics</i> , 2012, 124, 875-891.	3.6	72
17	Medieval herbal iconography and lexicography of <i>Cucumis</i> (cucumber and melon, Cucurbitaceae) in the Occident, 1300â€“1458. <i>Annals of Botany</i> , 2011, 108, 471-484.	2.9	29
18	Single Recessive Gene for Multiple Flowering in Summer Squash. <i>Hortscience: A Publication of the American Society for Horticultural Science</i> , 2010, 45, 1643-1644.	1.0	6

#	ARTICLE	IF	CITATIONS
19	The Cucurbitaceae and Solanaceae illustrated in medieval manuscripts known as the Tacuinum Sanitatis. <i>Annals of Botany</i> , 2009, 103, 1187-1205.	2.9	26
20	Genes for "Reverse" Fruit Striping in Squash (<i>Cucurbita pepo</i>). <i>Journal of Heredity</i> , 2009, 100, 371-379.	2.4	16
21	Summer Squash. , 2008, , 351-379.		23
22	The Cucurbits of Mediterranean Antiquity: Identification of Taxa from Ancient Images and Descriptions. <i>Annals of Botany</i> , 2007, 100, 1441-1457.	2.9	100
23	First Known Image of <i>Cucurbita</i> in Europe, 1503â€“1508. <i>Annals of Botany</i> , 2006, 98, 41-47.	2.9	45
24	The Cucurbit Images (1515â€“1518) of the Villa Farnesina, Rome. <i>Annals of Botany</i> , 2006, 97, 165-176.	2.9	42
25	Dual Role of the Pigmentation Gene B in Affecting Carotenoid and Vitamin E Content in Squash (<i>Cucurbita pepo</i>) Mesocarp. <i>Journal of Agricultural and Food Chemistry</i> , 2005, 53, 9759-9763.	5.2	31
26	The Genes of Pumpkin and Squash. <i>Hortscience: A Publication of the American Society for Horticultural Science</i> , 2005, 40, 1620-1630.	1.0	85
27	Title is missing!. <i>Euphytica</i> , 2002, 124, 121-128.	1.2	15
28	Multiple allelism at a major locus affecting fruit coloration in <i>Cucurbita pepo</i> . <i>Euphytica</i> , 2002, 125, 149-153.	1.2	9
29	Gene for broad, contiguous dark stripes in cocozelle squash (<i>Cucurbita pepo</i>). <i>Euphytica</i> , 2000, 115, 191-196.	1.2	12
30	Paintings (1769â€“1774) by A. N. Duchesne and the History of <i>Cucurbita pepo</i> . <i>Annals of Botany</i> , 2000, 85, 815-830.	2.9	19
31	First two publications by Duchesne of <i>Cucurbita moschata</i> (Cucurbitaceae). <i>Taxon</i> , 2000, 49, 305-319.	0.7	10
32	Summer Squash: History, Diversity, and Distribution. <i>HortTechnology</i> , 1996, 6, 6-13.	0.9	34
33	Genes for intense fruit pigmentation of squash. <i>Journal of Heredity</i> , 1986, 77, 403-409.	2.4	37