

Patricia A Labosky

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/421343/publications.pdf>

Version: 2024-02-01

50
papers

5,727
citations

145106

33
h-index

223390

49
g-index

51
all docs

51
docs citations

51
times ranked

8329
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | The NIH "BEST" programs: Institutional programs, the program evaluation, and early data. FASEB Journal, 2020, 34, 3570-3582. | 0.2 | 39 |
| 2 | Evolution of a functional taxonomy of career pathways for biomedical trainees. Journal of Clinical and Translational Science, 2018, 2, 63-65. | 0.3 | 16 |
| 3 | FOXD3 Regulates Pluripotent Stem Cell Potential by Simultaneously Initiating and Repressing Enhancer Activity. Cell Stem Cell, 2016, 18, 104-117. | 5.2 | 93 |
| 4 | The origin and implementation of the Broadening Experiences in Scientific Training programs: an NIH common fund initiative. FASEB Journal, 2016, 30, 507-514. | 0.2 | 71 |
| 5 | The NIH Extracellular RNA Communication Consortium. Journal of Extracellular Vesicles, 2015, 4, 27493. | 5.5 | 60 |
| 6 | Vascular endothelial growth factor coordinates islet innervation via vascular scaffolding. Development (Cambridge), 2014, 141, 1480-1491. | 1.2 | 77 |
| 7 | Transcriptional targets of Foxd3 in murine ES cells. Stem Cell Research, 2014, 12, 233-240. | 0.3 | 24 |
| 8 | Embryonic domains of the aorta derived from diverse origins exhibit distinct properties that converge into a common phenotype in the adult. Journal of Molecular and Cellular Cardiology, 2014, 69, 88-96. | 0.9 | 49 |
| 9 | Ground-State Transcriptional Requirements for Skin-Derived Precursors. Stem Cells and Development, 2013, 22, 1779-1788. | 1.1 | 7 |
| 10 | A dynamic code of dorsal neural tube genes regulates the segregation between neurogenic and melanogenic neural crest cells. Development (Cambridge), 2013, 140, 2269-2279. | 1.2 | 77 |
| 11 | Neural crest and Schwann cell progenitor-derived melanocytes are two spatially segregated populations similarly regulated by Foxd3. Proceedings of the National Academy of Sciences of the United States of America, 2013, 110, 12709-12714. | 3.3 | 92 |
| 12 | Transgenic Mouse Models. Methods in Molecular Biology, 2013, 1032, 1-17. | 0.4 | 3 |
| 13 | Renshaw cell interneuron specialization is controlled by a temporally restricted transcription factor program. Development (Cambridge), 2012, 139, 179-190. | 1.2 | 96 |
| 14 | Isolation and Culture of Neural Crest Cells from Embryonic Murine Neural Tube. Journal of Visualized Experiments, 2012, , e4134. | 0.2 | 19 |
| 15 | Enteric nervous system specific deletion of Foxd3 disrupts glial cell differentiation and activates compensatory enteric progenitors. Developmental Biology, 2012, 363, 373-387. | 0.9 | 38 |
| 16 | Neural crest stem cell multipotency requires Foxd3 to maintain neural potential and repress mesenchymal fates. Development (Cambridge), 2011, 138, 641-652. | 1.2 | 97 |
| 17 | Influence and timing of arrival of murine neural crest on pancreatic beta cell development and maturation. Developmental Biology, 2011, 349, 321-330. | 0.9 | 41 |
| 18 | A predicted hairpin cluster correlates with barriers to PCR, sequencing and possibly BAC recombineering. Scientific Reports, 2011, 1, 106. | 1.6 | 21 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Functional interaction between Foxd3 and Pax3 in cardiac neural crest development. <i>Genesis</i> , 2011, 49, 10-23. | 0.8 | 24 |
| 20 | Loss of Foxd3 Results in Decreased β -Cell Proliferation and Glucose Intolerance During Pregnancy. <i>Endocrinology</i> , 2011, 152, 4589-4600. | 1.4 | 30 |
| 21 | Conditional Gene Targeting in Mouse Pancreatic β -Cells. <i>Diabetes</i> , 2010, 59, 3090-3098. | 0.3 | 288 |
| 22 | Transcriptional Control of Neural Crest Development. <i>Colloquium Series on Developmental Biology</i> , 2010, 1, 1-227. | 0.2 | 30 |
| 23 | Epigenetic Priming of a Pre-B Cell-Specific Enhancer through Binding of Sox2 and Foxd3 at the ESC Stage. <i>Cell Stem Cell</i> , 2010, 7, 114-126. | 5.2 | 79 |
| 24 | Endocardial cells are a distinct endothelial lineage derived from Flk1+ multipotent cardiovascular progenitors. <i>Developmental Biology</i> , 2009, 333, 78-89. | 0.9 | 106 |
| 25 | Targeted loss of Arx results in a developmental epilepsy mouse model and recapitulates the human phenotype in heterozygous females. <i>Brain</i> , 2009, 132, 1563-1576. | 3.7 | 178 |
| 26 | Regulation of Embryonic Stem Cell Self-Renewal and Pluripotency by Foxd3. <i>Stem Cells</i> , 2008, 26, 2475-2484. | 1.4 | 93 |
| 27 | Requirement for Foxd3 in the maintenance of neural crest progenitors. <i>Development (Cambridge)</i> , 2008, 135, 1615-1624. | 1.2 | 171 |
| 28 | Identification of Arx transcriptional targets in the developing basal forebrain. <i>Human Molecular Genetics</i> , 2008, 17, 3740-3760. | 1.4 | 121 |
| 29 | Mouse Primordial Germ Cells. <i>Methods in Molecular Biology</i> , 2008, 461, 187-199. | 0.4 | 4 |
| 30 | SULF1 and SULF2 regulate heparan sulfate-mediated GDNF signaling for esophageal innervation. <i>Development (Cambridge)</i> , 2007, 134, 3327-3338. | 1.2 | 148 |
| 31 | Resident Endothelial Precursors in Muscle, Adipose, and Dermis Contribute to Postnatal Vasculogenesis. <i>Stem Cells</i> , 2007, 25, 3101-3110. | 1.4 | 77 |
| 32 | Expression and shifting subcellular localization of the transcription factor, Foxd3, in embryonic and adult pancreas. <i>Gene Expression Patterns</i> , 2006, 6, 971-977. | 0.3 | 27 |
| 33 | FoxD3 regulation of Nodal in the Spemann organizer is essential for <i>Xenopus</i> dorsal mesoderm development. <i>Development (Cambridge)</i> , 2006, 133, 4827-4838. | 1.2 | 44 |
| 34 | HIF-2 α regulates Oct-4: effects of hypoxia on stem cell function, embryonic development, and tumor growth. <i>Genes and Development</i> , 2006, 20, 557-570. | 2.7 | 721 |
| 35 | Neural Crest Stem Cells. , 2006, 589, 206-212. | | 33 |
| 36 | Foxd3 is required in the trophoblast progenitor cell lineage of the mouse embryo. <i>Developmental Biology</i> , 2005, 285, 126-137. | 0.9 | 90 |

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 37 | A dermal niche for multipotent adult skin-derived precursor cells. <i>Nature Cell Biology</i> , 2004, 6, 1082-1093. | 4.6 | 692 |
| 38 | Electroporation of Murine Embryonic Stem Cells: A Step-by-Step Guide. <i>Stem Cells</i> , 2004, 22, 243-249. | 1.4 | 51 |
| 39 | Online with Stem Cells. <i>Developmental Cell</i> , 2003, 5, 827-828. | 3.1 | 0 |
| 40 | Requirement for Foxd3 in maintaining pluripotent cells of the early mouse embryo. <i>Genes and Development</i> , 2002, 16, 2650-2661. | 2.7 | 323 |
| 41 | The winged helix gene, Foxb1, controls development of mammary glands and regions of the CNS that regulate the milk-ejection reflex. <i>Genesis</i> , 2001, 29, 60-71. | 0.8 | 16 |
| 42 | The winged-helix transcription factor Foxd3 suppresses interneuron differentiation and promotes neural crest cell fate. <i>Development (Cambridge)</i> , 2001, 128, 4127-4138. | 1.2 | 247 |
| 43 | Mesendoderm Induction and Reversal of Left-Right Pattern by Mouse Gdf1, a Vg1-Related Gene. <i>Developmental Biology</i> , 2000, 227, 495-509. | 0.9 | 65 |
| 44 | The winged helix transcription factor Hfh2 is expressed in neural crest and spinal cord during mouse development. <i>Mechanisms of Development</i> , 1998, 76, 185-190. | 1.7 | 102 |
| 45 | A Role for Indian hedgehog in Extraembryonic Endoderm Differentiation in F9 Cells and the Early Mouse Embryo. <i>Developmental Biology</i> , 1997, 187, 298-310. | 0.9 | 53 |
| 46 | The Chromosomal Mapping of Four Genes Encoding Winged Helix Proteins Expressed Early in Mouse Development. <i>Genomics</i> , 1996, 34, 241-245. | 1.3 | 13 |
| 47 | Effects on blood pressure and exploratory behaviour of mice lacking angiotensin II type-2 receptor. <i>Nature</i> , 1995, 377, 748-750. | 13.7 | 821 |
| 48 | The Pbx Homeobox Gene Is X-Linked and Exclusively Expressed in Extraembryonic Tissues during Early Murine Development. <i>Developmental Biology</i> , 1994, 166, 170-179. | 0.9 | 106 |
| 49 | Embryonic Germ Cell Lines and Their Derivation from Mouse Primordial Germ Cells. <i>Novartis Foundation Symposium</i> , 1994, 182, 157-178. | 1.2 | 40 |
| 50 | Homeobox-Containing Genes in Teratocarcinoma Embryoid Bodies: A Possible Role for Hox-D12 (Hox-4.7) in Establishing the Extraembryonic Endoderm Lineage in the Mouse. <i>Developmental Biology</i> , 1993, 159, 232-244. | 0.9 | 8 |