

Martin Turner

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4216272/publications.pdf>

Version: 2024-02-01

185
papers

19,119
citations

16791

66
h-index

13635

134
g-index

201
all docs

201
docs citations

201
times ranked

20936
citing authors

#	ARTICLE	IF	CITATIONS
1	Correction to: ORFLine: a bioinformatic pipeline to prioritize small open reading frames identifies candidate secreted small proteins from lymphocytes. <i>Bioinformatics</i> , 2022, 38, 2673-2673.	1.8	1
2	Polypyrimidine tract binding protein 1 regulates the activation of mouse CD8 T cells. <i>European Journal of Immunology</i> , 2022, 52, 1058-1068.	1.6	5
3	A functional screen of RNA binding proteins identifies genes that promote or limit the accumulation of CD138+ plasma cells. <i>ELife</i> , 2022, 11, .	2.8	5
4	The timing of differentiation and potency of CD8 effector function is set by RNA binding proteins. <i>Nature Communications</i> , 2022, 13, 2274.	5.8	25
5	ORFLine: a bioinformatic pipeline to prioritize small open reading frames identifies candidate secreted small proteins from lymphocytes. <i>Bioinformatics</i> , 2021, 37, 3152-3159.	1.8	7
6	RNA Binding Proteins As Regulators of Oxidative Stress Identified by a Targeted CRISPR-Cas9 Single Guide RNA Library. <i>CRISPR Journal</i> , 2021, 4, 427-437.	1.4	8
7	Essential requirement for polypyrimidine tract binding proteins 1 and 3 in the maturation and maintenance of mature B cells in mice. <i>European Journal of Immunology</i> , 2021, 51, 2266-2273.	1.6	5
8	Sequential inverse dysregulation of the RNA helicases DDX3X and DDX3Y facilitates MYC-driven lymphomagenesis. <i>Molecular Cell</i> , 2021, 81, 4059-4075.e11.	4.5	42
9	A RAC-GEF network critical for early intestinal tumourigenesis. <i>Nature Communications</i> , 2021, 12, 56.	5.8	11
10	Efficient homing of antibody-secreting cells to the bone marrow requires RNA-binding protein ZFP36L1. <i>Journal of Experimental Medicine</i> , 2021, 218, .	4.2	19
11	The RNA-binding protein HuR is required for maintenance of the germinal centre response. <i>Nature Communications</i> , 2021, 12, 6556.	5.8	10
12	Dynamic Post-Transcriptional Events Governing CD8+ T Cell Homeostasis and Effector Function. <i>Trends in Immunology</i> , 2020, 41, 240-254.	2.9	39
13	Polypyrimidine tract-binding proteins are essential for B cell development. <i>ELife</i> , 2020, 9, .	2.8	25
14	Myeloid Tribbles 1 induces early atherosclerosis via enhanced foam cell expansion. <i>Science Advances</i> , 2019, 5, eaax9183.	4.7	50
15	Alternative Translation Initiation Generates a Functionally Distinct Isoform of the Stress-Activated Protein Kinase MK2. <i>Cell Reports</i> , 2019, 27, 2859-2870.e6.	2.9	22
16	Signalling circuits that direct early B-cell development. <i>Biochemical Journal</i> , 2019, 476, 769-778.	1.7	13
17	RNA-binding proteins in hematopoiesis and hematological malignancy. <i>Blood</i> , 2019, 133, 2365-2373.	0.6	52
18	Membrane Cholesterol Efflux Drives Tumor-Associated Macrophage Reprogramming and Tumor Progression. <i>Cell Metabolism</i> , 2019, 29, 1376-1389.e4.	7.2	261

#	ARTICLE	IF	CITATIONS
19	MicroRNA-155 is essential for the optimal proliferation and survival of plasmablast B cells. <i>Life Science Alliance</i> , 2019, 2, e201800244.	1.3	17
20	RNA-binding proteins control gene expression and cell fate in the immune system. <i>Nature Immunology</i> , 2018, 19, 120-129.	7.0	168
21	The RNA-binding protein PTBP1 is necessary for B cell selection in germinal centers. <i>Nature Immunology</i> , 2018, 19, 267-278.	7.0	63
22	The RNA-binding proteins Zfp361l1 and Zfp361l2 act redundantly in myogenesis. <i>Skeletal Muscle</i> , 2018, 8, 37.	1.9	22
23	Initial Identification of a Blood-Based Chromosome Conformation Signature for Aiding in the Diagnosis of Amyotrophic Lateral Sclerosis. <i>EBioMedicine</i> , 2018, 33, 169-184.	2.7	17
24	Uncovering the Role of RNA-Binding Proteins in Gene Expression in the Immune System. <i>Frontiers in Immunology</i> , 2018, 9, 1094.	2.2	60
25	Translational repression of pre-formed cytokine-encoding mRNA prevents chronic activation of memory T cells. <i>Nature Immunology</i> , 2018, 19, 828-837.	7.0	90
26	Antigen phagocytosis by B cells is required for a potent humoral response. <i>EMBO Reports</i> , 2018, 19, .	2.0	44
27	Transcriptome Analysis of Infected and Bystander Type 2 Alveolar Epithelial Cells during Influenza A Virus Infection Reveals <i>In Vivo</i> Wnt Pathway Downregulation. <i>Journal of Virology</i> , 2018, 92, .	1.5	50
28	RNA-binding proteins mind the GAPs. <i>Nature Immunology</i> , 2017, 18, 146-148.	7.0	2
29	Cell cycle <i>RNA</i> regulons coordinating early lymphocyte development. <i>Wiley Interdisciplinary Reviews RNA</i> , 2017, 8, e1419.	3.2	11
30	Maintenance of the marginal-zone B cell compartment specifically requires the RNA-binding protein ZFP36L1. <i>Nature Immunology</i> , 2017, 18, 683-693.	7.0	59
31	Tia1 dependent regulation of mRNA subcellular location and translation controls p53 expression in B cells. <i>Nature Communications</i> , 2017, 8, 530.	5.8	48
32	Characterization of the B Cell Transcriptome Bound by RNA-Binding Proteins with iCLIP. <i>Methods in Molecular Biology</i> , 2017, 1623, 159-179.	0.4	5
33	RNA-binding protein ZFP36L1 maintains posttranscriptional regulation of bile acid metabolism. <i>Journal of Clinical Investigation</i> , 2017, 127, 3741-3754.	3.9	45
34	RNA-binding proteins ZFP36L1 and ZFP36L2 promote cell quiescence. <i>Science</i> , 2016, 352, 453-459.	6.0	142
35	The RNA-Binding Proteins Zfp361l1 and Zfp361l2 Enforce the Thymic β^2 -Selection Checkpoint by Limiting DNA Damage Response Signaling and Cell Cycle Progression. <i>Journal of Immunology</i> , 2016, 197, 2673-2685.	0.4	63
36	The RNA-binding protein TTP is a global post-transcriptional regulator of feedback control in inflammation. <i>Nucleic Acids Research</i> , 2016, 44, gkw474.	6.5	128

#	ARTICLE	IF	CITATIONS
37	Anaplastic large cell lymphoma arises in thymocytes and requires transient TCR expression for thymic egress. <i>Nature Communications</i> , 2016, 7, 10087.	5.8	65
38	GIMAP1 Is Essential for the Survival of Naive and Activated B Cells In Vivo. <i>Journal of Immunology</i> , 2016, 196, 207-216.	0.4	26
39	Deletion of AU-Rich Elements within the Bcl2 3'UTR Reduces Protein Expression and B Cell Survival In Vivo. <i>PLoS ONE</i> , 2015, 10, e0116899.	1.1	11
40	Generation of functionally distinct isoforms of PTBP3 by alternative splicing and translation initiation. <i>Nucleic Acids Research</i> , 2015, 43, 5586-5600.	6.5	37
41	Improving access to medicines: empowering patients in the quest to improve treatment for rare lethal diseases. <i>Journal of Medical Ethics</i> , 2015, 41, 987-989.	1.0	8
42	The RNA-binding protein HuR is essential for the B cell antibody response. <i>Nature Immunology</i> , 2015, 16, 415-425.	7.0	125
43	The Role of p110 δ in the Development and Activation of B Lymphocytes. <i>Advances in Experimental Medicine and Biology</i> , 2015, 850, 119-135.	0.8	4
44	RNA binding proteins as regulators of immune cell biology. <i>Clinical and Experimental Immunology</i> , 2015, 183, 37-49.	1.1	50
45	MicroRNA-155 controls affinity-based selection by protecting c-MYC+ B cells from apoptosis. <i>Journal of Clinical Investigation</i> , 2015, 126, 377-388.	3.9	41
46	Generation and Characterisation of Mice Deficient in the Multi-GTPase Domain Containing Protein, GIMAP8. <i>PLoS ONE</i> , 2014, 9, e110294.	1.1	11
47	PI3K Signaling in B Cell and T Cell Biology. <i>Frontiers in Immunology</i> , 2014, 5, 557.	2.2	22
48	Noncoding RNA and its associated proteins as regulatory elements of the immune system. <i>Nature Immunology</i> , 2014, 15, 484-491.	7.0	165
49	The miR-155 \leftrightarrow PU.1 axis acts on Pax5 to enable efficient terminal B cell differentiation. <i>Journal of Experimental Medicine</i> , 2014, 211, 2183-2198.	4.2	83
50	Inactivation of PI(3)K p110 δ breaks regulatory T-cell-mediated immune tolerance to cancer. <i>Nature</i> , 2014, 510, 407-411.	13.7	450
51	The microRNA miR-155 controls CD8+ T cell responses by regulating interferon signaling. <i>Nature Immunology</i> , 2013, 14, 593-602.	7.0	249
52	Pharmacological Inhibition of Glycogen Synthase Kinase 3 Regulates T Cell Development In Vitro. <i>PLoS ONE</i> , 2013, 8, e58501.	1.1	15
53	RNA-binding proteins as a point of convergence of the PI3K and p38 MAPK pathways. <i>Frontiers in Immunology</i> , 2012, 3, 398.	2.2	36
54	Essential Role for Thymosin β 4 in Regulating Vascular Smooth Muscle Cell Development and Vessel Wall Stability. <i>Circulation Research</i> , 2012, 111, e89-102.	2.0	54

#	ARTICLE	IF	CITATIONS
55	An Emerging Role of RNA-Binding Proteins as Multifunctional Regulators of Lymphocyte Development and Function. <i>Advances in Immunology</i> , 2012, 115, 161-185.	1.1	15
56	A New Mechanism of Gene Regulation Mediated by Noncoding RNA. <i>Journal of Immunology</i> , 2012, 189, 3-4.	0.4	4
57	Pten Loss in CD4 T Cells Enhances Their Helper Function but Does Not Lead to Autoimmunity or Lymphoma. <i>Journal of Immunology</i> , 2012, 188, 5935-5943.	0.4	31
58	Regulation of lymphocyte development and function by RNA-binding proteins. <i>Current Opinion in Immunology</i> , 2012, 24, 160-165.	2.4	18
59	T Cell Receptor Internalization from the Immunological Synapse Is Mediated by TC21 and RhoG GTPase-Dependent Phagocytosis. <i>Immunity</i> , 2011, 35, 208-222.	6.6	152
60	Interaction of Ras with P110 β Is Required for Thymic β -Selection in the Mouse. <i>Journal of Immunology</i> , 2011, 187, 4667-4675.	0.4	9
61	Is Transcription the Dominant Force During Dynamic Changes in Gene Expression?. <i>Advances in Experimental Medicine and Biology</i> , 2011, 780, 1-13.	0.8	12
62	Putative GTPase GIMAP1 is critical for the development of mature B and T lymphocytes. <i>Blood</i> , 2010, 115, 3249-3257.	0.6	48
63	Deletion of the RNA-binding proteins ZFP36L1 and ZFP36L2 leads to perturbed thymic development and T lymphoblastic leukemia. <i>Nature Immunology</i> , 2010, 11, 717-724.	7.0	187
64	A novel Rac-dependent checkpoint in B cell development controls entry into the splenic white pulp and cell survival. <i>Journal of Experimental Medicine</i> , 2010, 207, 837-853.	4.2	55
65	Phosphoinositide 3-Kinase Activity in T Cells Regulates the Magnitude of the Germinal Center Reaction. <i>Journal of Immunology</i> , 2010, 185, 4042-4052.	0.4	200
66	Thymic development beyond β -selection requires phosphatidylinositol 3-kinase activation by CXCR4. <i>Journal of Experimental Medicine</i> , 2010, 207, 247-261.	4.2	143
67	MicroRNA 125b inhibition of B cell differentiation in germinal centers. <i>International Immunology</i> , 2010, 22, 583-592.	1.8	141
68	The development of mature B lymphocytes requires the combined function of CD19 and the p110 β subunit of PI3K. <i>Self/nonself</i> , 2010, 1, 144-153.	2.0	8
69	Stromal cell-derived factor 1 α and CXCR4: newly defined requirements for efficient thymic β -selection. <i>Trends in Immunology</i> , 2010, 31, 370-376.	2.9	26
70	Signaling Pathways in T Follicular Helper Cells. <i>Journal of Immunology</i> , 2010, 184, 6563-6568.	0.4	42
71	A novel Rac-dependent checkpoint in B cell development controls entry into the splenic white pulp and cell survival. <i>Journal of Cell Biology</i> , 2010, 189, i1-i1.	2.3	0
72	Vav GEFs regulate macrophage morphology and adhesion-induced Rac and Rho activation. <i>Experimental Cell Research</i> , 2009, 315, 3345-3358.	1.2	39

#	ARTICLE	IF	CITATIONS
73	Detection of protein-protein interactions by ribosome display and protein in situ immobilisation. <i>New Biotechnology</i> , 2009, 26, 277-281.	2.4	4
74	Cutting Edge: The Foxp3 Target miR-155 Contributes to the Development of Regulatory T Cells. <i>Journal of Immunology</i> , 2009, 182, 2578-2582.	0.4	350
75	Rac GTPases play critical roles in early T-cell development. <i>Blood</i> , 2009, 113, 3990-3998.	0.6	64
76	THE ROLE OF PI3K SIGNALLING IN THE B CELL RESPONSE TO ANTIGEN. <i>Advances in Experimental Medicine and Biology</i> , 2009, 633, 43-53.	0.8	16
77	B-cell responses to B-cell activation factor of the TNF family (BAFF) are impaired in the absence of PI3K delta. <i>European Journal of Immunology</i> , 2008, 38, 3543-3548.	1.6	86
78	Detection of transforming growth factor-beta in rheumatoid arthritis synovial tissue: lack of effect on spontaneous cytokine production in joint cell cultures. <i>Clinical and Experimental Immunology</i> , 2008, 81, 278-285.	1.1	83
79	Regulation of B-cell differentiation by microRNAs and RNA-binding proteins. <i>Biochemical Society Transactions</i> , 2008, 36, 1191-1193.	1.6	16
80	Activation of the Small GTPase Rac2 via the B Cell Receptor Regulates B Cell Adhesion and Immunological-Synapse Formation. <i>Immunity</i> , 2008, 28, 88-99.	6.6	148
81	The Effect of Deleting p110 δ on the Phenotype and Function of PTEN-Deficient B Cells. <i>Journal of Immunology</i> , 2008, 180, 739-746.	0.4	40
82	Tribbles-2 is a novel regulator of inflammatory activation of monocytes. <i>International Immunology</i> , 2008, 20, 1543-1550.	1.8	53
83	Regulation of B- and T-cell differentiation by a single microRNA. <i>Biochemical Society Transactions</i> , 2008, 36, 531-533.	1.6	65
84	Cutting Edge: The PI3K p110 δ Is Required for Down-Regulation of RAG Expression in Immature B Cells. <i>Journal of Immunology</i> , 2007, 178, 1981-1985.	0.4	52
85	The 3BP2 Adapter Protein Is Required for Optimal B-Cell Activation and Thymus-Independent Type 2 Humoral Response. <i>Molecular and Cellular Biology</i> , 2007, 27, 3109-3122.	1.1	45
86	Vav family proteins are required for optimal regulation of PLC β 2 by integrin α IIb β 3. <i>Biochemical Journal</i> , 2007, 401, 753-761.	1.7	44
87	The p110 δ catalytic isoform of PI3K is a key player in NK-cell development and cytokine secretion. <i>Blood</i> , 2007, 110, 3202-3208.	0.6	83
88	microRNA-155 Regulates the Generation of Immunoglobulin Class-Switched Plasma Cells. <i>Immunity</i> , 2007, 27, 847-859.	6.6	724
89	Requirement of bic/microRNA-155 for Normal Immune Function. <i>Science</i> , 2007, 316, 608-611.	6.0	1,786
90	Syk and Slp-76 Mutant Mice Reveal a Cell-Autonomous Hematopoietic Cell Contribution to Vascular Development. <i>Developmental Cell</i> , 2006, 11, 349-361.	3.1	115

#	ARTICLE	IF	CITATIONS
91	Phospholipase C- β 2 is essential for NK cell cytotoxicity and innate immunity to malignant and virally infected cells. <i>Blood</i> , 2006, 107, 994-1002.	0.6	120
92	Cellular Notch responsiveness is defined by phosphoinositide 3-kinase-dependent signals. <i>BMC Cell Biology</i> , 2006, 7, 10.	3.0	70
93	G β 1 β s and the Ras binding domain of p110 β are both important regulators of PI3K β signalling in neutrophils. <i>Nature Cell Biology</i> , 2006, 8, 1303-1309.	4.6	167
94	Identification of an imprinting control region affecting the expression of all transcripts in the <i>Gnas</i> cluster. <i>Nature Genetics</i> , 2006, 38, 350-355.	9.4	176
95	Synergistic activation of PKD by the B cell antigen receptor and CD19 requires PI3K, Vav1 and PLC β 3. <i>Cellular Signalling</i> , 2006, 18, 1455-1460.	1.7	13
96	The RNA binding protein Zfp361 is required for normal vascularisation and post-transcriptionally regulates VEGF expression. <i>Developmental Dynamics</i> , 2006, 235, 3144-3155.	0.8	93
97	A genome-scale assessment of peripheral blood B-cell molecular homeostasis in patients with rheumatoid arthritis. <i>Rheumatology</i> , 2006, 45, 1466-1476.	0.9	38
98	RhoG Regulates the Neutrophil NADPH Oxidase. <i>Journal of Immunology</i> , 2006, 176, 5314-5320.	0.4	37
99	Differential Requirements of PI3K Subunits for BCR or BCR/CD19-Induced ERK Activation. , 2006, 584, 43-52.		2
100	Vav proteins regulate peripheral B-cell survival. <i>Blood</i> , 2005, 106, 2391-2398.	0.6	46
101	The role of endothelial PI3K β activity in neutrophil trafficking. <i>Blood</i> , 2005, 106, 150-157.	0.6	169
102	Vav proteins are required for B-lymphocyte responses to LPS. <i>Blood</i> , 2005, 106, 635-640.	0.6	48
103	Sequential activation of class IB and class IA PI3K is important for the primed respiratory burst of human but not murine neutrophils. <i>Blood</i> , 2005, 106, 1432-1440.	0.6	274
104	Phosphatidylinositol 3-kinase is required for the transcriptional activation of cyclin D2 in BCR activated primary mouse B α lymphocytes. <i>European Journal of Immunology</i> , 2005, 35, 2748-2761.	1.6	15
105	Role of the p110 β PI 3-kinase in integrin and ITAM receptor signalling in platelets. <i>Platelets</i> , 2005, 16, 191-202.	1.1	47
106	Cutting Edge: T Cell Development Requires the Combined Activities of the p110 β and p110 δ Catalytic Isoforms of Phosphatidylinositol 3-Kinase. <i>Journal of Immunology</i> , 2005, 175, 2783-2787.	0.4	142
107	Vav1 and Vav2 play different roles in macrophage migration and cytoskeletal organization. <i>Experimental Cell Research</i> , 2005, 310, 303-310.	1.2	40
108	The Role of Endothelial PI3K β Activity in Neutrophil Trafficking.. <i>Blood</i> , 2005, 106, 3891-3891.	0.6	0

#	ARTICLE	IF	CITATIONS
109	Differential Regulation of TCR-mediated Gene Transcription by Vav Family Members. <i>Journal of Experimental Medicine</i> , 2004, 199, 429-434.	4.2	35
110	Vav1 and Vav3 Have Critical but Redundant Roles in Mediating Platelet Activation by Collagen. <i>Journal of Biological Chemistry</i> , 2004, 279, 53955-53962.	1.6	91
111	Vav-Dependent and Vav-Independent Phosphatidylinositol 3-Kinase Activation in Murine B Cells Determined by the Nature of the Stimulus. <i>Journal of Immunology</i> , 2004, 173, 3209-3214.	0.4	46
112	Immunological Function in Mice Lacking the Rac-Related GTPase RhoG. <i>Molecular and Cellular Biology</i> , 2004, 24, 719-729.	1.1	62
113	A cis-acting control region is required exclusively for the tissue-specific imprinting of Gnas. <i>Nature Genetics</i> , 2004, 36, 894-899.	9.4	157
114	The tyrosine kinase Syk is required for light chain isotype exclusion but dispensable for the negative selection of B cells. <i>European Journal of Immunology</i> , 2004, 34, 1102-1110.	1.6	19
115	PLC β 2 regulates Bcl-2 levels and is required for survival rather than differentiation of marginal zone and follicular B cells. <i>European Journal of Immunology</i> , 2004, 34, 2237-2247.	1.6	27
116	Mechanisms and implications of phosphoinositide 3-kinase $\hat{\Gamma}$ in promoting neutrophil trafficking into inflamed tissue. <i>Blood</i> , 2004, 103, 3448-3456.	0.6	198
117	BCR activation of PI3K is Vav-independent in murine B cells. <i>Biochemical Society Transactions</i> , 2004, 32, 781-784.	1.6	3
118	The p110 $\hat{\Gamma}$ subunit of phosphoinositide 3-kinase is required for the lipopolysaccharide response of mouse B cells. <i>Biochemical Society Transactions</i> , 2004, 32, 789-791.	1.6	23
119	RhoG regulates gene expression and the actin cytoskeleton in lymphocytes. <i>Oncogene</i> , 2003, 22, 330-342.	2.6	46
120	Regulation of Vav Localization in Membrane Rafts by Adaptor Molecules Grb2 and BLNK. <i>Immunity</i> , 2003, 18, 777-787.	6.6	59
121	A Crucial Role for the p110 $\hat{\Gamma}$ Subunit of Phosphatidylinositol 3-Kinase in B Cell Development and Activation. <i>Journal of Experimental Medicine</i> , 2002, 196, 753-763.	4.2	417
122	Vav1, but not Vav2, contributes to platelet aggregation by CRP and thrombin, but neither is required for regulation of phospholipase C. <i>Blood</i> , 2002, 100, 3561-3569.	0.6	48
123	B-cell development and antigen receptor signalling. <i>Biochemical Society Transactions</i> , 2002, 30, 812-815.	1.6	9
124	The tyrosine kinase Lyn is required for B cell development beyond the T1 stage in the spleen: rescue by over-expression of Bcl-2. <i>European Journal of Immunology</i> , 2002, 32, 1029-1034.	1.6	46
125	Natural cytotoxicity uncoupled from the Syk and ZAP-70 intracellular kinases. <i>Nature Immunology</i> , 2002, 3, 288-294.	7.0	105
126	VAV proteins as signal integrators for multi-subunit immune-recognition receptors. <i>Nature Reviews Immunology</i> , 2002, 2, 476-486.	10.6	312

#	ARTICLE	IF	CITATIONS
127	The Role of Vav Proteins in B Cell Responses. <i>Advances in Experimental Medicine and Biology</i> , 2002, 512, 29-34.	0.8	4
128	Structural Organization of the Mouse Phosphatidylinositol 3-Kinase p110d Gene. <i>Biochemical and Biophysical Research Communications</i> , 2001, 280, 1328-1332.	1.0	5
129	Signal transduction through Vav-2 participates in humoral immune responses and B cell maturation. <i>Nature Immunology</i> , 2001, 2, 542-547.	7.0	169
130	Vav Is Required for Cyclin D2 Induction and Proliferation of Mouse B Lymphocytes Activated via the Antigen Receptor. <i>Journal of Biological Chemistry</i> , 2001, 276, 41040-41048.	1.6	31
131	Functional Dichotomy in Natural Killer Cell Signaling. <i>Journal of Experimental Medicine</i> , 2001, 193, 1413-1424.	4.2	75
132	Tyrosine kinase SYK: essential functions for immunoreceptor signalling. <i>Trends in Immunology</i> , 2000, 21, 148-154.	7.5	376
133	Development of T-leukaemias in CD45 tyrosine phosphatase-deficient mutant lck mice. <i>EMBO Journal</i> , 2000, 19, 4644-4654.	3.5	48
134	Vav-2 controls NFAT-dependent transcription in B- but not T-lymphocytes. <i>EMBO Journal</i> , 2000, 19, 6173-6184.	3.5	73
135	A New Look at Syk in $\hat{\alpha}\hat{\beta}$ and $\hat{\beta}\hat{\gamma}$ T Cell Development Using Chimeric Mice with a Low Competitive Hematopoietic Environment. <i>Journal of Immunology</i> , 2000, 164, 5140-5145.	0.4	22
136	Genetic and Pharmacological Analyses of Syk Function in $\hat{\alpha}\hat{\beta}$ Signaling in Platelets. <i>Blood</i> , 1999, 93, 2645-2652.	0.6	162
137	Tyrosine Phosphorylation of SLP-76 Is Downstream of Syk following Stimulation of the Collagen Receptor in Platelets. <i>Journal of Biological Chemistry</i> , 1999, 274, 5963-5971.	1.6	102
138	The Rho-family GTP exchange factor Vav is a critical transducer of T cell receptor signals to the calcium, ERK, and NF- $\hat{\kappa}$ B pathways. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1999, 96, 3035-3040.	3.3	235
139	Defective immunoglobulin class switching in Vav-deficient mice is attributable to compromised T cell help. <i>European Journal of Immunology</i> , 1999, 29, 477-487.	1.6	48
140	The CD45 tyrosine phosphatase regulates CD3-induced signal transduction and T cell development in recombina-se-deficient mice: restoration of pre-TCR function by active p56lck. <i>European Journal of Immunology</i> , 1999, 29, 2376-2384.	1.6	36
141	Greatly reduced efficiency of both positive and negative selection of thymocytes in CD45 tyrosine phosphatase-deficient mice. <i>European Journal of Immunology</i> , 1999, 29, 2923-2933.	1.6	67
142	The CD45 tyrosine phosphatase regulates CD3-induced signal transduction and T cell development in recombina-se-deficient mice: restoration of pre-TCR function by active p56lck. , 1999, 29, 2376.		2
143	Greatly reduced efficiency of both positive and negative selection of thymocytes in CD45 tyrosine phosphatase-deficient mice. , 1999, 29, 2923.		2
144	Collagen Mediates Changes in Intracellular Calcium in Primary Mouse Megakaryocytes Through syk-Dependent and -Independent Pathways. <i>Blood</i> , 1999, 93, 3847-3855.	0.6	21

#	ARTICLE	IF	CITATIONS
145	Genetic and Pharmacological Analyses of Syk Function in $\alpha\text{IIb}\beta_3$ Signaling in Platelets. <i>Blood</i> , 1999, 93, 2645-2652.	0.6	16
146	Collagen Mediates Changes in Intracellular Calcium in Primary Mouse Megakaryocytes Through syk-Dependent and -Independent Pathways. <i>Blood</i> , 1999, 93, 3847-3855.	0.6	0
147	Redundant role of the Syk protein tyrosine kinase in mouse NK cell differentiation. <i>Journal of Immunology</i> , 1999, 163, 1769-74.	0.4	30
148	CD19 as a Membrane-Anchored Adaptor Protein of B Lymphocytes: Costimulation of Lipid and Protein Kinases by Recruitment of Vav. <i>Immunity</i> , 1998, 8, 635-645.	6.6	177
149	Syk Tyrosine Kinase Is Required for the Positive Selection of Immature B Cells into the Recirculating B Cell Pool. <i>Journal of Experimental Medicine</i> , 1997, 186, 2013-2021.	4.2	147
150	A Critical Role for Syk in Signal Transduction and Phagocytosis Mediated by $\text{Fc}\gamma_3$ Receptors on Macrophages. <i>Journal of Experimental Medicine</i> , 1997, 186, 1027-1039.	4.2	471
151	Syk and Fyn Are Required by Mouse Megakaryocytes for the Rise in Intracellular Calcium Induced by a Collagen-related Peptide. <i>Journal of Biological Chemistry</i> , 1997, 272, 27539-27542.	1.6	55
152	A Requirement for the Rho-Family GTP Exchange Factor Vav in Positive and Negative Selection of Thymocytes. <i>Immunity</i> , 1997, 7, 451-460.	6.6	268
153	The $\text{Fc}\gamma_3$ receptor β -chain and the tyrosine kinase Syk are essential for activation of mouse platelets by collagen. <i>EMBO Journal</i> , 1997, 16, 2333-2341.	3.5	416
154	Critical role for the tyrosine kinase Syk in signalling through the high affinity IgE receptor of mast cells. <i>Oncogene</i> , 1996, 13, 2595-605.	2.6	249
155	Defective antigen receptor-mediated proliferation of B and T cells in the absence of Vav. <i>Nature</i> , 1995, 374, 467-470.	13.7	399
156	Perinatal lethality and blocked B-cell development in mice lacking the tyrosine kinase Syk. <i>Nature</i> , 1995, 378, 298-302.	13.7	706
157	Characterization of ligand binding by the human p55 tumour-necrosis-factor receptor. Involvement of individual cysteine-rich repeats. <i>FEBS Journal</i> , 1994, 223, 831-840.	0.2	13
158	Cytokine Assays: Role in Evaluation of the Pathogenesis of Autoimmunity. <i>Immunological Reviews</i> , 1991, 119, 105-123.	2.8	86
159	Production of Interleukin-1 and Interleukin-6 by Human Keratinocytes and Squamous Cell Carcinoma Cell Lines. <i>Journal of Investigative Dermatology</i> , 1991, 96, 771-776.	0.3	74
160	Induction of the interleukin 1 receptor antagonist protein by transforming growth factor- β_2 . <i>European Journal of Immunology</i> , 1991, 21, 1635-1639.	1.6	181
161	Expression of granulocyte-macrophage colony-stimulating factor in rheumatoid arthritis: Regulation by tumor necrosis factor. <i>European Journal of Immunology</i> , 1991, 21, 2575-2579.	1.6	288
162	CD4+ T-Cell Clones from Autoimmune Thyroid Tissue Cannot be Classified According to their Lymphokine Production. <i>Scandinavian Journal of Immunology</i> , 1990, 32, 433-440.	1.3	40

#	ARTICLE	IF	CITATIONS
163	Detection of interleukin 8 biological activity in synovial fluids from patients with rheumatoid arthritis and production of interleukin 8 mRNA by isolated synovial cells. <i>European Journal of Immunology</i> , 1990, 20, 2141-2144.	1.6	288
164	Cloning of human tumor necrosis factor (TNF) receptor cDNA and expression of recombinant soluble TNF-binding protein.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1990, 87, 7380-7384.	3.3	130
165	Effects of interferon alpha on autocrine growth factor loops in B lymphoproliferative disorders.. <i>Journal of Experimental Medicine</i> , 1990, 172, 1729-1734.	4.2	73
166	Transforming growth factor β^2 induces the production of interleukin 6 by human peripheral blood mononuclear cells. <i>Cytokine</i> , 1990, 2, 211-216.	1.4	80
167	Granulocyte-macrophage colony stimulating factor induces both HLA-DR expression and cytokine production by human monocytes. <i>Cytokine</i> , 1990, 2, 60-67.	1.4	51
168	Interleukin 7 and interleukin 4 stimulate human thymocyte growth through distinct mechanisms. <i>Cytokine</i> , 1990, 2, 55-59.	1.4	8
169	Interleukin 7 (murine pre-B cell growth factor/lymphopoietin 1) stimulates thymocyte growth: regulation by transforming growth factor beta. <i>European Journal of Immunology</i> , 1989, 19, 783-786.	1.6	66
170	Intrathyroidal cytokine production in thyroid disease. <i>Journal of Autoimmunity</i> , 1989, 2, 171-176.	3.0	12
171	Transforming growth factor beta regulates thyroid growth. Role in the pathogenesis of nontoxic goiter.. <i>Journal of Clinical Investigation</i> , 1989, 83, 764-770.	3.9	125
172	Intrathyroidal Cytokine Production in Thyroid Disease. , 1989, , 171-176.		1
173	Effect of cytokines on HLA-DR and IL-1 production by a monocytic tumour, THP-1. <i>Immunology</i> , 1989, 66, 170-5.	2.0	18
174	Analysis of intrathyroidal cytokine production in thyroid autoimmune disease: thyroid follicular cells produce interleukin-1 alpha and interleukin-6. <i>Clinical and Experimental Immunology</i> , 1989, 77, 324-30.	1.1	94
175	Regulation of expression of human IL-1 alpha and IL-1 beta genes. <i>Journal of Immunology</i> , 1989, 143, 3556-61.	0.4	67
176	Modulation of cytokine production by transforming growth factor-beta. <i>Journal of Immunology</i> , 1989, 142, 4295-300.	0.4	260
177	Excessive production of interleukin 6/B cell stimulatory factor-2 in rheumatoid arthritis. <i>European Journal of Immunology</i> , 1988, 18, 1797-1802.	1.6	790
178	TUMOUR NECROSIS FACTOR AS AN AUTOCRINE TUMOUR GROWTH FACTOR FOR CHRONIC B-CELL MALIGNANCIES. <i>Lancet, The</i> , 1988, 331, 969-971.	6.3	250
179	Post-transcriptional control of IL-1 gene expression in the acute monocytic leukemia line THP-1. <i>Biochemical and Biophysical Research Communications</i> , 1988, 156, 830-839.	1.0	25
180	Comparison of patterns of expression of tumour necrosis factor, lymphotoxin and interleukin-6 mRNA. <i>Biochemical and Biophysical Research Communications</i> , 1988, 153, 1144-1151.	1.0	27

#	ARTICLE	IF	CITATIONS
181	Role of HLA Class II and Cytokine Expression in Rheumatoid Arthritis. Scandinavian Journal of Rheumatology, 1988, 17, 39-46.	0.6	8
182	Interleukin-1 and tumour necrosis factor mRNA expression in rheumatoid arthritis: prolonged production of IL-1 alpha. Clinical and Experimental Immunology, 1988, 73, 449-55.	1.1	279
183	Does the maternal kidney contribute to the increased circulating 1,25-dihydroxyvitamin D concentrations during pregnancy?. Mineral and Electrolyte Metabolism, 1988, 14, 246-52.	1.1	61
184	Effects of tumour necrosis factor and alpha interferon on chronic B cell malignancies. , 1988, 30, 317-9.		1
185	Human T cells from autoimmune and normal individuals can produce tumor necrosis factor. European Journal of Immunology, 1987, 17, 1807-1814.	1.6	94