

Jian Wang

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/422345/jian-wang-publications-by-year.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

41
papers

776
citations

17
h-index

26
g-index

46
ext. papers

907
ext. citations

2.9
avg, IF

4.68
L-index

#	Paper	IF	Citations
41	o-Nitroacetanilide Solubility in Several Aqueous Solutions and Its Mathematical Correlation. <i>Journal of Chemical & Engineering Data</i> , 2022 , 67, 786-796	2.8	
40	Solubility Modeling, Solvent Effect, and Dissolution Properties of 4-Nitrophenylacetic Acid in Thirteen Solvents Ranging from 283.15 to 328.15 K. <i>Journal of Chemical & Engineering Data</i> , 2020 , 65, 2894-2902	2.8	10
39	Solubility of monobenzene in aqueous co-solvent mixtures of several alcohols: Determination, modelling and thermodynamic aspects analysis. <i>Journal of Chemical Thermodynamics</i> , 2020 , 142, 106023 ^{2.9}	2.9	2
38	4-Chloro-2-nitroaniline Solubility in Several Pure Solvents: Determination, Modeling, and Solvent Effect Analysis. <i>Journal of Chemical & Engineering Data</i> , 2020 , 65, 222-232	2.8	10
37	Equilibrium Solubility Investigation and Preferential Solvation of 2,6-Dichloro-4-nitroaniline Dissolved in Four Aqueous Mixtures of Isopropanol, Acetonitrile, n-Propanol, and N-Methyl-2-pyrrolidone. <i>Journal of Chemical & Engineering Data</i> , 2020 , 65, 2912-2920	2.8	4
36	Equilibrium Solubility and Dissolution Property Analysis of 2-Nitrophenylacetic Acid in 13 Pure Solvents at Elevated Temperatures. <i>Journal of Chemical & Engineering Data</i> , 2020 , 65, 4157-4165	2.8	4
35	Saturated Solubility of 2-Acrylamide-2-methylpropanesulfonic Acid in 14 Neat Organic Solvents from 283.15 to 328.15 K. <i>Journal of Chemical & Engineering Data</i> , 2020 , 65, 4692-4698	2.8	2
34	4,4'-Methylene-bis-(2-chloroaniline) Dissolved in Some Neat Solvents: Saturated Solubility, Mixing Properties, and Solvent Effect. <i>Journal of Chemical & Engineering Data</i> , 2020 , 65, 5018-5029	2.8	
33	Physicochemical properties of switchable-hydrophilicity solvent systems: N,N-Dimethylcyclohexylamine, water and carbon dioxide. <i>Journal of Chemical Thermodynamics</i> , 2019 , 133, 1-9	2.9	7
32	Equilibrium solubility of dinitolmide in several neat solvents and binary aqueous co-solvent mixtures: Experimental determination and thermodynamic analysis. <i>Journal of Chemical Thermodynamics</i> , 2019 , 132, 373-382	2.9	24
31	Solubility of d-Aspartic Acid in Several Neat Solvents: Determination, Modeling, and Solvent Effect Analysis. <i>Journal of Chemical & Engineering Data</i> , 2019 , 64, 2904-2910	2.8	12
30	Solubility Measurement and Thermodynamic Model Correlation and Evaluation of 2-Chloro-5-nitroaniline in 12 Pure Solvents. <i>Journal of Chemical & Engineering Data</i> , 2019 , 64, 1357-1365 ^{2.8}	2.8	11
29	Solubility and Solution Thermodynamics of 3-Nitrophthalonitrile in 12 Neat Solvents at Temperatures from 278.15 to 323.15 K. <i>Journal of Chemical & Engineering Data</i> , 2019 , 64, 3250-3259 ^{2.8}	2.8	20
28	Solubility Study and Mixing Property of 3,5-Dinitro-2-methylbenzoic Acid in 13 Pure Solvents from 288.15 to 333.15 K. <i>Journal of Chemical & Engineering Data</i> , 2019 , 64, 3652-3660	2.8	12
27	Solubility and Solution Thermodynamics of 2,6-Dichloro-4-nitroaniline in 12 Pure Solvents at Temperatures from 278.15 to 323.15 K. <i>Journal of Chemical & Engineering Data</i> , 2019 , 64, 5869-5877 ^{2.8}	2.8	11
26	Thermodynamic Solubility of Gatifloxacin in Nonaqueous Solvent Mixtures of N,N-Dimethylformamide Plus Isopropanol/Methanol/n-Propanol/Ethanol. <i>Journal of Chemical & Engineering Data</i> , 2019 , 64, 6071-6077	2.8	
25	Measurement and Correlation of Solubility of Gatifloxacin in 12 Pure Solvents from 273.15 K to 318.15 K. <i>Journal of Chemical & Engineering Data</i> , 2019 , 64, 676-681	2.8	3

24	Comment on Density, Sound Speed, and Viscosity of Dihydropyridine Derivatives in Dimethyl Sulfoxide at Different Temperatures <i>Journal of Chemical & Engineering Data</i> , 2018 , 63, 3165-3166	2.8	
23	Density, viscosity, refractive index and conductivity of switchable hydrophilicity solvent systems: N,N-Dimethylbenzylamine, water and carbon dioxide. <i>Fluid Phase Equilibria</i> , 2018 , 459, 94-102	2.5	3
22	Solubility of Terephthaldialdehyde in N-Methyl-2-pyrrolidone and Solid-Liquid Phase Equilibrium for Ternary Systems of Terephthalic Acid + Terephthaldialdehyde + N,N-Dimethylformamide/N-Methyl-2-pyrrolidone. <i>Journal of Chemical & Engineering Data</i> , 2018 , 63, 2081-2090	2.8	5
21	Solubility of 2-nitro-p-phenylenediamine in nine pure solvents and mixture of (methanol+N-methyl-2-pyrrolidone) from T=(283.15 to 318.15)K: Determination and modelling. <i>Journal of Chemical Thermodynamics</i> , 2017 , 108, 45-58	2.9	39
20	Solubility and solution thermodynamics of 4-hydroxybenzaldehyde in twelve organic solvents from T = (278.15 to 318.15) K. <i>Journal of Molecular Liquids</i> , 2017 , 237, 226-235	6	30
19	Solubility determination and thermodynamic functions of 2-chlorophenothiazine in nine organic solvents from T = 283.15 K to T = 318.15 K and mixing properties of solutions. <i>Journal of Chemical Thermodynamics</i> , 2017 , 106, 132-144	2.9	42
18	Solubility of amiodarone hydrochloride in aqueous co-solvent mixtures revisited: IBKI preferential solvation analysis. <i>Journal of Chemical Thermodynamics</i> , 2017 , 112, 276-282	2.9	11
17	Determination and thermodynamic modelling for 2-methyl-6-nitroaniline solubility in binary solvent mixtures of ethyl acetate + (methanol, ethanol, n -propanol and isopropanol). <i>Journal of Chemical Thermodynamics</i> , 2017 , 105, 404-413	2.9	43
16	Determination and correlation of terephthaldialdehyde solubility in (ethanol, isopropanol, ethyl acetate, isopentanol) + N,N-dimethylformamide mixed solvents at temperatures from 273.15 K to 318.15 K. <i>Journal of Chemical Thermodynamics</i> , 2017 , 105, 327-336	2.9	39
15	Thermodynamic modelling for solubility of 5-chloro-1-methyl-4-nitroimidazole in eleven organic solvents from T = (283.15 to 318.15) K. <i>Journal of Chemical Thermodynamics</i> , 2017 , 105, 58-70	2.9	33
14	Thermodynamic modelling for solubility of 4-nitrobenzaldehyde in different solvents at temperature range from (273.15 to 313.15) K and mixing properties of solutions. <i>Journal of Chemical Thermodynamics</i> , 2017 , 104, 50-60	2.9	7
13	Solubility Measurement and Thermodynamic Modeling of 4-Nitrophthalimide in Twelve Pure Solvents at Elevated Temperatures Ranging from (273.15 to 323.15) K. <i>Journal of Chemical & Engineering Data</i> , 2016 , 61, 2525-2535	2.8	68
12	Solubility determination and thermodynamic models for dehydroepiandrosterone acetate in mixed solvents of (ethyl acetate + methanol), (ethyl acetate + ethanol) and (ethyl acetate + isopropanol). <i>Journal of Chemical Thermodynamics</i> , 2016 , 101, 372-379	2.9	84
11	Solubility modelling and thermodynamic dissolution functions of phthalimide in ten organic solvents. <i>Journal of Chemical Thermodynamics</i> , 2016 , 94, 160-168	2.9	17
10	Solubility Modeling of 4-(Methylsulfonyl)benzaldehyde in Nine Organic Solvents at Elevated Temperatures. <i>Journal of Chemical & Engineering Data</i> , 2016 , 61, 1657-1666	2.8	10
9	Solubility and Dissolution Thermodynamics for 1,3,5-Trichlorobenzene in Organic Solvents. <i>Journal of Chemical & Engineering Data</i> , 2016 , 61, 380-390	2.8	26
8	Solubility determination and thermodynamic dissolution functions of 1,3-diphenylguanidine in nine organic solvents at evaluated temperatures. <i>Journal of Chemical Thermodynamics</i> , 2016 , 99, 86-95	2.9	9
7	Solubility of 3-chloro-N-phenylphthalimide in ten organic solvents from T = (288.15 to 323.15) K: Determination and modelling. <i>Journal of Chemical Thermodynamics</i> , 2016 , 96, 187-195	2.9	20

6	Solubility determination and thermodynamic modeling of 2,4-dinitroaniline in nine organic solvents from T = (278.15 to 318.15) K and mixing properties of solutions. <i>Journal of Chemical Thermodynamics</i> , 2016 , 102, 178-187	2.9	22
5	Solubility determination and thermodynamic modelling of terephthalaldehyde in ten organic solvents from T = (273.15 to 318.15) K and mixing properties of solutions. <i>Journal of Chemical Thermodynamics</i> , 2016 , 102, 188-198	2.9	63
4	Determination and Correlation of Solid-Liquid Phase Equilibrium and Phase Diagram for a Multicomponent System of Mixed Dibasic Acids. Ternary System of Succinic Acid + Adipic Acid + Ethanol. <i>Journal of Chemical & Engineering Data</i> , 2016 , 61, 2105-2113	2.8	17
3	Determination and thermodynamic modeling of solid-liquid phase equilibrium for 3,5-dichloroaniline in pure solvents and ternary 3,5-dichloroaniline + 1,3,5-trichlorobenzene + toluene system. <i>Journal of Chemical Thermodynamics</i> , 2016 , 98, 288-297	2.9	13
2	Solubility measurement and modelling of 1,8-dinitronaphthalene in nine organic solvents from T=(273.15 to 308.15)K and mixing properties of solutions. <i>Journal of Chemical Thermodynamics</i> , 2015 , 90, 259-269	2.9	22
1	Determination and prediction of solid-liquid phase equilibrium for quaternary system of terephthalic acid+isophthalic acid+phthalic acid+N-methyl-2-pyrrolidone at 303.15K and 313.15K. <i>Fluid Phase Equilibria</i> , 2015 , 397, 103-110	2.5	19