

Mnica Vazquez-Del Mercado

List of Publications by Citations

Source:

<https://exaly.com/author-pdf/4226577/monica-vazquez-del-mercado-publications-by-citations.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

109
papers

1,349
citations

19
h-index

30
g-index

132
ext. papers

1,653
ext. citations

3.5
avg, IF

3.88
L-index

#	Paper	IF	Citations
109	The EuroMyositis registry: an international collaborative tool to facilitate myositis research. <i>Annals of the Rheumatic Diseases</i> , 2018 , 77, 30-39	2.4	105
108	Clinical interpretation of antinuclear antibody tests in systemic rheumatic diseases. <i>Modern Rheumatology</i> , 2009 , 19, 219-228	3.3	88
107	Implications in the difference of anti-Mi-2 and -p155/140 autoantibody prevalence in two dermatomyositis cohorts from Mexico City and Guadalajara. <i>Arthritis Research and Therapy</i> , 2013 , 15, R48	5.7	50
106	T(H)1/T(H)2 cytokine profile, metalloprotease-9 activity and hormonal status in pregnant rheumatoid arthritis and systemic lupus erythematosus patients. <i>Clinical and Experimental Immunology</i> , 2003 , 131, 377-84	6.2	49
105	Clinical interpretation of antinuclear antibody tests in systemic rheumatic diseases. <i>Modern Rheumatology</i> , 2009 , 19, 219-28	3.3	44
104	IL-17 increased in the third trimester in healthy women with term labor. <i>American Journal of Reproductive Immunology</i> , 2011 , 65, 99-103	3.8	42
103	Update of the Mexican College of Rheumatology guidelines for the pharmacologic treatment of rheumatoid arthritis. <i>Reumatología Clínica</i> , 2014 , 10, 227-40	0.9	37
102	A controlled clinical trial with pirfenidone in the treatment of pathological skin scarring caused by burns in pediatric patients. <i>Annals of Plastic Surgery</i> , 2012 , 68, 22-8	1.7	35
101	Anti-Cyclic Citrullinated Peptide Antibodies and Severity of Interstitial Lung Disease in Women with Rheumatoid Arthritis. <i>Journal of Immunology Research</i> , 2015 , 2015, 151626	4.5	31
100	Polymorphisms C677T and A1298C in the MTHFR gene in Mexican patients with rheumatoid arthritis treated with methotrexate: implication with elevation of transaminases. <i>Pharmacogenomics Journal</i> , 2011 , 11, 287-91	3.5	31
99	Prolactin has a pathogenic role in systemic lupus erythematosus. <i>Immunologic Research</i> , 2017 , 65, 512-523		30
98	Expression of ICAM1 and VCAM1 Serum Levels in Rheumatoid Arthritis Clinical Activity. Association with Genetic Polymorphisms. <i>Disease Markers</i> , 2009 , 26, 119-126	3.2	30
97	The +49A>G CTLA-4 polymorphism is associated with rheumatoid arthritis in Mexican population. <i>Clinica Chimica Acta</i> , 2010 , 411, 725-8	6.2	29
96	Tumor necrosis factor alpha-308 and -238 polymorphisms in rheumatoid arthritis. Association with messenger RNA expression and sTNF-alpha. <i>Journal of Investigative Medicine</i> , 2008 , 56, 937-43	2.9	27
95	Serum levels of anticyclic citrullinated peptide antibodies, interleukin-6, tumor necrosis factor- α and C-reactive protein are associated with increased carotid intima-media thickness: a cross-sectional analysis of a cohort of rheumatoid arthritis patients without cardiovascular risk factors. <i>Journal of Rheumatology</i> , 2015 , 2015, 3426-32	3	24
94	Inhibitors of MAPK pathway ERK1/2 or p38 prevent the IL-1 β -induced up-regulation of SRP72 autoantigen in Jurkat cells. <i>Journal of Biological Chemistry</i> , 2010 , 285, 32824-32833	5.4	23
93	Genotype and allele frequency of PAI-1 promoter polymorphism in healthy subjects from the west of Mexico. Association with biochemical and hematological parameters. <i>Annales De Génétique</i> , 2004 , 47, 155-62		21

92	Serum levels of adiponectin and leptin as biomarkers of proteinuria in lupus nephritis. <i>PLoS ONE</i> , 2017 , 12, e0184056	3.7	20
91	Autoantibodies to transcription intermediary factor TIF1 β associated with dermatomyositis. <i>Arthritis Research and Therapy</i> , 2012 , 14, R79	5.7	20
90	Reduced IgG anti-small nuclear ribonucleoprotein autoantibody production in systemic lupus erythematosus patients with positive IgM anti-cytomegalovirus antibodies. <i>Arthritis Research and Therapy</i> , 2009 , 11, R27	5.7	19
89	Massive myeloid sarcoma affecting the central nervous system, mediastinum, retroperitoneum, liver, and rectum associated with acute myeloblastic leukaemia: a case report. <i>Journal of Clinical Pathology</i> , 2005 , 58, 325-7	3.9	19
88	Autoantibodies to Argonaute 2 (Su antigen). <i>Advances in Experimental Medicine and Biology</i> , 2013 , 768, 45-59	3.6	18
87	The Impact of LEP G-2548A and LEPR Gln223Arg Polymorphisms on Adiposity, Leptin, and Leptin-Receptor Serum Levels in a Mexican Mestizo Population. <i>BioMed Research International</i> , 2015 , 2015, 539408	3	18
86	The -174G/C and -572G/C interleukin 6 promoter gene polymorphisms in mexican patients with rheumatoid arthritis: a case-control study. <i>Clinical and Developmental Immunology</i> , 2013 , 2013, 959084		18
85	Tumor necrosis factor receptor 2 M196R polymorphism in rheumatoid arthritis and osteoarthritis: relationship with sTNFR2 levels and clinical features. <i>Rheumatology International</i> , 2006 , 27, 53-9	3.6	18
84	Detection of autoantibodies to DSF70/LEDGFp75 in Mexican Hispanics using multiple complementary assay platforms. <i>Autoimmunity Highlights</i> , 2017 , 8, 1	3.7	17
83	Circulating TNFR1 and TNFR2 levels correlated with the disease activity score (DAS28) in rheumatoid arthritis. <i>Scandinavian Journal of Rheumatology</i> , 2009 , 38, 332-5	1.9	16
82	Fibrogenic polymorphisms (TGF-beta, PAI-1, AT) in Mexican patients with established liver fibrosis. Potential correlation with pirfenidone treatment. <i>Journal of Investigative Medicine</i> , 2008 , 56, 944-53	2.9	16
81	Genotype Ser413/Ser of PAI-2 polymorphism Ser413/Cys is associated with anti-phospholipid syndrome and systemic lupus erythematosus in a familial case: comparison with healthy controls. <i>Scandinavian Journal of Rheumatology</i> , 2007 , 36, 206-10	1.9	16
80	Pirfenidone Is an Agonistic Ligand for PPAR α and Improves NASH by Activation of SIRT1/LKB1/pAMPK. <i>Hepatology Communications</i> , 2020 , 4, 434-449	6	15
79	Increase levels of apo-A1 and apo B are associated in knee osteoarthritis: lack of association with VEGF-460 T/C and +405 C/G polymorphisms. <i>Rheumatology International</i> , 2008 , 29, 63-8	3.6	15
78	YKL-40 expression in CD14+ liver cells in acute and chronic injury. <i>World Journal of Gastroenterology</i> , 2011 , 17, 3830-5	5.6	15
77	Serum leptin and serum leptin/serum leptin receptor ratio imbalance in obese rheumatoid arthritis patients positive for anti-cyclic citrullinated peptide antibodies. <i>Arthritis Research and Therapy</i> , 2015 , 17, 335	5.7	13
76	Association of ADIPOQ +45T>G polymorphism with body fat mass and blood levels of soluble adiponectin and inflammation markers in a Mexican-Mestizo population. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2012 , 5, 369-78	3.4	13
75	Tofacitinib in juvenile idiopathic arthritis: a double-blind, placebo-controlled, withdrawal phase 3 randomised trial. <i>Lancet, The</i> , 2021 , 398, 1984-1996	40	13

74	Autoantibodies to Mi-2 alpha and Mi-2 beta in patients with idiopathic inflammatory myopathy. <i>Rheumatology</i> , 2019 , 58, 1655-1661	3.9	12
73	Disease duration of rheumatoid arthritis is a predictor of vascular stiffness: a cross-sectional study in patients without known cardiovascular comorbidities: A STROBE-compliant article. <i>Medicine (United States)</i> , 2017 , 96, e7862	1.8	12
72	The 482Ser of PPARGC1A and 12Pro of PPARG2 Alleles Are Associated with Reduction of Metabolic Risk Factors Even Obesity in a Mexican-Mestizo Population. <i>BioMed Research International</i> , 2015 , 2015, 285491	3	12
71	The -844 G/A PAI-1 polymorphism is associated with mRNA expression in rheumatoid arthritis. <i>Rheumatology International</i> , 2008 , 28, 355-60	3.6	12
70	Pulmonary function in ankylosing spondylitis: association with clinical variables. <i>Rheumatology International</i> , 2013 , 33, 2351-8	3.6	11
69	The ACTN3 R577X polymorphism is associated with inflammatory myopathies in a Mexican population. <i>Scandinavian Journal of Rheumatology</i> , 2012 , 41, 396-400	1.9	11
68	MTHFR C677T, MTHFR A1298C, and OPG A163G Polymorphisms in Mexican Patients with Rheumatoid Arthritis and Osteoporosis. <i>Disease Markers</i> , 2012 , 32, 109-114	3.2	11
67	A functional Ser(413)/Ser(413) PAI-2 polymorphism is associated with susceptibility and damage index score in systemic lupus erythematosus. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2009 , 15, 233-8	3.3	11
66	Prolonged-release pirfenidone prevents obesity-induced cardiac steatosis and fibrosis in a mouse NASH model. <i>Cardiovascular Drugs and Therapy</i> , 2021 , 35, 927-938	3.9	10
65	Liver fibrosis secondary to bile duct injury: correlation of Smad7 with TGF-beta and extracellular matrix proteins. <i>BMC Gastroenterology</i> , 2009 , 9, 81	3	10
64	Inverse Relationship of the CMKLR1 Relative Expression and Chemerin Serum Levels in Obesity with Dysmetabolic Phenotype and Insulin Resistance. <i>Mediators of Inflammation</i> , 2016 , 2016, 3085390	4.3	10
63	Association of interleukin-10 promoter haplotypes with disease susceptibility and IL-10 levels in Mexican patients with systemic lupus erythematosus. <i>Clinical and Experimental Medicine</i> , 2015 , 15, 439-46	4.9	9
62	FAS -670A>G promoter polymorphism is associated with soluble Fas levels in primary Sjögren's syndrome. <i>Genetics and Molecular Research</i> , 2014 , 13, 4831-8	1.2	9
61	Modifications in lipid levels are independent of serum TNF- α in rheumatoid arthritis: results of an observational 24-week cohort study comparing patients receiving etanercept plus methotrexate or methotrexate as monotherapy. <i>BioMed Research International</i> , 2014 , 2014, 510305	3	9
60	-383 A/C tumor necrosis factor receptor 1 polymorphism and ankylosing spondylitis in Mexicans: a preliminary study. <i>Rheumatology International</i> , 2012 , 32, 2565-8	3.6	9
59	Amebic monocyte locomotion inhibitory factor peptide ameliorates inflammation in CIA mouse model by downregulation of cell adhesion, inflammation/chemotaxis, and matrix metalloproteinases genes. <i>Inflammation Research</i> , 2010 , 59, 1041-51	7.2	9
58	Capsaicin induces a protective effect on gastric mucosa along with decreased expression of inflammatory molecules in a gastritis model. <i>Journal of Functional Foods</i> , 2019 , 59, 345-351	5.1	8
57	Procollagen Type I and III Aminoterminal Propeptide Levels and Severity of Interstitial Lung Disease in Mexican Women With Progressive Systemic Sclerosis. <i>Archivos De Bronconeumologia</i> , 2015 , 51, 440-8	0.7	8

56	High prevalence of autoantibodies to RNA helicase A in Mexican patients with systemic lupus erythematosus. <i>Arthritis Research and Therapy</i> , 2010 , 12, R6	5.7	8
55	Distribution of -844 G/A and Hind III C/G PAI-1 polymorphisms and plasma PAI-1 levels in Mexican subjects: comparison of frequencies between populations. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2008 , 14, 220-6	3.3	8
54	MTHFR C677T, MTHFR A1298C, and OPG A163G polymorphisms in Mexican patients with rheumatoid arthritis and osteoporosis. <i>Disease Markers</i> , 2012 , 32, 109-14	3.2	8
53	Ageing influences the relationship of circulating miR-33a and miR- 33b levels with insulin resistance and adiposity. <i>Diabetes and Vascular Disease Research</i> , 2019 , 16, 244-253	3.3	8
52	The Montreal Cognitive Assessment Test: A Useful Tool in Screening of Cognitive Impairment in Patients With Systemic Lupus Erythematosus. <i>Journal of Clinical Rheumatology</i> , 2019 , 25, 325-328	1.1	8
51	Downregulation of hippocampal NR2A/2B subunits related to cognitive impairment in a pristane-induced lupus BALB/c mice. <i>PLoS ONE</i> , 2019 , 14, e0217190	3.7	7
50	Low levels of CD36 in peripheral blood monocytes in subclinical atherosclerosis in rheumatoid arthritis: a cross-sectional study in a Mexican population. <i>BioMed Research International</i> , 2014 , 2014, 736786	3	7
49	Circulating E-selectin and tumor necrosis factor-alpha in extraarticular involvement and joint disease activity in rheumatoid arthritis. <i>Rheumatology International</i> , 2009 , 29, 281-6	3.6	7
48	Polymorphism of the beta3-adrenergic receptor and lipid profile in patients with rheumatoid arthritis and systemic lupus erythematosus treated with chloroquine. <i>Rheumatology International</i> , 2003 , 23, 99-103	3.6	7
47	CCL2 Serum Levels and Adiposity Are Associated with the Polymorphic Phenotypes -2518A on CCL2 and 641LE on CCR2 in a Mexican Population with Insulin Resistance. <i>Journal of Diabetes Research</i> , 2016 , 2016, 5675739	3.9	7
46	Subclinical parameters of arterial stiffness and arteriosclerosis correlate with QRISK3 in systemic lupus erythematosus. <i>PLoS ONE</i> , 2018 , 13, e0207520	3.7	7
45	Prescription for antiresorptive therapy in Mexican patients with rheumatoid arthritis: is it time to reevaluate the strategies for osteoporosis prevention?. <i>Rheumatology International</i> , 2013 , 33, 145-50	3.6	6
44	FCGR3A V(176) polymorphism for systemic lupus erythematosus susceptibility in Mexican population. <i>Rheumatology International</i> , 2011 , 31, 1065-8	3.6	6
43	A new PCR-RFLP assay for -1123 G>C polymorphism in the PTPN22 gene: allele and genotype frequencies in a western Mexican population. <i>Clinical Chemistry and Laboratory Medicine</i> , 2009 , 47, 491-3 ^{5.9}		6
42	IgA antihistone antibodies in isoniazid-treated tuberculosis patients. <i>Autoimmunity</i> , 1995 , 20, 105-11	3	6
41	The distribution of CD56(dim) CD16+ and CD56(bright) CD16- cells are associated with prolactin levels during pregnancy and menstrual cycle in healthy women. <i>American Journal of Reproductive Immunology</i> , 2011 , 65, 433-7	3.8	5
40	Plasminogen activator inhibitor-1 C/G polymorphism in relation to plasma levels in rheumatoid arthritis. <i>Clinical and Experimental Medicine</i> , 2009 , 9, 223-8	4.9	5
39	Disease Activity Score on 28 Joints and Polypharmacy Are Independent Predictors for Health-Related Quality of Life Evaluated by INCAVISA in Patients With Rheumatoid Arthritis. <i>Journal of Clinical Rheumatology</i> , 2016 , 22, 399-404	1.1	5

38	sE-Selectin expression and A561C polymorphism in relation to rheumatoid arthritis clinical activity. <i>Journal of Rheumatology</i> , 2006 , 33, 1968-72	4.1	5
37	Potential Chronotherapeutic Optimization of Antimalarials in Systemic Lupus Erythematosus: Is Toll-Like Receptor 9 Expression Dependent on the Circadian Cycle in Humans?. <i>Frontiers in Immunology</i> , 2018 , 9, 1497	8.4	4
36	Plasminogen activator inhibitor-1 polymorphisms (-844 G>A and HindIII C>G) in systemic lupus erythematosus: association with clinical variables. <i>Clinical and Experimental Medicine</i> , 2011 , 11, 11-7	4.9	4
35	The -383A>C TNFRI polymorphism is associated with soluble levels and clinical activity in rheumatoid arthritis. <i>Rheumatology International</i> , 2010 , 30, 655-9	3.6	4
34	Evaluation of lipid profile, macular toxicity and clinical manifestations according to APO E genotype in systemic lupus erythematosus and rheumatoid arthritis patients treated with chloroquine. <i>Scandinavian Journal of Rheumatology</i> , 2002 , 31, 32-7	1.9	4
33	Procollagen Type I and III Aminoterminal Propeptide Levels and Severity of Interstitial Lung Disease in Mexican Women With Progressive Systemic Sclerosis. <i>Archivos De Bronconeumologia</i> , 2015 , 51, 440-448	8.7	3
32	Increased CD28 serum levels are not associated with specific clinical activity in systemic lupus erythematosus. <i>Rheumatology International</i> , 2011 , 31, 1321-1324	3.6	3
31	Placental cytokine expression and NF-kappaB activity in women with rheumatic diseases. <i>Scandinavian Journal of Rheumatology</i> , 2006 , 35, 160-2	1.9	3
30	Myositis-Specific Antibodies and Myositis-Associated Antibodies in Patients With Idiopathic Inflammatory Myopathies From the PANLAR Myositis Study Group. <i>Journal of Clinical Rheumatology</i> , 2021 , 27, e302-e306	1.1	3
29	Enalapril Influence on Arterial Stiffness in Rheumatoid Arthritis Women: A Randomized Clinical Trial. <i>Frontiers in Medicine</i> , 2019 , 6, 341	4.9	2
28	Influenza Vaccine and Autoimmune Diseases 2015 , 175-184		2
27	Coexistence of anti-RNA polymerase III and anti-U1RNP antibodies in patients with systemic lupus erythematosus: two cases without features of scleroderma. <i>Lupus</i> , 2012 , 21, 68-74	2.6	2
26	Lipid profile and antibodies against oxidized low density lipoprotein in alcoholic cirrhosis. <i>Hepatology Research</i> , 1999 , 15, 95-103	5.1	2
25	IgG Anti-ghrelin Immune Complexes Are Increased in Rheumatoid Arthritis Patients Under Biologic Therapy and Are Related to Clinical and Metabolic Markers. <i>Frontiers in Endocrinology</i> , 2019 , 10, 252	5.7	1
24	Adenovirus Biodistribution is Modified in Sensitive Animals Compared to Naïve Animals. <i>Molecular Biotechnology</i> , 2020 , 62, 260-272	3	1
23	Response to: Qantisynthetase syndrome or what else? Different perspectives indicate the need for new classification criteriaQy Cavagna. <i>Annals of the Rheumatic Diseases</i> , 2018 , 77, e51	2.4	1
22	The DNA co-vaccination using Sm antigen and IL-10 as prophylactic experimental therapy ameliorates nephritis in a model of lupus induced by pristane. <i>PLoS ONE</i> , 2021 , 16, e0259114	3.7	1
21	Geographical Latitude Remains as an Important Factor for the Prevalence of Some Myositis Autoantibodies: A Systematic Review. <i>Frontiers in Immunology</i> , 2021 , 12, 672008	8.4	1

20	Low CD36 and LOX-1 Levels and CD36 Gene Subexpression Are Associated with Metabolic Dysregulation in Older Individuals with Abdominal Obesity. <i>Journal of Diabetes Research</i> , 2016 , 2016, 5678946	3.9	1
19	Evaluation of medical ethics competencies in rheumatology: local experience during national accreditation process. <i>Journal of Medical Ethics</i> , 2019 , 45, 839-842	2.5	1
18	Extrapulmonary Tuberculosis Infection in Mexican Patients With Idiopathic Inflammatory Myopathies. <i>Journal of Clinical Rheumatology</i> , 2015 , 21, 331-2	1.1	0
17	Going Further: Comprehensive Disease Control of Rheumatoid Arthritis, Targeting Cytokines and Chemokines. <i>Journal of Clinical Rheumatology</i> , 2021 , 27, e432-e439	1.1	0
16	Evaluaci3n cl3nica y ultrasonogr3fica de la gl3ndula tiroides en pacientes con artritis reumatoide. <i>Gaceta Medica De Mexico</i> , 2018 , 154, 432-437	0.3	0
15	Presence of anti-TIF-1 anti-Ro52, anti-SSA/Ro60 and anti-Su/Ago2 antibodies in breast cancer: a cross-sectional study. <i>Immunopharmacology and Immunotoxicology</i> , 2021 , 43, 328-333	3.2	0
14	Contribution of rs3211938 polymorphism at CD36 to glucose levels, oxidized low-density lipoproteins, insulin resistance, and body mass index in Mexican mestizos with type-2 diabetes from western Mexico. <i>Nutricion Hospitalaria</i> , 2021 , 38, 742-748	1	0
13	CCR2/CCL2 and CMKLR1/RVE1 chemokines system levels are associated with insulin resistance in rheumatoid arthritis. <i>PLoS ONE</i> , 2021 , 16, e0246054	3.7	0
12	Osteopontin: another piece in the systemic lupus erythematosus immunopathology puzzle. <i>Clinical and Experimental Rheumatology</i> , 2022 , 40, 173-182	2.2	0
11	Good response to mycophenolate mofetil on treatment of interstitial lung disease in polymyositis associated with antisynthetase syndrome positive for anti-EJ and anti-Ro52 antibodies. <i>Clinical Rheumatology</i> , 2020 , 39, 2837-2839	3.9	
10	AB0029 Relationship SCD36 Levels and Disease Activity in Patients with Rheumatoid Arthritis. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 906.3-907	2.4	
9	AB0635 Clinical and Autoantibodies Features in Autoimmune/Inflammatory Syndrome Induced by Adjuvants: Mineral Oils and Silicone. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 1015.3-1016	2.4	
8	FRI0081 Titers of Anti-Ccp Antibodies, IL-6, Tnf-Alpha and CRP Are Associated with Increased Carotid Intima-Media Thickness: Cross-Sectional Analysis of A Rheumatoid Arthritis Group without Cardiovascular Risk Factors. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 410.3-411	2.4	
7	SAT0135 Low Levels of CD36 in Peripheral Blood Monocytes in Subclinical Atherosclerosis in Rheumatoid Arthritis: A Cross-Sectional Study of A Mexican Population. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 639.2-640	2.4	
6	AB0371 Lepr 668 A>g Snp is Associated with Titers of Anti-CCP Antibodies and Erythrocyte Sedimentation Rate in Mexican Mestizo Patients with Rheumatoid Arthritis. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 1017.2-1017	2.4	
5	AB0936 Tuberculosis Infection in Mexican Patients with Idiopathic Inflammatory Myopathies: Report of three Cases and Review of the Literature. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 1212.3-1212	2.4	
4	AB0381 Clinical and Ultrasound Assessment of Thyroid in Rheumatoid Arthritis Mexican Patients. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 1020.2-1020	2.4	
3	Influence of Anti-TNF and Disease Modifying Antirheumatic Drugs Therapy on Pulmonary Forced Vital Capacity Associated to Ankylosing Spondylitis: A 2-Year Follow-Up Observational Study. <i>Journal of Immunology Research</i> , 2015 , 2015, 980147	4.5	

- 2 Caspase 3 blocking avoids the expression of autoantigens triggered by apoptosis in neonatal Balb/c mice skin. *Reumatismo*, **2011**, 63, 29-37 1.1
- 1 Sensorineural Hearing Impairment and Subclinical Atherosclerosis in Rheumatoid Arthritis Patients Without Traditional Cardiovascular Risk Factors. *Archives of Rheumatology*, **2016**, 31, 208-214 0.9