Xiongwen Chen

List of Publications by Citations

Source: https://exaly.com/author-pdf/4229491/xiongwen-chen-publications-by-citations.pdf

Version: 2024-04-19

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

97
papers

5,824
h-index

75
g-index

114
ext. papers

6,754
ext. citations

9
avg, IF

L-index

#	Paper	IF	Citations
97	A peptide encoded by a transcript annotated as long noncoding RNA enhances SERCA activity in muscle. <i>Science</i> , 2016 , 351, 271-5	33.3	439
96	Cellular basis of abnormal calcium transients of failing human ventricular myocytes. <i>Circulation Research</i> , 2003 , 92, 651-8	15.7	363
95	Ca2+- and mitochondrial-dependent cardiomyocyte necrosis as a primary mediator of heart failure. Journal of Clinical Investigation, 2007, 117, 2431-44	15.9	317
94	Bone marrow cells adopt the cardiomyogenic fate in vivo. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007 , 104, 17783-8	11.5	261
93	L-type Ca2+ channel density and regulation are altered in failing human ventricular myocytes and recover after support with mechanical assist devices. <i>Circulation Research</i> , 2002 , 91, 517-24	15.7	229
92	The mitochondrial Na/Ca exchanger is essential for Ca homeostasis and viability. <i>Nature</i> , 2017 , 545, 93-9	₹ 0.4	203
91	G protein-coupled receptor kinase 2 ablation in cardiac myocytes before or after myocardial infarction prevents heart failure. <i>Circulation Research</i> , 2008 , 103, 413-22	15.7	186
90	Ca2+ influx-induced sarcoplasmic reticulum Ca2+ overload causes mitochondrial-dependent apoptosis in ventricular myocytes. <i>Circulation Research</i> , 2005 , 97, 1009-17	15.7	154
89	GDF11 does not rescue aging-related pathological hypertrophy. Circulation Research, 2015, 117, 926-32	15.7	124
88	Decreased cardiac L-type Call+ channel activity induces hypertrophy and heart failure in mice. <i>Journal of Clinical Investigation</i> , 2012 , 122, 280-90	15.9	123
87	Plasma long non-coding RNA, CoroMarker, a novel biomarker for diagnosis of coronary artery disease. <i>Clinical Science</i> , 2015 , 129, 675-85	6.5	119
86	How to Improve the Survival of Transplanted Mesenchymal Stem Cell in Ischemic Heart?. <i>Stem Cells International</i> , 2016 , 2016, 9682757	5	115
85	Protein kinase C{alpha}, but not PKC{beta} or PKC{gamma}, regulates contractility and heart failure susceptibility: implications for ruboxistaurin as a novel therapeutic approach. <i>Circulation Research</i> , 2009 , 105, 194-200	15.7	113
84	CaMKII negatively regulates calcineurin-NFAT signaling in cardiac myocytes. <i>Circulation Research</i> , 2009 , 105, 316-25	15.7	104
83	Cardiotoxic and cardioprotective features of chronic 🛭 adrenergic signaling. <i>Circulation Research</i> , 2013 , 112, 498-509	15.7	102
82	Adolescent feline heart contains a population of small, proliferative ventricular myocytes with immature physiological properties. <i>Circulation Research</i> , 2007 , 100, 536-44	15.7	102
81	Dedifferentiation, Proliferation, and Redifferentiation of Adult Mammalian Cardiomyocytes After Ischemic Injury. <i>Circulation</i> , 2017 , 136, 834-848	16.7	101

(2008-2016)

80	Crucial Role of miR-433 in Regulating Cardiac Fibrosis. <i>Theranostics</i> , 2016 , 6, 2068-2083	12.1	100
79	Paroxetine-mediated GRK2 inhibition reverses cardiac dysfunction and remodeling after myocardial infarction. <i>Science Translational Medicine</i> , 2015 , 7, 277ra31	17.5	99
78	Calcium influx through Cav1.2 is a proximal signal for pathological cardiomyocyte hypertrophy. Journal of Molecular and Cellular Cardiology, 2011 , 50, 460-70	5.8	88
77	Transient receptor potential channels contribute to pathological structural and functional remodeling after myocardial infarction. <i>Circulation Research</i> , 2014 , 115, 567-580	15.7	84
76	L-type Ca2+ channel alpha 1c subunit isoform switching in failing human ventricular myocardium. Journal of Molecular and Cellular Cardiology, 2000 , 32, 973-84	5.8	82
75	LETM1-dependent mitochondrial Ca2+ flux modulates cellular bioenergetics and proliferation. <i>FASEB Journal</i> , 2014 , 28, 4936-49	0.9	8o
74	Prolyl hydroxylase domain protein 2 silencing enhances the survival and paracrine function of transplanted adipose-derived stem cells in infarcted myocardium. <i>Circulation Research</i> , 2013 , 113, 288-3	66.7	77
73	Adrenergic regulation of cardiac contractility does not involve phosphorylation of the cardiac ryanodine receptor at serine 2808. <i>Circulation Research</i> , 2008 , 102, e65-72	15.7	77
72	STIM1 elevation in the heart results in aberrant Call+ handling and cardiomyopathy. <i>Journal of Molecular and Cellular Cardiology</i> , 2015 , 87, 38-47	5.8	76
71	Hyperphosphorylation of the cardiac ryanodine receptor at serine 2808 is not involved in cardiac dysfunction after myocardial infarction. <i>Circulation Research</i> , 2012 , 110, 831-40	15.7	75
70	alpha1G-dependent T-type Ca2+ current antagonizes cardiac hypertrophy through a NOS3-dependent mechanism in mice. <i>Journal of Clinical Investigation</i> , 2009 , 119, 3787-96	15.9	75
69	Circulating RncRNA OTTHUMT00000387022Rfrom monocytes as a novel biomarker for coronary artery disease. <i>Cardiovascular Research</i> , 2016 , 112, 714-724	9.9	74
68	Mitochondrial DNA oxidative damage contributes to cardiomyocyte ischemia/reperfusion-injury in rats: cardioprotective role of lycopene. <i>Journal of Cellular Physiology</i> , 2015 , 230, 2128-41	7	72
67	Repair of the injured adult heart involves new myocytes potentially derived from resident cardiac stem cells. <i>Circulation Research</i> , 2011 , 108, 1226-37	15.7	72
66	Alterations in early action potential repolarization causes localized failure of sarcoplasmic reticulum Ca2+ release. <i>Circulation Research</i> , 2005 , 96, 543-50	15.7	69
65	Targeting HO-1 by Epigallocatechin-3-Gallate Reduces Contrast-Induced Renal Injury via Anti-Oxidative Stress and Anti-Inflammation Pathways. <i>PLoS ONE</i> , 2016 , 11, e0149032	3.7	66
64	Neonatal Transplantation Confers Maturation of PSC-Derived Cardiomyocytes Conducive to Modeling Cardiomyopathy. <i>Cell Reports</i> , 2017 , 18, 571-582	10.6	63
63	Ca2+ influx through T- and L-type Ca2+ channels have different effects on myocyte contractility and induce unique cardiac phenotypes. <i>Circulation Research</i> , 2008 , 103, 1109-19	15.7	63

62	Insulin inhibits cardiac contractility by inducing a Gi-biased [2-adrenergic signaling in hearts. <i>Diabetes</i> , 2014 , 63, 2676-89	0.9	60
61	Activation of Cannabinoid Receptor 2 Ameliorates DSS-Induced Colitis through Inhibiting NLRP3 Inflammasome in Macrophages. <i>PLoS ONE</i> , 2016 , 11, e0155076	3.7	57
60	Phosphodiesterases coordinate cAMP propagation induced by two stimulatory G protein-coupled receptors in hearts. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012 , 109, 6578-83	11.5	56
59	Pim-1 kinase antagonizes aspects of myocardial hypertrophy and compensation to pathological pressure overload. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008 , 105, 13889-94	11.5	55
58	Regulated overexpression of the A1-adenosine receptor in mice results in adverse but reversible changes in cardiac morphology and function. <i>Circulation</i> , 2006 , 114, 2240-50	16.7	53
57	Negative inotropic effects of high-mobility group box 1 protein in isolated contracting cardiac myocytes. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2008 , 294, H1490-6	5.2	48
56	GSK-3alpha directly regulates beta-adrenergic signaling and the response of the heart to hemodynamic stress in mice. <i>Journal of Clinical Investigation</i> , 2010 , 120, 2280-91	15.9	44
55	Therapeutic effect of a novel Wnt pathway inhibitor on cardiac regeneration after myocardial infarction. <i>Clinical Science</i> , 2017 , 131, 2919-2932	6.5	38
54	Analyses of caspase-1-regulated transcriptomes in various tissues lead to identification of novel IL-1\(\Pri\), IL-18- and sirtuin-1-independent pathways. <i>Journal of Hematology and Oncology</i> , 2017 , 10, 40	22.4	35
53	Inflammation and cardiac dysfunction during sepsis, muscular dystrophy, and myocarditis. <i>Burns and Trauma</i> , 2013 , 1, 109-21		35
52	Increasing cardiac contractility after myocardial infarction exacerbates cardiac injury and pump dysfunction. <i>Circulation Research</i> , 2010 , 107, 800-9	15.7	35
51	Autophagy Plays an Important Role in Anti-inflammatory Mechanisms Stimulated by Alpha7 Nicotinic Acetylcholine Receptor. <i>Frontiers in Immunology</i> , 2017 , 8, 553	8.4	33
50	Potential of cardiac stem/progenitor cells and induced pluripotent stem cells for cardiac repair in ischaemic heart disease. <i>Clinical Science</i> , 2013 , 125, 319-27	6.5	31
49	Inhibition of type 5 phosphodiesterase counteracts [2-adrenergic signalling in beating cardiomyocytes. <i>Cardiovascular Research</i> , 2015 , 106, 408-20	9.9	29
48	A cellular mechanism of muscle memory facilitates mitochondrial remodelling following resistance training. <i>Journal of Physiology</i> , 2018 , 596, 4413-4426	3.9	29
47	{beta}1-Adrenergic receptor activation induces mouse cardiac myocyte death through both L-type calcium channel-dependent and -independent pathways. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2010 , 299, H322-31	5.2	29
46	Cardiac G-protein-coupled receptor kinase 2 ablation induces a novel Ca2+ handling phenotype resistant to adverse alterations and remodeling after myocardial infarction. <i>Circulation</i> , 2012 , 125, 2108	-167 -187	29
45	Role of STIM1 (Stromal Interaction Molecule 1) in Hypertrophy-Related Contractile Dysfunction. <i>Circulation Research</i> , 2017 , 121, 125-136	15.7	27

(2013-2014)

44	Blunted cardiac beta-adrenergic response as an early indication of cardiac dysfunction in Duchenne muscular dystrophy. <i>Cardiovascular Research</i> , 2014 , 103, 60-71	9.9	27
43	Circulating "LncPPAR" From Monocytes as a Novel Biomarker for Coronary Artery Diseases. <i>Medicine (United States)</i> , 2016 , 95, e2360	1.8	27
42	A novel porcupine inhibitor blocks WNT pathways and attenuates cardiac hypertrophy. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2018 , 1864, 3459-3467	6.9	24
41	□Adrenergic stimulation increases Cav3.1 activity in cardiac myocytes through protein kinase A. <i>PLoS ONE</i> , 2012 , 7, e39965	3.7	24
40	Protease-activated receptor 4 deficiency offers cardioprotection after acute ischemia reperfusion injury. <i>Journal of Molecular and Cellular Cardiology</i> , 2016 , 90, 21-9	5.8	23
39	The A isoform of calmodulin kinase II mediates pathological cardiac hypertrophy by interfering with the HDAC4-MEF2 signaling pathway. <i>Biochemical and Biophysical Research Communications</i> , 2011 , 409, 125-30	3.4	22
38	Reduced effects of BAY K 8644 on L-type Ca2+ current in failing human cardiac myocytes are related to abnormal adrenergic regulation. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2008 , 294, H2257-67	5.2	22
37	Cardiomyocyte PKA Ablation Enhances Basal Contractility While Eliminates Cardiac I-Adrenergic Response Without Adverse Effects on the Heart. <i>Circulation Research</i> , 2019 , 124, 1760-1777	15.7	21
36	Enhanced basal contractility but reduced excitation-contraction coupling efficiency and beta-adrenergic reserve of hearts with increased Cav1.2 activity. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2010 , 299, H519-28	5.2	21
35	Gastrin Protects Against Myocardial Ischemia/Reperfusion Injury via Activation of RISK (Reperfusion Injury Salvage Kinase) and SAFE (Survivor Activating Factor Enhancement) Pathways. <i>Journal of the American Heart Association</i> , 2018 , 7,	6	20
34	Protein kinase D3 is a pivotal activator of pathological cardiac hypertrophy by selectively increasing the expression of hypertrophic transcription factors. <i>Journal of Biological Chemistry</i> , 2011 , 286, 40782-9	⊅ ·4	20
33	Calcineurin inhibition normalizes beta-adrenergic responsiveness in the spontaneously hypertensive rat. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2007 , 293, H3122-9	5.2	19
32	Interaction of the Joining Region in Junctophilin-2 With the L-Type Ca Channel Is Pivotal for Cardiac Dyad Assembly and Intracellular Ca Dynamics. <i>Circulation Research</i> , 2021 , 128, 92-114	15.7	19
31	Persistent increases in Ca(2+) influx through Cav1.2 shortens action potential and causes Ca(2+) overload-induced afterdepolarizations and arrhythmias. <i>Basic Research in Cardiology</i> , 2016 , 111, 4	11.8	18
30	Increased expression of Syne1/nesprin-1 facilitates nuclear envelope structure changes in embryonic stem cell differentiation. <i>Developmental Dynamics</i> , 2011 , 240, 2245-55	2.9	18
29	Caveolae-localized L-type Ca2+ channels do not contribute to function or hypertrophic signalling in the mouse heart. <i>Cardiovascular Research</i> , 2017 , 113, 749-759	9.9	17
28	Fatty Acid Oxidation Promotes Cardiomyocyte Proliferation Rate but Does Not Change Cardiomyocyte Number in Infant Mice. <i>Frontiers in Cell and Developmental Biology</i> , 2019 , 7, 42	5.7	17
27	Compartmentalization of 🛭 adrenergic signals in cardiomyocytes. <i>Trends in Cardiovascular Medicine</i> , 2013 , 23, 250-6	6.9	16

26	Evidence for K+-dependent HCO3- utilization in the marine diatom Phaeodactylum tricornutum. <i>Plant Physiology</i> , 2006 , 141, 731-6	6.6	16
25	Intracellular sodium determines frequency-dependent alterations in contractility in hypertrophied feline ventricular myocytes. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2007 , 292, H1129-38	5.2	15
24	T-type Call+ channels regulate the exit of cardiac myocytes from the cell cycle after birth. <i>Journal of Molecular and Cellular Cardiology</i> , 2013 , 62, 122-30	5.8	14
23	Regulation of L-type calcium channel by phospholemman in cardiac myocytes. <i>Journal of Molecular and Cellular Cardiology</i> , 2015 , 84, 104-11	5.8	13
22	Protein Kinase Inhibitor Peptide as a Tool to Specifically Inhibit Protein Kinase A. <i>Frontiers in Physiology</i> , 2020 , 11, 574030	4.6	12
21	Increasing T-type calcium channel activity by Padrenergic stimulation contributes to Padrenergic regulation of heart rates. <i>Journal of Physiology</i> , 2018 , 596, 1137-1151	3.9	10
20	Cardiac troponin I exacerbates myocardial ischaemia/reperfusion injury by inducing the adhesion of monocytes to vascular endothelial cells via a TLR4/NF- B -dependent pathway. <i>Clinical Science</i> , 2016 , 130, 2279-2293	6.5	10
19	Metformin promotes the survival of transplanted cardiosphere-derived cells thereby enhancing their therapeutic effect against myocardial infarction. <i>Stem Cell Research and Therapy</i> , 2017 , 8, 17	8.3	10
18	Reduced Myocardial Reserve in Young X-Linked Muscular Dystrophy Mice Diagnosed by Two-Dimensional Strain Analysis Combined with Stress Echocardiography. <i>Journal of the American Society of Echocardiography</i> , 2017 , 30, 815-827.e9	5.8	8
17	Combined administration of anisodamine and neostigmine rescued acute lethal crush syndrome through InAChR-dependent JAK2-STAT3 signaling. <i>Scientific Reports</i> , 2016 , 6, 37709	4.9	7
16	Cross talk between serine/threonine and tyrosine kinases regulates ADP-induced thromboxane generation in platelets. <i>Thrombosis and Haemostasis</i> , 2015 , 114, 558-68	7	7
15	miR-301a-PTEN-AKT Signaling Induces Cardiomyocyte Proliferation and Promotes Cardiac Repair Post-MI. <i>Molecular Therapy - Nucleic Acids</i> , 2020 , 22, 251-262	10.7	7
14	Syngeneic AAV Pseudo-particles Potentiate Gene Transduction of AAV Vectors. <i>Molecular Therapy - Methods and Clinical Development</i> , 2017 , 4, 149-158	6.4	6
13	Inhibition of angiotensin II Gq signaling augments beta-adrenergic receptor mediated effects in a renal artery stenosis model of high blood pressure. <i>Journal of Molecular and Cellular Cardiology</i> , 2009 , 46, 100-7	5.8	6
12	Preclinical cardiovascular changes in children with obesity: A real-time 3-dimensional speckle tracking imaging study. <i>PLoS ONE</i> , 2018 , 13, e0205177	3.7	6
11	Phosphorus limitation for the colony formation, growth and photosynthesis of an edible cyanobacterium, Nostoc sphaeroides. <i>Biotechnology Letters</i> , 2012 , 34, 137-43	3	5
10	LIGHT DEPENDENCY OF PHOTOSYNTHETIC RECOVERY DURING WETTING AND THE ACCLIMATION OF PHOTOSYNTHETIC APPARATUS TO LIGHT FLUCTUATION IN A TERRESTRIAL CYANOBACTERIUM NOSTOC COMMUNE(1). <i>Journal of Phycology</i> , 2011 , 47, 1063-71	3	4
9	Remodeling of repolarization and arrhythmia susceptibility in a myosin-binding protein C knockout mouse model. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2017 , 313, H620-H630	5.2	3

LIST OF PUBLICATIONS

8	Effect of Clibn photosynthetic bicarbonate uptake in two cyanobacteria Microcystis aeruginosa and Synechocystis PCC6803. <i>Science Bulletin</i> , 2009 , 54, 1197-1203	10.6	2
7	Physiological profiles associated with ceasing growth of unfertilized eggs produced by unmated queens in the subterranean termite Reticulitermes chinensis. <i>Biology Open</i> , 2016 , 5, 756-63	2.2	2
6	Hepatic PKA inhibition accelerates the lipid accumulation in liver. <i>Nutrition and Metabolism</i> , 2019 , 16, 69	4.6	1
5	Bazedoxifene Regulates Th17 Immune Response to Ameliorate Experimental Autoimmune myocarditis via Inhibition of STAT3 Activation. <i>Frontiers in Pharmacology</i> , 2020 , 11, 613160	5.6	1
4	A low voltage activated Ca current found in a subset of human ventricular myocytes. <i>Channels</i> , 2020 , 14, 231-245	3	О
3	Phenotype and Function of c-kit+-derived Amplifying Myocytes 2007 , 77-85		
2	Pharmacology of L-Type and T-Type Calcium Channels in the Heart 2004 , 133-142		
1	Research on the Expression and Regulatory Mechanism of Breast Cancer Susceptibility Gene-1 on Cell of Skin Cancer in Different Classification. <i>Journal of Biomaterials and Tissue Engineering</i> , 2021 , 11, 756-760	0.3	