Stephen R Wisniewski

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/4252540/stephen-r-wisniewski-publications-by-year.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

180 367 35,490 91 h-index g-index citations papers 6.2 6.5 40,037 370 L-index avg, IF ext. citations ext. papers

#	Paper	IF	Citations
367	Comparison of Intracranial Pressure Measurements Before and After Hypertonic Saline or Mannitol Treatment in Children With Severe Traumatic Brain Injury <i>JAMA Network Open</i> , 2022 , 5, e220891	10.4	3
366	Assessing Unexpected Circumstances That Lead to Modifications in Clinical Trial Design, Conduct, or Analysis: Application of the CONSERVE Reporting Guideline <i>JAMA - Journal of the American Medical Association</i> , 2022 , 327, 868-869	27.4	
365	Use of Antifibrinolytics in Pediatric Life-Threatening Hemorrhage: A Prospective Observational Multicenter Study. <i>Critical Care Medicine</i> , 2021 ,	1.4	2
364	Characteristics Associated with the Use of an Inpatient Rehabilitation or Skilled Nursing Facility after Acute Care in Children with Severe Traumatic Brain Injury. <i>Developmental Neurorehabilitation</i> , 2021 , 24, 466-477	1.8	1
363	Association of Pharmacological Interventions With Symptom Burden Reduction in Patients With Mild Traumatic Brain Injury: A Systematic Review. <i>JAMA Neurology</i> , 2021 , 78, 596-608	17.2	3
362	Effect of Antimicrobial Therapy on Respiratory Hospitalization or Death in Adults With Idiopathic Pulmonary Fibrosis: The CleanUP-IPF Randomized Clinical Trial. <i>JAMA - Journal of the American Medical Association</i> , 2021 , 325, 1841-1851	27.4	12
361	Life-Threatening Bleeding in Children: A Prospective Observational Study. <i>Critical Care Medicine</i> , 2021 , 49, 1943-1954	1.4	9
360	Cerebrospinal Fluid Sulfonylurea Receptor-1 is Associated with Intracranial Pressure and Outcome after Pediatric TBI: An Exploratory Analysis of the Cool Kids Trial. <i>Journal of Neurotrauma</i> , 2021 , 38, 16	1 <i>5</i> : 1 61	9 ⁶
359	Effect of vitamin D supplementation on total and allergen-specific IgE in children with asthma and low vitamin D levels. <i>Journal of Allergy and Clinical Immunology</i> , 2021 ,	11.5	2
358	Gene coexpression networks reveal novel molecular endotypes in alpha-1 antitrypsin deficiency. <i>Thorax</i> , 2021 , 76, 134-143	7.3	3
357	Survey to inform trial of low-titer group O whole-blood compared to conventional blood components for children with severe traumatic bleeding. <i>Transfusion</i> , 2021 , 61 Suppl 1, S43-S48	2.9	O
356	Vitamin D supplementation, lung function and asthma control in children with asthma and low vitamin D levels. <i>European Respiratory Journal</i> , 2021 , 58,	13.6	1
355	Violence-related distress and lung function in two longitudinal studies of youth. <i>European Respiratory Journal</i> , 2021 ,	13.6	2
354	Multi-omic analysis in injured humans: Patterns align with outcomes and treatment responses <i>Cell Reports Medicine</i> , 2021 , 2, 100478	18	4
353	Improving Cost-effectiveness and Access to Cognitive Behavior Therapy for Depression: Providing Remote-Ready, Computer-Assisted Psychotherapy in Times of Crisis and Beyond. <i>Psychotherapy and Psychosomatics</i> , 2020 , 89, 307-313	9.4	15
352	Effects of Collaborative Care for Comorbid Attention Deficit Hyperactivity Disorder Among Children With Behavior Problems in Pediatric Primary Care. <i>Clinical Pediatrics</i> , 2020 , 59, 787-800	1.2	1
351	Design and rationale of a multi-center, pragmatic, open-label randomized trial of antimicrobial therapy - the study of clinical efficacy of antimicrobial therapy strategy using pragmatic design in Idiopathic Pulmonary Fibrosis (CleanUP-IPF) clinical trial. <i>Respiratory Research</i> , 2020 , 21, 68	7.3	15

(2017-2020)

350	Effect of Vitamin D3 Supplementation on Severe Asthma Exacerbations in Children With Asthma and Low Vitamin D Levels: The VDKA Randomized Clinical Trial. <i>JAMA - Journal of the American Medical Association</i> , 2020 , 324, 752-760	27.4	36	
349	Paths to Successful Translation of New Therapies for Severe Traumatic Brain Injury in the Golden Age of Traumatic Brain Injury Research: A Pittsburgh Vision. <i>Journal of Neurotrauma</i> , 2020 , 37, 2353-23	37∮· ⁴	15	
348	Practice Patterns in Pharmacological and Non-Pharmacological Therapies for Children with Mild Traumatic Brain Injury: A Survey of 15 Canadian and United States Centers. <i>Journal of Neurotrauma</i> , 2019 , 36, 2886-2894	5.4	5	
347	Use of magnetic resonance imaging in severe pediatric traumatic brain injury: assessment of current practice. <i>Journal of Neurosurgery: Pediatrics</i> , 2019 , 23, 471-479	2.1	12	
346	Age and Mortality in Pediatric Severe Traumatic Brain Injury: Results from an International Study. <i>Neurocritical Care</i> , 2018 , 28, 302-313	3.3	22	
345	The Rapid Escalation of Fentanyl in Illicit Drug Evidence in Allegheny County, Pennsylvania, 2010-2016. <i>Public Health Reports</i> , 2018 , 133, 142-146	2.5	7	
344	Initiating Nutritional Support Before 72 Hours Is Associated With Favorable Outcome After Severe Traumatic Brain Injury in Children: A Secondary Analysis of a Randomized, Controlled Trial of Therapeutic Hypothermia. <i>Pediatric Critical Care Medicine</i> , 2018 , 19, 345-352	3	14	
343	Adjunctive Bright Light Therapy for Bipolar Depression: A Randomized Double-Blind Placebo-Controlled Trial. <i>American Journal of Psychiatry</i> , 2018 , 175, 131-139	11.9	86	
342	Improving the Efficiency of Psychotherapy for Depression: Computer-Assisted Versus Standard CBT. <i>American Journal of Psychiatry</i> , 2018 , 175, 242-250	11.9	48	
341	AF-CBT for Families Experiencing Physical Aggression or Abuse Served by the Mental Health or Child Welfare System: An Effectiveness Trial. <i>Child Maltreatment</i> , 2018 , 23, 319-333	2.8	12	
340	Presenting Characteristics Associated With Outcome in Children With Severe Traumatic Brain Injury: A Secondary Analysis From a Randomized, Controlled Trial of Therapeutic Hypothermia. <i>Pediatric Critical Care Medicine</i> , 2018 , 19, 957-964	3	8	
339	Tripartite Stratification of the Glasgow Coma Scale in Children with Severe Traumatic Brain Injury and Mortality: An Analysis from a Multi-Center Comparative Effectiveness Study. <i>Journal of Neurotrauma</i> , 2017 , 34, 2220-2229	5.4	21	
338	Effect of aerosolized antipseudomonals on Pseudomonas positivity and bronchiolitis obliterans syndrome after lung transplantation. <i>Transplant Infectious Disease</i> , 2017 , 19, e12688	2.7	5	
337	Abusive Head Trauma and Mortality-An Analysis From an International Comparative Effectiveness Study of Children With Severe Traumatic Brain Injury. <i>Critical Care Medicine</i> , 2017 , 45, 1398-1407	1.4	30	
336	Correlates of Postpartum Maternal Functioning in a Low-Income Obstetric Population. <i>Journal of the American Psychiatric Nurses Association</i> , 2017 , 23, 149-158	1.8	21	
335	Rescue alemtuzumab for refractory acute cellular rejection and bronchiolitis obliterans syndrome after lung transplantation. <i>Clinical Transplantation</i> , 2017 , 31, e12899	3.8	14	
334	Generating an efficient version of the Edinburgh Postnatal Depression Scale in an urban obstetrical population. <i>Journal of Affective Disorders</i> , 2017 , 208, 615-620	6.6	13	
333	Challenges and opportunities for pediatric severe TBI-review of the evidence and exploring a way forward. <i>Childis Nervous System</i> , 2017 , 33, 1663-1667	1.7	23	

332	Mood symptoms in pregnant and postpartum women with bipolar disorder: a naturalistic study. <i>Bipolar Disorders</i> , 2017 , 19, 295-304	3.8	11
331	Cost-Effective Drug Switch Options After Unsuccessful Treatment With an SSRI for Depression. <i>Psychiatric Services</i> , 2017 , 68, 81-87	3.3	1
330	Reply. Arthritis and Rheumatology, 2017 , 69, 242	9.5	
329	Are Patients with Childhood Onset of Insomnia and Depression More Difficult to Treat Than Are Those with Adult Onsets of These Disorders? A Report from the TRIAD Study. <i>Journal of Clinical Sleep Medicine</i> , 2017 , 13, 205-213	3.1	5
328	Phase I randomized clinical trial of N-acetylcysteine in combination with an adjuvant probenecid for treatment of severe traumatic brain injury in children. <i>PLoS ONE</i> , 2017 , 12, e0180280	3.7	31
327	One-Year Developmental Outcomes for Infants of Mothers With Bipolar Disorder. <i>Journal of Clinical Psychiatry</i> , 2017 , 78, 1083-1090	4.6	10
326	Telephone-Based Depression Care Management for Postpartum Women: A Randomized Controlled Trial. <i>Journal of Clinical Psychiatry</i> , 2017 , 78, 1369-1375	4.6	3
325	Clinical Profile, Etiology, and Treatment of Chronic Pancreatitis in North American Women: Analysis of a Large Multicenter Cohort. <i>Pancreas</i> , 2016 , 45, 934-40	2.6	19
324	Severe traumatic brain injury in children: a vision for the future. <i>Intensive Care Medicine</i> , 2016 , 42, 1618	-1629	3
323	Factors associated with onset timing, symptoms, and severity of depression identified in the postpartum period. <i>Journal of Affective Disorders</i> , 2016 , 203, 111-120	6.6	32
322	Derivation and External Validation of a Prediction Rule for Five-Year Mortality in Patients With Early Diffuse Cutaneous Systemic Sclerosis. <i>Arthritis and Rheumatology</i> , 2016 , 68, 993-1003	9.5	24
321	Intracranial Hypertension and Cerebral Hypoperfusion in Children With Severe Traumatic Brain Injury: Thresholds and Burden in Accidental and Abusive Insults. <i>Pediatric Critical Care Medicine</i> , 2016 , 17, 444-50	3	28
320	On-pump versus off-pump coronary artery bypass graft surgery among patients with type 2 diabetes in the Bypass Angioplasty Revascularization Investigation 2 Diabetes trial. <i>European Journal of Cardio-thoracic Surgery</i> , 2016 , 49, 406-16	3	13
319	Structure, Process, and Culture Differences of Pediatric Trauma Centers Participating in an International Comparative Effectiveness Study of Children with Severe Traumatic Brain Injury. <i>Neurocritical Care</i> , 2016 , 24, 353-60	3.3	11
318	Is third trimester serotonin reuptake inhibitor use associated with postpartum hemorrhage?. <i>Journal of Psychiatric Research</i> , 2016 , 73, 79-85	5.2	6
317	Efficacy of Cognitive-Behavioral Therapy for Insomnia Combined With Antidepressant Pharmacotherapy in Patients With Comorbid Depression and Insomnia: A Randomized Controlled Trial. <i>Journal of Clinical Psychiatry</i> , 2016 , 77, e1316-e1323	4.6	69
316	Effectiveness of Pharmacological Therapies for Intracranial Hypertension in Children With Severe Traumatic Brain InjuryResults From an Automated Data Collection System Time-Synched to Drug Administration. <i>Pediatric Critical Care Medicine</i> , 2016 , 17, 236-45	3	44
315	Agreement of Physicians and Nurses Performing Tender and Swollen Joint Counts in Rheumatoid Arthritis. <i>Journal of Clinical Rheumatology</i> , 2016 , 22, 30-4	1.1	4

314	Factors Associated with Postpartum Maternal Functioning in Women with Positive Screens for Depression. <i>Journal of Womenis Health</i> , 2016 , 25, 707-13	3	25	
313	Cerebrospinal Fluid Biomarkers of Simian Immunodeficiency Virus Encephalitis: CSF Biomarkers of SIV Encephalitis. <i>Journal of NeuroImmune Pharmacology</i> , 2016 , 11, 332-47	6.9	1	
312	Viral infections in outpatients with medically attended acute respiratory illness during the 2012-2013 influenza season. <i>BMC Infectious Diseases</i> , 2015 , 15, 87	4	17	
311	Intention to receive influenza vaccine after an acute respiratory illness. <i>American Journal of Health Behavior</i> , 2015 , 39, 573-81	1.9	2	
310	Anaphylaxis management before and after implementation of guidelines in the pediatric emergency department. <i>Journal of Allergy and Clinical Immunology: in Practice</i> , 2015 , 3, 604-6.e2	5.4	7	
309	Maternal-Infant interaction in women with unipoloar and bipolar depression. <i>Applied Nursing Research</i> , 2015 , 28, 381-3	1.8	12	
308	Suicidal ideation in depressed postpartum women: Associations with childhood trauma, sleep disturbance and anxiety. <i>Journal of Psychiatric Research</i> , 2015 , 66-67, 95-104	5.2	60	
307	Symptoms of depressed mood, disturbed sleep, and sexual problems in midlife women: cross-sectional data from the Study of Women's Health Across the Nation. <i>Journal of Womenis Health</i> , 2015 , 24, 119-26	3	52	
306	Rationale and Design of the Genomic Research in Alpha-1 Antitrypsin Deficiency and Sarcoidosis (GRADS) Study. Sarcoidosis Protocol. <i>Annals of the American Thoracic Society</i> , 2015 , 12, 1561-71	4.7	42	
305	Surgery versus nonsurgical treatment of lumbar spinal stenosis: a randomized trial. <i>Annals of Internal Medicine</i> , 2015 , 162, 465-73	8	85	
304	Rationale and Design of the Genomic Research in Alpha-1 Antitrypsin Deficiency and Sarcoidosis Study. Alpha-1 Protocol. <i>Annals of the American Thoracic Society</i> , 2015 , 12, 1551-60	4.7	6	
303	Pre-treatment insomnia as a predictor of single and combination antidepressant outcomes: a CO-MED report. <i>Journal of Affective Disorders</i> , 2015 , 174, 157-64	6.6	16	
302	DOES SCREENING WITH THE MDQ AND EPDS IMPROVE IDENTIFICATION OF BIPOLAR DISORDER IN AN OBSTETRICAL SAMPLE?. <i>Depression and Anxiety</i> , 2015 , 32, 518-26	8.4	81	
301	Effect of administration of neuromuscular blocking agents in children with severe traumatic brain injury on acute complication rates and outcomes: a secondary analysis from a randomized, controlled trial of therapeutic hypothermia. <i>Pediatric Critical Care Medicine</i> , 2015 , 16, 352-8	3	12	
300	Chronic pancreatitis pain pattern and severity are independent of abdominal imaging findings. <i>Clinical Gastroenterology and Hepatology</i> , 2015 , 13, 552-60; quiz e28-9	6.9	102	
299	Prevalence and impact of obsessive-compulsive symptoms in depression: a STAR*D report. <i>Journal of Clinical Psychiatry</i> , 2015 , 76, 1668-74	4.6	12	
298	Transdermal Estradiol Treatment for Postpartum Depression: A Pilot, Randomized Trial. <i>Journal of Clinical Psychopharmacology</i> , 2015 , 35, 389-95	1.7	24	
297	Theoretical approaches to maternal-infant interaction: which approach best discriminates between mothers with and without postpartum depression?. <i>Archives of Psychiatric Nursing</i> , 2014 , 28, 384-8	2.1	1	

296	The psychometric properties of the Barkin Index of Maternal Functioning. <i>JOGNN - Journal of Obstetric, Gynecologic, and Neonatal Nursing</i> , 2014 , 43, 792-802	1.2	29
295	Hemorrhagic shock shifts the serum cytokine profile from pro- to anti-inflammatory after experimental traumatic brain injury in mice. <i>Journal of Neurotrauma</i> , 2014 , 31, 1386-95	5.4	36
294	Influenza and other respiratory virus infections in outpatients with medically attended acute respiratory infection during the 2011-12 influenza season. <i>Influenza and Other Respiratory Viruses</i> , 2014 , 8, 397-405	5.6	32
293	Derivation and validation of a prediction rule for two-year mortality in early diffuse cutaneous systemic sclerosis. <i>Arthritis and Rheumatology</i> , 2014 , 66, 1616-24	9.5	29
292	Abnormal screening for gestational diabetes, maternal mood disorder, and preterm birth. <i>Bipolar Disorders</i> , 2014 , 16, 308-17	3.8	10
291	Clinical and functional outcomes of patients who experience partial response to citalopram: secondary analysis of STAR*D. <i>Journal of Psychiatric Practice</i> , 2014 , 20, 178-87	1.3	14
290	Collaborative care outcomes for pediatric behavioral health problems: a cluster randomized trial. <i>Pediatrics</i> , 2014 , 133, e981-92	7.4	146
289	Detection of influenza virus infection using two PCR methods. <i>Advances in Virology</i> , 2014 , 2014, 274679	9 1.9	4
288	Energy expenditure in children after severe traumatic brain injury. <i>Pediatric Critical Care Medicine</i> , 2014 , 15, 242-9	3	26
287	Brief, unidimensional melancholia rating scales are highly sensitive to the effect of citalopram and may have biological validity: implications for the research domain criteria (RDoC). <i>Journal of Affective Disorders</i> , 2014 , 163, 18-24	6.6	30
286	Impact of prenatal exposure to serotonin reuptake inhibitors or maternal major depressive disorder on infant developmental outcomes. <i>Journal of Clinical Psychiatry</i> , 2014 , 75, 1088-95	4.6	35
285	CBT for treatment resistant depression. <i>Lancet, The</i> , 2013 , 381, 352-3	40	5
284	Brain imaging predictors and the international study to predict optimized treatment for depression: study protocol for a randomized controlled trial. <i>Trials</i> , 2013 , 14, 224	2.8	31
283	Hypothermia in paediatric traumatic brain injuryauthors' reply. Lancet Neurology, The, 2013, 12, 849-8.	5 0 4.1	1
282	Prospective evaluation of nighttime hot flashes during pregnancy and postpartum. <i>Fertility and Sterility</i> , 2013 , 100, 1667-72	4.8	6
281	Effect of hyperoxia on resuscitation of experimental combined traumatic brain injury and hemorrhagic shock in mice. <i>Anesthesiology</i> , 2013 , 118, 649-63	4.3	21
280	The clinical relevance of self-reported premenstrual worsening of depressive symptoms in the management of depressed outpatients: a STAR*D report. <i>Journal of Womenis Health</i> , 2013 , 22, 219-29	3	20
279	Comparison of hypothermia and normothermia after severe traumatic brain injury in children (Cool Kids): a phase 3, randomised controlled trial. <i>Lancet Neurology, The</i> , 2013 , 12, 546-53	24.1	178

(2012-2013)

278	Findings from the Sequenced Treatment Alternatives to Relieve Depression (STAR*D) study. Journal of Womenis Health, 2013, 22, 121-31	3	30
277	Response to Grzeskowiak and Morrison. <i>American Journal of Psychiatry</i> , 2013 , 170, 1364-5	11.9	
276	MRI assessment of cerebral blood flow after experimental traumatic brain injury combined with hemorrhagic shock in mice. <i>Journal of Cerebral Blood Flow and Metabolism</i> , 2013 , 33, 129-36	7.3	31
275	Relationship between obesity and depression: characteristics and treatment outcomes with antidepressant medication. <i>Psychosomatic Medicine</i> , 2013 , 75, 863-72	3.7	34
274	Disturbed sleep and inflammatory cytokines in depressed and nondepressed pregnant women: an exploratory analysis of pregnancy outcomes. <i>Psychosomatic Medicine</i> , 2013 , 75, 670-81	3.7	63
273	Differences in medical therapy goals for children with severe traumatic brain injury-an international study. <i>Pediatric Critical Care Medicine</i> , 2013 , 14, 811-8	3	53
272	Onset timing, thoughts of self-harm, and diagnoses in postpartum women with screen-positive depression findings. <i>JAMA Psychiatry</i> , 2013 , 70, 490-8	14.5	516
271	Increase in work productivity of depressed individuals with improvement in depressive symptom severity. <i>American Journal of Psychiatry</i> , 2013 , 170, 633-41	11.9	84
270	Clinical and sociodemographic characteristics associated with suicidal ideation in depressed outpatients. <i>Canadian Journal of Psychiatry</i> , 2013 , 58, 113-22	4.8	25
269	The impact of HAART on the respiratory complications of HIV infection: longitudinal trends in the MACS and WIHS cohorts. <i>PLoS ONE</i> , 2013 , 8, e58812	3.7	32
268	Combining Medications to Enhance Depression Outcomes (CO-MED): Acute and Long-Term Outcomes of a Single-Blind Randomized Study. <i>Focus (American Psychiatric Publishing)</i> , 2012 , 10, 492-50	5 ^{1.1}	1
267	What Did STAR?D Teach Us? Results From a Large-Scale, Practical, Clinical Trial for Patients With Depression. <i>Focus (American Psychiatric Publishing)</i> , 2012 , 10, 510-517	1.1	
266	Common genetic variants in the CLDN2 and PRSS1-PRSS2 loci alter risk for alcohol-related and sporadic pancreatitis. <i>Nature Genetics</i> , 2012 , 44, 1349-54	36.3	223
265	Baseline depression severity as a predictor of single and combination antidepressant treatment outcome: results from the CO-MED trial. <i>European Neuropsychopharmacology</i> , 2012 , 22, 183-99	1.2	48
264	Risk factors for mortality in children with abusive head trauma. <i>Journal of Pediatrics</i> , 2012 , 161, 716-722	.e. 6	49
263	How influenza vaccination policy may affect vaccine logistics. <i>Vaccine</i> , 2012 , 30, 4517-23	4.1	18
262	Validity of a pediatric version of the Glasgow Outcome Scale-Extended. <i>Journal of Neurotrauma</i> , 2012 , 29, 1126-39	5.4	162
261	Mortality among foreign nationals in Chiang Mai City, Thailand, 2010 to 2011. <i>Journal of Travel Medicine</i> , 2012 , 19, 334-51	12.9	4

260	Cognitive therapy for anxious depression in STAR(*) D: what have we learned?. <i>Journal of Affective Disorders</i> , 2012 , 142, 213-8	6.6	37
259	Local spatial and temporal processes of influenza in Pennsylvania, USA: 2003-2009. <i>PLoS ONE</i> , 2012 , 7, e34245	3.7	16
258	Sensitivity, specificity, and predictive power of the "Brief Risk-resilience Index for SCreening," a brief pan-diagnostic web screen for emotional health. <i>Brain and Behavior</i> , 2012 , 2, 576-89	3.4	13
257	What are the clinical implications of new onset or worsening anxiety during the first two weeks of SSRI treatment for depression?. <i>Depression and Anxiety</i> , 2012 , 29, 94-101	8.4	20
256	Effect of concurrent substance use disorder on the effectiveness of single and combination antidepressant medications for the treatment of major depression: an exploratory analysis of a single-blind randomized trial. <i>Depression and Anxiety</i> , 2012 , 29, 111-22	8.4	13
255	Effects of heart disease on depression treatment: results from the COMED study. <i>General Hospital Psychiatry</i> , 2012 , 34, 24-34	5.6	4
254	Depression treatment in patients with general medical conditions: results from the CO-MED trial. <i>Annals of Family Medicine</i> , 2012 , 10, 23-33	2.9	28
253	Implementation of AF-CBT by community practitioners serving child welfare and mental health: a randomized trial. <i>Child Maltreatment</i> , 2012 , 17, 32-46	2.8	46
252	Brain tissue oxygen monitoring after severe traumatic brain injury in children: relationship to outcome and association with other clinical parameters. <i>Journal of Neurosurgery: Pediatrics</i> , 2012 , 10, 383-91	2.1	36
	Response to Blier Letter. <i>American Journal of Psychiatry</i> , 2012 , 169, 95-96		2
251	Response to biler Letter. American Journal of Psychiatry, 2012, 103, 33-30	11.9	2
251 250	Severe brief pressure-controlled hemorrhagic shock after traumatic brain injury exacerbates functional deficits and long-term neuropathological damage in mice. <i>Journal of Neurotrauma</i> , 2012 , 29, 2192-208	5.4	43
	Severe brief pressure-controlled hemorrhagic shock after traumatic brain injury exacerbates functional deficits and long-term neuropathological damage in mice. <i>Journal of Neurotrauma</i> , 2012 ,		
250	Severe brief pressure-controlled hemorrhagic shock after traumatic brain injury exacerbates functional deficits and long-term neuropathological damage in mice. <i>Journal of Neurotrauma</i> , 2012 , 29, 2192-208	5.4	43
250 249	Severe brief pressure-controlled hemorrhagic shock after traumatic brain injury exacerbates functional deficits and long-term neuropathological damage in mice. <i>Journal of Neurotrauma</i> , 2012 , 29, 2192-208 Postpartum lipid levels in women with major depression. <i>Journal of Womenis Health</i> , 2012 , 21, 534-8	5.4	43
250249248	Severe brief pressure-controlled hemorrhagic shock after traumatic brain injury exacerbates functional deficits and long-term neuropathological damage in mice. <i>Journal of Neurotrauma</i> , 2012 , 29, 2192-208 Postpartum lipid levels in women with major depression. <i>Journal of Womenis Health</i> , 2012 , 21, 534-8 Disturbed sleep, a novel risk factor for preterm birth?. <i>Journal of Womenis Health</i> , 2012 , 21, 54-60 Correlates and outcomes of depressed out-patients with greater and fewer anxious symptoms: a	5·4 3 3	43 11 44
250249248247	Severe brief pressure-controlled hemorrhagic shock after traumatic brain injury exacerbates functional deficits and long-term neuropathological damage in mice. <i>Journal of Neurotrauma</i> , 2012 , 29, 2192-208 Postpartum lipid levels in women with major depression. <i>Journal of Womenis Health</i> , 2012 , 21, 534-8 Disturbed sleep, a novel risk factor for preterm birth?. <i>Journal of Womenis Health</i> , 2012 , 21, 54-60 Correlates and outcomes of depressed out-patients with greater and fewer anxious symptoms: a CO-MED report. <i>International Journal of Neuropsychopharmacology</i> , 2012 , 15, 1387-99 Relationship between hyperglycemia and outcome in children with severe traumatic brain injury.	5·4 3 3 5.8	43 11 44 18
250249248247246	Severe brief pressure-controlled hemorrhagic shock after traumatic brain injury exacerbates functional deficits and long-term neuropathological damage in mice. <i>Journal of Neurotrauma</i> , 2012 , 29, 2192-208 Postpartum lipid levels in women with major depression. <i>Journal of Womenis Health</i> , 2012 , 21, 534-8 Disturbed sleep, a novel risk factor for preterm birth?. <i>Journal of Womenis Health</i> , 2012 , 21, 54-60 Correlates and outcomes of depressed out-patients with greater and fewer anxious symptoms: a CO-MED report. <i>International Journal of Neuropsychopharmacology</i> , 2012 , 15, 1387-99 Relationship between hyperglycemia and outcome in children with severe traumatic brain injury. <i>Pediatric Critical Care Medicine</i> , 2012 , 13, 85-91	5·4 3 5.8 3	43 11 44 18 55

242	Short-term repeatability of diurnal intraocular pressure patterns in glaucomatous individuals. <i>Ophthalmology</i> , 2011 , 118, 47-51	7.3	64
241	Residual symptoms in depressed outpatients who respond by 50% but do not remit to antidepressant medication. <i>Journal of Clinical Psychopharmacology</i> , 2011 , 31, 180-6	1.7	123
240	The relationship between adverse events during selective serotonin reuptake inhibitor treatment for major depressive disorder and nonremission in the suicide assessment methodology study. Journal of Clinical Psychopharmacology, 2011, 31, 31-8	1.7	13
239	Changes in depressive symptoms and social functioning in the sequenced treatment alternatives to relieve depression study. <i>Journal of Nervous and Mental Disease</i> , 2011 , 199, 807-10	1.8	15
238	Reduction in the incidence of influenza A but not influenza B associated with use of hand sanitizer and cough hygiene in schools: a randomized controlled trial. <i>Pediatric Infectious Disease Journal</i> , 2011 , 30, 921-6	3.4	61
237	Intracranial pressure-monitoring systems in children with traumatic brain injury: combining therapeutic and diagnostic tools. <i>Pediatric Critical Care Medicine</i> , 2011 , 12, 560-5	3	38
236	International Study to Predict Optimized Treatment for Depression (iSPOT-D), a randomized clinical trial: rationale and protocol. <i>Trials</i> , 2011 , 12, 4	2.8	126
235	Changes in sleep quality, but not hormones predict time to postpartum depression recurrence. Journal of Affective Disorders, 2011 , 130, 378-84	6.6	104
234	Randomized comparison of selective serotonin reuptake inhibitor (escitalopram) monotherapy and antidepressant combination pharmacotherapy for major depressive disorder with melancholic features: a CO-MED report. <i>Journal of Affective Disorders</i> , 2011 , 133, 467-76	6.6	42
233	Suicide and suicide attempts in the Systematic Treatment Enhancement Program for Bipolar Disorder (STEP-BD). <i>Journal of Affective Disorders</i> , 2011 , 133, 423-7	6.6	46
232	Impact of changing the measles vaccine vial size on Niger's vaccine supply chain: a computational model. <i>BMC Public Health</i> , 2011 , 11, 425	4.1	50
231	Assessing anxious features in depressed outpatients. <i>International Journal of Methods in Psychiatric Research</i> , 2011 , 20, e69-82	4.3	15
230	Sleep disturbances in depressed and nondepressed pregnant women. <i>Depression and Anxiety</i> , 2011 , 28, 676-85	8.4	77
229	Depression treatment and maternal functioning. <i>Depression and Anxiety</i> , 2011 , 28, 1020-6	8.4	28
228	Children of depressed mothers 1 year after remission of maternal depression: findings from the STAR*D-Child study. <i>American Journal of Psychiatry</i> , 2011 , 168, 593-602	11.9	124
227	Combining medications to enhance depression outcomes (CO-MED): acute and long-term outcomes of a single-blind randomized study. <i>American Journal of Psychiatry</i> , 2011 , 168, 689-701	11.9	266
226	Agility and perturbation training techniques in exercise therapy for reducing pain and improving function in people with knee osteoarthritis: a randomized clinical trial. <i>Physical Therapy</i> , 2011 , 91, 452-6	3 ·3	89
225	Effects of race and ethnicity on depression treatment outcomes: the CO-MED trial. <i>Psychiatric Services</i> , 2011 , 62, 1167-79	3.3	25

224	Mother-infant antidepressant concentrations, maternal depression, and perinatal events. <i>Journal of Clinical Psychiatry</i> , 2011 , 72, 994-1001	4.6	43
223	Effect of antidepressant medication treatment on suicidal ideation and behavior in a randomized trial: an exploratory report from the Combining Medications to Enhance Depression Outcomes Study. <i>Journal of Clinical Psychiatry</i> , 2011 , 72, 1322-32	4.6	34
222	Concise Health Risk Tracking scale: a brief self-report and clinician rating of suicidal risk. <i>Journal of Clinical Psychiatry</i> , 2011 , 72, 757-64	4.6	81
221	Concise Associated Symptoms Tracking scale: a brief self-report and clinician rating of symptoms associated with suicidality. <i>Journal of Clinical Psychiatry</i> , 2011 , 72, 765-74	4.6	42
220	Endothelin-1 is increased in cerebrospinal fluid and associated with unfavorable outcomes in children after severe traumatic brain injury. <i>Journal of Neurotrauma</i> , 2010 , 27, 1819-25	5.4	52
219	Assessment of functioning in new mothers. <i>Journal of Womenis Health</i> , 2010 , 19, 1493-9	3	43
218	Development of the Barkin Index of Maternal Functioning. Journal of Womenis Health, 2010, 19, 2239-	463	59
217	Esynuclein levels are elevated in cerebrospinal fluid following traumatic brain injury in infants and children: the effect of therapeutic hypothermia. <i>Developmental Neuroscience</i> , 2010 , 32, 385-95	2.2	37
216	Relationship of intracranial pressure and cerebral perfusion pressure with outcome in young children after severe traumatic brain injury. <i>Developmental Neuroscience</i> , 2010 , 32, 413-9	2.2	41
215	Mechanical vs manual manipulation for low back pain: an observational cohort study. <i>Journal of Manipulative and Physiological Therapeutics</i> , 2010 , 33, 193-200	1.3	12
214	Does comorbid substance use disorder impair recovery from major depression with SSRI treatment? An analysis of the STAR*D level one treatment outcomes. <i>Drug and Alcohol Dependence</i> , 2010 , 107, 161-70	4.9	59
213	Diurnal intraocular pressure patterns are not repeatable in the short term in healthy individuals. <i>Ophthalmology</i> , 2010 , 117, 1700-4	7.3	56
212	Do atypical features affect outcome in depressed outpatients treated with citalopram?. <i>International Journal of Neuropsychopharmacology</i> , 2010 , 13, 15-30	5.8	41
211	Disposition of chiral and racemic fluoxetine and norfluoxetine across childbearing. <i>Journal of Clinical Psychopharmacology</i> , 2010 , 30, 381-6	1.7	20
210	Early adverse events and attrition in selective serotonin reuptake inhibitor treatment: a suicide assessment methodology study report. <i>Journal of Clinical Psychopharmacology</i> , 2010 , 30, 259-66	1.7	17
209	The influence of menopause status and postmenopausal use of hormone therapy on presentation of major depression in women. <i>Menopause</i> , 2010 , 17, 828-39	2.5	43
208	Insomnia in patients with depression: a STAR*D report. CNS Spectrums, 2010, 15, 394-404	1.8	102
207	Dysfunctional Attitudes, Attributional Styles, and Phase of Illness in Bipolar Disorder. <i>Cognitive Therapy and Research</i> , 2010 , 34, 24-34	2.7	14

206	The impact of diabetes on depression treatment outcomes. General Hospital Psychiatry, 2010, 32, 33-41	5.6	20
205	What did STAR*D teach us? Results from a large-scale, practical, clinical trial for patients with depression. <i>Psychiatric Services</i> , 2009 , 60, 1439-45	3.3	368
204	Predictors of pain and function outcome after rehabilitation in patients with patellofemoral pain syndrome. <i>Journal of Rehabilitation Medicine</i> , 2009 , 41, 604-12	3.4	93
203	Effectiveness Study of Venlafaxine-XR Combined with Aripiprazole for Chronic or Recurrent Major Depressive Disorder. <i>Australian and New Zealand Journal of Psychiatry</i> , 2009 , 43, 956-967	2.6	8
202	Identifying risk for attrition during treatment for depression. <i>Psychotherapy and Psychosomatics</i> , 2009 , 78, 372-9	9.4	20
201	Cognitive Therapy Augmentation versus CT Switch Treatment: A STAR*D Report. <i>International Journal of Cognitive Therapy</i> , 2009 , 2, 66-87	1.4	1
200	Can phase III trial results of antidepressant medications be generalized to clinical practice? A STAR*D report. <i>American Journal of Psychiatry</i> , 2009 , 166, 599-607	11.9	171
199	What predicts attrition in second step medication treatments for depression?: a STAR*D Report. <i>International Journal of Neuropsychopharmacology</i> , 2009 , 12, 459-73	5.8	58
198	Sex differences in response to citalopram: a STAR*D report. <i>Journal of Psychiatric Research</i> , 2009 , 43, 503-11	5.2	123
197	The functional impact of subsyndromal depressive symptoms in bipolar disorder: data from STEP-BD. <i>Journal of Affective Disorders</i> , 2009 , 114, 58-67	6.6	79
196	Clinical correlates of the worsening or emergence of suicidal ideation during SSRI treatment of depression: an examination of citalopram in the STAR*D study. <i>Journal of Affective Disorders</i> , 2009 , 117, 63-73	6.6	73
195	Income and attrition in the treatment of depression: a STAR*D report. <i>Depression and Anxiety</i> , 2009 , 26, 622-33	8.4	39
194	Diurnal mood variation in outpatients with major depressive disorder. <i>Depression and Anxiety</i> , 2009 , 26, 851-63	8.4	18
193	Sociodemographic, clinical, and treatment characteristics associated with worsened depression during treatment with citalopram: results of the NIMH STAR(*)D trial. <i>Depression and Anxiety</i> , 2009 , 26, 612-21	8.4	25
192	Early adversity in chronic depression: clinical correlates and response to pharmacotherapy. <i>Depression and Anxiety</i> , 2009 , 26, 701-10	8.4	113
191	Release bias in accessing medical records in clinical trials: a STAR*D report. <i>International Journal of Methods in Psychiatric Research</i> , 2009 , 18, 147-58	4.3	1
190	Running in place: implications of HIV incidence estimates among urban men who have sex with men in the United States and other industrialized countries. <i>AIDS and Behavior</i> , 2009 , 13, 615-29	4.3	92
189	Relationship between increases in pancreatic enzymes and cerebral events in children after traumatic brain injury. <i>Neurocritical Care</i> , 2009 , 11, 322-9	3.3	5

188	Global benefit-risk analysis of adjunctive aripiprazole in the treatment of patients with major depressive disorder. <i>Pharmacoepidemiology and Drug Safety</i> , 2009 , 18, 965-72	2.6	5
187	Retrospective age at onset of bipolar disorder and outcome during two-year follow-up: results from the STEP-BD study. <i>Bipolar Disorders</i> , 2009 , 11, 391-400	3.8	187
186	Concurrent anxiety and substance use disorders among outpatients with major depression: clinical features and effect on treatment outcome. <i>Drug and Alcohol Dependence</i> , 2009 , 99, 248-60	4.9	109
185	Associates of physical function and pain in patients with patellofemoral pain syndrome. <i>Archives of Physical Medicine and Rehabilitation</i> , 2009 , 90, 285-95	2.8	84
184	Relationships of fear, anxiety, and depression with physical function in patients with knee osteoarthritis. <i>Archives of Physical Medicine and Rehabilitation</i> , 2009 , 90, 1866-73	2.8	115
183	STAR*D: revising conventional wisdom. <i>CNS Drugs</i> , 2009 , 23, 627-47	6.7	172
182	The impact of nonclinical factors on care use for patients with depression: a STAR*D report. <i>CNS Neuroscience and Therapeutics</i> , 2009 , 15, 320-32	6.8	9
181	Metabolic assessment of aripiprazole as adjunctive therapy in major depressive disorder: a pooled analysis of 2 studies. <i>Journal of Clinical Psychopharmacology</i> , 2009 , 29, 362-7	1.7	14
180	Therapeutic hypothermia preserves antioxidant defenses after severe traumatic brain injury in infants and children. <i>Critical Care Medicine</i> , 2009 , 37, 689-95	1.4	122
179	Factors associated with concomitant psychotropic drug use in the treatment of major depression: a STAR*D Report. <i>CNS Spectrums</i> , 2009 , 14, 487-98	1.8	8
178	Do depressed patients with diabetes experience more side effects when treated with CitalopramThan their counterparts without diabetes? a STAR*D study. <i>Primary Care Companion To the Journal of Clinical Psychiatry</i> , 2009 , 11, 186-96		
177	Family history of depression and therapeutic outcome: findings from STAR*D. <i>Journal of Clinical Psychiatry</i> , 2009 , 70, 185-95	4.6	15
176	Voice response system to measure healthcare costs: a STAR*D report. <i>American Journal of Managed Care</i> , 2009 , 15, 153-62	2.1	3
175	Increases in cerebrospinal fluid caffeine concentration are associated with favorable outcome after severe traumatic brain injury in humans. <i>Journal of Cerebral Blood Flow and Metabolism</i> , 2008 , 28, 395-	40 ⁷ 1 ³	34
174	A comparison of baseline sociodemographic and clinical characteristics between major depressive disorder patients with and without diabetes: a STAR*D report. <i>Journal of Affective Disorders</i> , 2008 , 108, 113-20	6.6	13
173	Family history of completed suicide and characteristics of major depressive disorder: a STAR*D (sequenced treatment alternatives to relieve depression) study. <i>Journal of Affective Disorders</i> , 2008 , 108, 129-34	6.6	12
172	Antidepressant-associated chronic irritable dysphoria (ACID) in STEP-BD patients. <i>Journal of Affective Disorders</i> , 2008 , 111, 372-7	6.6	56
171	The FKBP5-gene in depression and treatment responsean association study in the Sequenced Treatment Alternatives to Relieve Depression (STAR*D) Cohort. <i>Biological Psychiatry</i> , 2008 , 63, 1103-1	o ^{7.9}	218

(2008-2008)

170	Sex differences in depression symptoms in treatment-seeking adults: confirmatory analyses from the Sequenced Treatment Alternatives to Relieve Depression study. <i>Comprehensive Psychiatry</i> , 2008 , 49, 238-46	7.3	156
169	Treatment outcomes for older depressed patients with earlier versus late onset of first depressive episode: a Sequenced Treatment Alternatives to Relieve Depression (STAR*D) report. <i>American Journal of Geriatric Psychiatry</i> , 2008 , 16, 58-64	6.5	32
168	Remission of maternal depression: relations to family functioning and youth internalizing and externalizing symptoms. <i>Journal of Clinical Child and Adolescent Psychology</i> , 2008 , 37, 714-24	5.4	47
167	Antecedentes familiares de suicidio consumado y caracter\(\textstyle{\textstyle{1}}\) ticas del trastorno depresivo mayor: estudio STAR*D (sequenced treatment alternatives to relieve depression). <i>Psiquiatria Biologica</i> , 2008 , 15, 195-201	0.2	
166	Selecting among second-step antidepressant medication monotherapies: predictive value of clinical, demographic, or first-step treatment features. <i>Archives of General Psychiatry</i> , 2008 , 65, 870-80		125
165	The prospective course of rapid-cycling bipolar disorder: findings from the STEP-BD. <i>American Journal of Psychiatry</i> , 2008 , 165, 370-7; quiz 410	11.9	129
164	Difference in treatment outcome in outpatients with anxious versus nonanxious depression: a STAR*D report. <i>American Journal of Psychiatry</i> , 2008 , 165, 342-51	11.9	594
163	Acute and Longer-Term Outcomes in Depressed Outpatients Requiring One or Several Treatment Steps: A STAR*D Report. <i>Focus (American Psychiatric Publishing)</i> , 2008 , 6, 128-142	1.1	9
162	Children of depressed mothers 1 year after the initiation of maternal treatment: findings from the STAR*D-Child Study. <i>American Journal of Psychiatry</i> , 2008 , 165, 1136-47	11.9	146
161	Cognitive Therapy Versus Medication in Augmentation and Switch Strategies as Second-Step Treatments: A STAR*D Report. <i>Focus (American Psychiatric Publishing)</i> , 2008 , 6, 104-119	1.1	
160	Depression outcomes of Spanish- and english-speaking Hispanic outpatients in STAR*D. <i>Psychiatric Services</i> , 2008 , 59, 1273-84	3.3	26
159	Caregiver burden and health in bipolar disorder: a cluster analytic approach. <i>Journal of Nervous and Mental Disease</i> , 2008 , 196, 484-91	1.8	50
158	An open pilot study of the combination of escitalopram and bupropion-SR for outpatients with major depressive disorder. <i>Journal of Psychiatric Practice</i> , 2008 , 14, 271-80	1.3	34
157	Course and Severity of Maternal Depression: Associations with Family Functioning and Child Adjustment. <i>Journal of Youth and Adolescence</i> , 2008 , 37, 906-916	4.5	68
156	Primary versus specialty care outcomes for depressed outpatients managed with measurement-based care: results from STAR*D. <i>Journal of General Internal Medicine</i> , 2008 , 23, 551-60	4	74
155	Clinically Relevant Findings from STAR*D. <i>Psychiatric Annals</i> , 2008 , 38,	0.5	11
154	Does the duration of index episode affect the treatment outcome of major depressive disorder? A STAR*D report. <i>Journal of Clinical Psychiatry</i> , 2008 , 69, 1246-56	4.6	31
153	Response to a selective serotonin reuptake inhibitor (citalopram) in major depressive disorder with melancholic features: a STAR*D report. <i>Journal of Clinical Psychiatry</i> , 2008 , 69, 1847-55	4.6	80

152	Effectiveness of adjunctive antidepressant treatment for bipolar depression. <i>New England Journal of Medicine</i> , 2007 , 356, 1711-22	59.2	671
151	Is Psychosocial Management Effective?Reply. Archives of General Psychiatry, 2007, 64, 1452		
150	Clinical differences among depressed patients with and without a history of suicide attempts: findings from the STAR*D trial. <i>Journal of Affective Disorders</i> , 2007 , 97, 77-84	6.6	59
149	Influences of hormone-based contraception on depressive symptoms in premenopausal women with major depression. <i>Psychoneuroendocrinology</i> , 2007 , 32, 843-53	5	61
148	Family history of mood disorder and characteristics of major depressive disorder: a STAR*D (sequenced treatment alternatives to relieve depression) study. <i>Journal of Psychiatric Research</i> , 2007 , 41, 214-21	5.2	49
147	Electronic source materials in clinical research: acceptability and validity of symptom self-rating in major depressive disorder. <i>Journal of Psychiatric Research</i> , 2007 , 41, 737-43	5.2	18
146	The STAR*D Project results: a comprehensive review of findings. <i>Current Psychiatry Reports</i> , 2007 , 9, 449-59	9.1	443
145	Adjunctive antidepressant use and symptomatic recovery among bipolar depressed patients with concomitant manic symptoms: findings from the STEP-BD. <i>American Journal of Psychiatry</i> , 2007 , 164, 1348-55	11.9	161
144	Maximizing the adequacy of medication treatment in controlled trials and clinical practice: STAR(*)D measurement-based care. <i>Neuropsychopharmacology</i> , 2007 , 32, 2479-89	8.7	106
143	Cognitive therapy versus medication in augmentation and switch strategies as second-step treatments: a STAR*D report. <i>American Journal of Psychiatry</i> , 2007 , 164, 739-52	11.9	246
142	Insured and non-insured depressed outpatients: how do they compare?. <i>Annals of Clinical Psychiatry</i> , 2007 , 19, 73-82	1.4	4
141	Distal support and community living among individuals diagnosed with schizophrenia and schizoaffective disorder. <i>Psychiatry (New York)</i> , 2007 , 70, 1-11	1	13
140	Acceptability of second-step treatments to depressed outpatients: a STAR*D report. <i>American Journal of Psychiatry</i> , 2007 , 164, 753-60	11.9	56
139	Association of GRIK4 with outcome of antidepressant treatment in the STAR*D cohort. <i>American Journal of Psychiatry</i> , 2007 , 164, 1181-8	11.9	169
138	Predictors of attrition during initial (citalopram) treatment for depression: a STAR*D report. <i>American Journal of Psychiatry</i> , 2007 , 164, 1189-97	11.9	106
137	Intensive psychosocial intervention enhances functioning in patients with bipolar depression: results from a 9-month randomized controlled trial. <i>American Journal of Psychiatry</i> , 2007 , 164, 1340-7	11.9	186
136	Effect of age at onset on the course of major depressive disorder. <i>American Journal of Psychiatry</i> , 2007 , 164, 1539-46	11.9	295
135	Serum and cerebrospinal fluid magnesium in severe traumatic brain injury outcome. <i>Journal of Neurotrauma</i> , 2007 , 24, 1347-54	5.4	20

134	Association between a functional serotonin transporter promoter polymorphism and citalopram treatment in adult outpatients with major depression. <i>Archives of General Psychiatry</i> , 2007 , 64, 783-92		182	
133	Major depression symptoms in primary care and psychiatric care settings: a cross-sectional analysis. <i>Annals of Family Medicine</i> , 2007 , 5, 126-34	2.9	79	
132	Ethnicity/race and outcome in the treatment of depression: results from STAR*D. <i>Medical Care</i> , 2007 , 45, 1043-51	3.1	43	
131	Comparison of quality of life measures in a depressed population. <i>Journal of Nervous and Mental Disease</i> , 2007 , 195, 219-25	1.8	27	
130	Psychosocial treatments for bipolar depression: a 1-year randomized trial from the Systematic Treatment Enhancement Program. <i>Archives of General Psychiatry</i> , 2007 , 64, 419-26		379	
129	Pain in depression: STAR*D study findings. <i>Journal of Psychosomatic Research</i> , 2007 , 63, 113-22	4.1	64	
128	Major depressive disorder and comorbid cardiac disease: is there a depressive subtype with greater cardiovascular morbidity? Results from the STAR*D study. <i>Psychosomatics</i> , 2007 , 48, 418-25	2.6	24	
127	Multiplex assessment of cytokine and chemokine levels in cerebrospinal fluid following severe pediatric traumatic brain injury: effects of moderate hypothermia. <i>Journal of Neurotrauma</i> , 2007 , 24, 1707-17	5.4	118	
126	Diurnal Mood Variation in Outpatients With Major Depressive Disorder. <i>Journal of Clinical Psychiatry</i> , 2007 , 68, 1339-1347	4.6	29	
125	Are depressed outpatients with and without a family history of substance use disorder different? A baseline analysis of the STAR*D cohort. <i>Journal of Clinical Psychiatry</i> , 2007 , 68, 1931-8	4.6	10	
124	Clinical features of depression in outpatients with and without co-occurring general medical conditions in STAR*D: confirmatory analysis. <i>Primary Care Companion To the Journal of Clinical Psychiatry</i> , 2007 , 9, 7-15		44	
123	Diurnal mood variation in outpatients with major depressive disorder: implications for DSM-V from an analysis of the Sequenced Treatment Alternatives to Relieve Depression Study data. <i>Journal of Clinical Psychiatry</i> , 2007 , 68, 1339-47	4.6	12	
122	Presenting characteristics of depressed outpatients as a function of recurrence: preliminary findings from the STAR*D clinical trial. <i>Journal of Psychiatric Research</i> , 2006 , 40, 59-69	5.2	78	
121	Predictors of Recurrence in Bipolar Disorder: Primary Outcomes From the Systematic Treatment Enhancement Program for Bipolar Disorder (STEP-BD). <i>Focus (American Psychiatric Publishing)</i> , 2006 , 4, 553-561	1.1	6	
120	Mitochondrial heat shock protein 60 is increased in cerebrospinal fluid following pediatric traumatic brain injury. <i>Developmental Neuroscience</i> , 2006 , 28, 336-41	2.2	23	
119	Predictors of recurrence in bipolar disorder: primary outcomes from the Systematic Treatment Enhancement Program for Bipolar Disorder (STEP-BD). <i>American Journal of Psychiatry</i> , 2006 , 163, 217-2	24 ^{11.9}	473	
118	Remissions in maternal depression and child psychopathology: a STAR*D-child report. <i>JAMA - Journal of the American Medical Association</i> , 2006 , 295, 1389-98	27.4	442	
117	A Comparison of Mirtazapine and Nortriptyline Following Two Consecutive Failed Medication Treatments for Depressed Outpatients: A STAR*D Report. <i>American Journal of Psychiatry</i> , 2006 , 163, 1161-1172	11.9	203	

116	Pharmacological Treatment Patterns at Study Entry for the First 500 STEP-BD Participants. <i>Psychiatric Services</i> , 2006 , 57, 660-5	3.3	101
115	Clinical and demographic factors associated with DSM-IV melancholic depression. <i>Annals of Clinical Psychiatry</i> , 2006 , 18, 91-8	1.4	86
114	A comparison of lithium and T(3) augmentation following two failed medication treatments for depression: a STAR*D report. <i>American Journal of Psychiatry</i> , 2006 , 163, 1519-30; quiz 1665	11.9	380
113	Treatment-resistant bipolar depression: a STEP-BD equipoise randomized effectiveness trial of antidepressant augmentation with lamotrigine, inositol, or risperidone. <i>American Journal of Psychiatry</i> , 2006 , 163, 210-6	11.9	190
112	Evaluation of outcomes with citalopram for depression using measurement-based care in STAR*D: implications for clinical practice. <i>American Journal of Psychiatry</i> , 2006 , 163, 28-40	11.9	2607
111	Medication augmentation after the failure of SSRIs for depression. <i>New England Journal of Medicine</i> , 2006 , 354, 1243-52	59.2	748
110	Acute and longer-term outcomes in depressed outpatients requiring one or several treatment steps: a STAR*D report. <i>American Journal of Psychiatry</i> , 2006 , 163, 1905-17	11.9	3098
109	TREAD: TReatment with Exercise Augmentation for Depression: study rationale and design. <i>Clinical Trials</i> , 2006 , 3, 291-305	2.2	60
108	Bupropion-SR, sertraline, or venlafaxine-XR after failure of SSRIs for depression. <i>New England Journal of Medicine</i> , 2006 , 354, 1231-42	59.2	778
107	Variation in the gene encoding the serotonin 2A receptor is associated with outcome of antidepressant treatment. <i>American Journal of Human Genetics</i> , 2006 , 78, 804-814	11	396
106	An evaluation of the quick inventory of depressive symptomatology and the hamilton rating scale for depression: a sequenced treatment alternatives to relieve depression trial report. <i>Biological Psychiatry</i> , 2006 , 59, 493-501	7.9	300
105	Prevention of missing data in clinical research studies. <i>Biological Psychiatry</i> , 2006 , 59, 997-1000	7.9	27
104	Comparisons of perceived quality of life across clinical states in bipolar disorder: data from the first 2000 Systematic Treatment Enhancement Program for Bipolar Disorder (STEP-BD) participants. <i>Comprehensive Psychiatry</i> , 2006 , 47, 161-8	7.3	47
103	Enrolling research subjects from clinical practice: ethical and procedural issues in the Sequenced Treatment Alternatives to Relieve Depression (STAR*D) trial. <i>Psychiatry Research</i> , 2006 , 141, 193-200	9.9	15
102	Omega-3 fatty acids in bipolar disorder: clinical and research considerations. <i>Prostaglandins Leukotrienes and Essential Fatty Acids</i> , 2006 , 75, 315-21	2.8	20
101	Self-rated global measure of the frequency, intensity, and burden of side effects. <i>Journal of Psychiatric Practice</i> , 2006 , 12, 71-9	1.3	176
100	Tranylcypromine versus venlafaxine plus mirtazapine following three failed antidepressant medication trials for depression: a STAR*D report. <i>American Journal of Psychiatry</i> , 2006 , 163, 1531-41; quiz 1666	11.9	244
99	Suicidal Ideation and Pharmacotherapy Among STEP-BD Patients. <i>Focus (American Psychiatric Publishing)</i> , 2006 , 4, 565-572	1.1	

(2005-2006)

98	What clinical and symptom features and comorbid disorders characterize outpatients with anxious major depressive disorder: a replication and extension. <i>Canadian Journal of Psychiatry</i> , 2006 , 51, 823-35	4.8	115
97	Dr. Trivedi and Colleagues Reply. American Journal of Psychiatry, 2006, 163, 1293-1294	11.9	4
96	Psychotherapy, symptom outcomes, and role functioning over one year among patients with bipolar disorder. <i>Psychiatric Services</i> , 2006 , 57, 959-65	3.3	15
95	Prospective 12-month course of bipolar disorder in out-patients with and without comorbid anxiety disorders. <i>British Journal of Psychiatry</i> , 2006 , 189, 20-5	5.4	164
94	Prospective predictors of suicide and suicide attempts in 1,556 patients with bipolar disorders followed for up to 2 years. <i>Bipolar Disorders</i> , 2006 , 8, 566-75	3.8	91
93	Substance use disorder comorbidity in major depressive disorder: a confirmatory analysis of the STAR*D cohort. <i>American Journal on Addictions</i> , 2006 , 15, 278-85	3.7	70
92	Positron emission tomography imaging of peripheral benzodiazepine receptor binding in human immunodeficiency virus-infected subjects with and without cognitive impairment. <i>Journal of NeuroVirology</i> , 2006 , 12, 262-71	3.9	46
91	Towards a noninvasive method for determination of patient-specific wall strength distribution in abdominal aortic aneurysms. <i>Annals of Biomedical Engineering</i> , 2006 , 34, 1098-106	4.7	151
90	A comparison of mirtazapine and nortriptyline following two consecutive failed medication treatments for depressed outpatients: a STAR*D report. <i>American Journal of Psychiatry</i> , 2006 , 163, 116	1 -72 9	100
89	Are antidepressants associated with new-onset suicidality in bipolar disorder? A prospective study of participants in the Systematic Treatment Enhancement Program for Bipolar Disorder (STEP-BD). <i>Journal of Clinical Psychiatry</i> , 2006 , 67, 48-55	4.6	61
88	Children of currently depressed mothers: a STAR*D ancillary study. <i>Journal of Clinical Psychiatry</i> , 2006 , 67, 126-36	4.6	103
87	Factors associated with health-related quality of life among outpatients with major depressive disorder: a STAR*D report. <i>Journal of Clinical Psychiatry</i> , 2006 , 67, 185-95	4.6	113
86	Baseline predictors of serious adverse events at one year among patients with bipolar disorder in STEP-BD. <i>Psychiatric Services</i> , 2005 , 56, 1541-8	3.3	18
85	Contributors to suicidal ideation among bipolar patients with and without a history of suicide attempts. <i>Suicide and Life-Threatening Behavior</i> , 2005 , 35, 671-80	3.9	27
84	Age-Related Characteristics of Depression: A Preliminary STAR*D Report. <i>American Journal of Geriatric Psychiatry</i> , 2005 , 13, 852-860	6.5	42
83	Quality improvement methods as applied to a multicenter effectiveness trialSTAR*D. <i>Contemporary Clinical Trials</i> , 2005 , 26, 95-112	2.3	6
82	Substance use disorder comorbidity in major depressive disorder: an exploratory analysis of the Sequenced Treatment Alternatives to Relieve Depression cohort. <i>Comprehensive Psychiatry</i> , 2005 , 46, 81-9	7.3	65
81	Care satisfaction, hope, and life functioning among adults with bipolar disorder: data from the first 1000 participants in the Systematic Treatment Enhancement Program. <i>Comprehensive Psychiatry</i> , 2005 , 46, 98-104	7.3	20

80	Response to Zhang et al. (2005): loss-of-function mutation in tryptophan hydroxylase-2 identified in unipolar major depression. Neuron 45, 11-16. <i>Neuron</i> , 2005 , 48, 702-3; author reply 705-6	13.9	25
79	Perceived criticism from family members as a predictor of the one-year course of bipolar disorder. <i>Psychiatry Research</i> , 2005 , 136, 101-11	9.9	78
78	Clinical and diagnostic implications of lifetime attention-deficit/hyperactivity disorder comorbidity in adults with bipolar disorder: data from the first 1000 STEP-BD participants. <i>Biological Psychiatry</i> , 2005 , 57, 1467-73	7.9	206
77	Revisiting depressive-prone bipolar disorder: polarity of initial mood episode and disease course among bipolar I systematic treatment enhancement program for bipolar disorder participants. <i>Biological Psychiatry</i> , 2005 , 58, 549-53	7.9	64
76	Mild hypothermia improves survival after prolonged, traumatic hemorrhagic shock in pigs. <i>Journal of Trauma</i> , 2005 , 59, 291-9; discussion 299-301		33
75	Gender differences in bipolar disorder: retrospective data from the first 500 STEP-BD participants. <i>Bipolar Disorders</i> , 2005 , 7, 465-70	3.8	91
74	Cytochrome c, a biomarker of apoptosis, is increased in cerebrospinal fluid from infants with inflicted brain injury from child abuse. <i>Journal of Cerebral Blood Flow and Metabolism</i> , 2005 , 25, 919-27	7.3	81
73	Characterization of the effects of adenosine receptor agonists on cerebral blood flow in uninjured and traumatically injured rat brain using continuous arterial spin-labeled magnetic resonance imaging. <i>Journal of Cerebral Blood Flow and Metabolism</i> , 2005 , 25, 1596-612	7.3	29
72	Gender differences in depression: findings from the STAR*D study. <i>Journal of Affective Disorders</i> , 2005 , 87, 141-50	6.6	292
71	Comorbid psychiatric disorders in depressed outpatients: demographic and clinical features. Journal of Affective Disorders, 2005 , 87, 43-55	6.6	185
70	Development of an efficient SAS macro to perform permutation tests for two independent samples. <i>Computer Methods and Programs in Biomedicine</i> , 2005 , 79, 179-87	6.9	2
69	A direct comparison of presenting characteristics of depressed outpatients from primary vs. specialty care settings: preliminary findings from the STAR*D clinical trial. <i>General Hospital Psychiatry</i> , 2005 , 27, 87-96	5.6	56
68	Prevalence and correlates of tobacco use in bipolar disorder: data from the first 2000 participants in the Systematic Treatment Enhancement Program. <i>General Hospital Psychiatry</i> , 2005 , 27, 321-8	5.6	101
67	Characteristics of insured and noninsured outpatients with depression in STAR(*)D. <i>Psychiatric Services</i> , 2005 , 56, 995-1004	3.3	25
66	Suicidal ideation and pharmacotherapy among STEP-BD patients. <i>Psychiatric Services</i> , 2005 , 56, 1534-40	3.3	19
65	Age-related characteristics of depression: a preliminary STAR*D report. <i>American Journal of Geriatric Psychiatry</i> , 2005 , 13, 852-60	6.5	25
64	Prevalence and Clinical Correlates of Irritability in Major Depressive Disorder. <i>Journal of Clinical Psychiatry</i> , 2005 , 66, 159-166	4.6	53
63	Does recovery from substance use disorder matter in patients with bipolar disorder?. <i>Journal of Clinical Psychiatry</i> , 2005 , 66, 730-5; quiz 808-9	4.6	94

(2004-2005)

62	Clinical and demographic features of atypical depression in outpatients with major depressive disorder: preliminary findings from STAR*D. <i>Journal of Clinical Psychiatry</i> , 2005 , 66, 1002-11	4.6	133
61	Prevalence and clinical correlates of irritability in major depressive disorder: a preliminary report from the Sequenced Treatment Alternatives to Relieve Depression study. <i>Journal of Clinical Psychiatry</i> , 2005 , 66, 159-66; quiz 147, 273-4	4.6	20
60	Continuous versus intermittent cerebrospinal fluid drainage after severe traumatic brain injury in children: effect on biochemical markers. <i>Journal of Neurotrauma</i> , 2004 , 21, 1113-22	5.4	80
59	Positive Aspects of Caregiving: Contributions of the REACH Project to the Development of New Measures for Alzheimer Caregiving. <i>Research on Aging</i> , 2004 , 26, 429-453	3	308
58	Assessment of the macrophage marker quinolinic acid in cerebrospinal fluid after pediatric traumatic brain injury: insight into the timing and severity of injury in child abuse. <i>Journal of Neurotrauma</i> , 2004 , 21, 1123-30	5.4	26
57	Marked gender effect on lipid peroxidation after severe traumatic brain injury in adult patients. Journal of Neurotrauma, 2004 , 21, 1-8	5.4	175
56	Anxiety disorder comorbidity in bipolar disorder patients: data from the first 500 participants in the Systematic Treatment Enhancement Program for Bipolar Disorder (STEP-BD). <i>American Journal of Psychiatry</i> , 2004 , 161, 2222-9	11.9	423
55	Web-based communications and management of a multi-center clinical trial: the Sequenced Treatment Alternatives to Relieve Depression (STAR*D) project. <i>Clinical Trials</i> , 2004 , 1, 387-98	2.2	38
54	Demographic and diagnostic characteristics of the first 1000 patients enrolled in the Systematic Treatment Enhancement Program for Bipolar Disorder (STEP-BD). <i>Bipolar Disorders</i> , 2004 , 6, 460-9	3.8	97
53	Methods of testing feasibility for sequenced treatment alternatives to relieve depression (STAR*D). <i>Journal of Psychiatric Research</i> , 2004 , 38, 241-8	5.2	6
52	Suicide risk management for the sequenced treatment alternatives to relieve depression study: applied NIMH guidelines. <i>Journal of Psychiatric Research</i> , 2004 , 38, 583-9	5.2	22
51	Sequenced treatment alternatives to relieve depression (STAR*D): rationale and design. <i>Contemporary Clinical Trials</i> , 2004 , 25, 119-42		725
50	Clinical features of depressed outpatients with and without co-occurring general medical conditions in STAR*D. <i>General Hospital Psychiatry</i> , 2004 , 26, 421-9	5.6	83
49	The Implementation of Cognitive Therapy in STAR*D. Cognitive Therapy and Research, 2004, 28, 819-833	3 2.7	5
48	Bone mineral density and prevalent vertebral fractures in men and women. <i>Osteoporosis International</i> , 2004 , 15, 32-7	5.3	46
47	Long-term implications of early onset in bipolar disorder: data from the first 1000 participants in the systematic treatment enhancement program for bipolar disorder (STEP-BD). <i>Biological Psychiatry</i> , 2004 , 55, 875-81	7.9	618
46	Alzheimer's pathology in human temporal cortex surgically excised after severe brain injury. <i>Experimental Neurology</i> , 2004 , 190, 192-203	5.7	302
45	Factors that differentiate early vs. later onset of major depression disorder. <i>Psychiatry Research</i> , 2004 , 129, 127-40	9.9	103

44	Pharmacotherapy for bipolar disorder and comorbid conditions: baseline data from STEP-BD. Journal of Clinical Psychopharmacology, 2004 , 24, 512-20	1.7	71
43	Vascular endothelial growth factor is increased in cerebrospinal fluid after traumatic brain injury in infants and children. <i>Neurosurgery</i> , 2004 , 54, 605-11; discussion 611-2	3.2	47
42	Phenomenology of rapid-cycling bipolar disorder: data from the first 500 participants in the Systematic Treatment Enhancement Program. <i>American Journal of Psychiatry</i> , 2004 , 161, 1902-8	11.9	85
41	Clinical correlates and symptom patterns of anxious depression among patients with major depressive disorder in STAR*D. <i>Psychological Medicine</i> , 2004 , 34, 1299-308	6.9	261
40	Psychosocial service utilization by patients with bipolar disorders: data from the first 500 participants in the Systematic Treatment Enhancement Program. <i>Journal of Psychiatric Practice</i> , 2004 , 10, 81-7	1.3	18
39	PET imaging of brain macrophages using the peripheral benzodiazepine receptor in a macaque model of neuroAIDS. <i>Journal of Clinical Investigation</i> , 2004 , 113, 981-989	15.9	34
38	PET imaging of brain macrophages using the peripheral benzodiazepine receptor in a macaque model of neuroAIDS. <i>Journal of Clinical Investigation</i> , 2004 , 113, 981-9	15.9	30
37	Acute etomidate treatment reduces cognitive deficits and histopathology in rats with traumatic brain injury. <i>Critical Care Medicine</i> , 2003 , 31, 2222-7	1.4	33
36	Introduction to the special section on Resources for Enhancing Alzheimer's Caregiver Health (REACH). <i>Psychology and Aging</i> , 2003 , 18, 357-60	3.6	60
35	The Resources for Enhancing Alzheimer's Caregiver Health (REACH): project design and baseline characteristics. <i>Psychology and Aging</i> , 2003 , 18, 375-84	3.6	171
34	Plasma and cerebrospinal fluid alpha1-antichymotrypsin levels in Alzheimer's disease: correlation with cognitive impairment. <i>Annals of Neurology</i> , 2003 , 53, 81-90	9.4	75
33	F2-isoprostane and neuron-specific enolase in cerebrospinal fluid after severe traumatic brain injury in infants and children. <i>Journal of Neurotrauma</i> , 2003 , 20, 781-6	5.4	74
32	Psychotic Symptoms in Alzheimer Disease: Evidence for Subtypes. <i>American Journal of Geriatric Psychiatry</i> , 2003 , 11, 406-413	6.5	48
31	Rationale, design, and methods of the systematic treatment enhancement program for bipolar disorder (STEP-BD). <i>Biological Psychiatry</i> , 2003 , 53, 1028-42	7.9	421
30	Upregulation of choline acetyltransferase activity in hippocampus and frontal cortex of elderly subjects with mild cognitive impairment. <i>Annals of Neurology</i> , 2002 , 51, 145-55	9.4	553
29	Apolipoprotein E and alpha-1-antichymotrypsin genotypes do not predict time to psychosis in Alzheimer's disease. <i>Journal of Geriatric Psychiatry and Neurology</i> , 2002 , 15, 24-30	3.8	16
28	Neuron-specific enolase and S100B in cerebrospinal fluid after severe traumatic brain injury in infants and children. <i>Pediatrics</i> , 2002 , 109, E31	7.4	169
27	Diabetes and postcataract extraction endophthalmitis. Current Opinion in Ophthalmology, 2002, 13, 147	-5.1	17

26	Cerebrospinal fluid procalcitonin and severe traumatic brain injury in children. <i>Pediatric Critical Care Medicine</i> , 2002 , 3, 39-44	3	8
25	Increased adenosine in cerebrospinal fluid after severe traumatic brain injury in infants and children: association with severity of injury and excitotoxicity. <i>Critical Care Medicine</i> , 2001 , 29, 2287-93	1.4	59
24	Alterations of striatal dopamine receptor binding in Alzheimer disease are associated with Lewy body pathology and antemortem psychosis. <i>Archives of Neurology</i> , 2001 , 58, 466-72		56
23	Strengthening clinical effectiveness trials: equipoise-stratified randomization. <i>Biological Psychiatry</i> , 2001 , 50, 792-801	7.9	124
22	Excitatory amino acid concentrations in ventricular cerebrospinal fluid after severe traumatic brain injury in infants and children: the role of child abuse. <i>Journal of Pediatrics</i> , 2001 , 138, 18-25	3.6	116
21	The Th1 versus Th2 cytokine profile in cerebrospinal fluid after severe traumatic brain injury in infants and children. <i>Pediatric Critical Care Medicine</i> , 2001 , 2, 260-264	3	38
20	No long-term benefit from hypothermia after severe traumatic brain injury with secondary insult in rats. <i>Critical Care Medicine</i> , 2000 , 28, 3218-23	1.4	35
19	Psychotic symptoms in Alzheimer's disease are not associated with more severe neuropathologic features. <i>International Psychogeriatrics</i> , 2000 , 12, 547-58	3.4	48
18	Interleukin-8 is increased in cerebrospinal fluid of children with severe head injury. <i>Critical Care Medicine</i> , 2000 , 28, 929-34	1.4	158
17	Isoflurane improves long-term neurologic outcome versus fentanyl after traumatic brain injury in rats. <i>Journal of Neurotrauma</i> , 2000 , 17, 1179-89	5.4	82
16	Increases in bcl-2 protein in cerebrospinal fluid and evidence for programmed cell death in infants and children after severe traumatic brain injury. <i>Journal of Pediatrics</i> , 2000 , 137, 197-204	3.6	94
15	Inducible nitric oxide synthase is an endogenous neuroprotectant after traumatic brain injury in rats and mice. <i>Journal of Clinical Investigation</i> , 1999 , 104, 647-56	15.9	179
14	Psychiatric medication and abnormal behavior as predictors of progression in probable Alzheimer disease. <i>Archives of Neurology</i> , 1999 , 56, 1266-72		160
13	Factors affecting survival rates for acute vertebrobasilar artery occlusions treated with intra-arterial thrombolytic therapy: a meta-analytical approach. <i>Neurosurgery</i> , 1999 , 45, 539-45; discussion 545-8	3.2	86
12	Apolipoprotein E polymorphism: A new genetic marker of hip fracture riskThe Study of Osteoporotic Fractures. <i>Journal of Bone and Mineral Research</i> , 1999 , 14, 1175-81	6.3	86
11	Quinolinic acid in the cerebrospinal fluid of children after traumatic brain injury. <i>Critical Care Medicine</i> , 1999 , 27, 493-7	1.4	50
10	Quinolinic acid is increased in CSF and associated with mortality after traumatic brain injury in humans. <i>Journal of Cerebral Blood Flow and Metabolism</i> , 1998 , 18, 610-5	7.3	57
9	Genetic association of two chromosome 14 genes (presenilin 1 and alpha 1-antichymotrypsin) with Alzheimer's disease. <i>Annals of Neurology</i> , 1998 , 44, 387-90	9.4	24

8	Additional procedures after the initial vitrectomy or tap-biopsy in the Endophthalmitis Vitrectomy Study. <i>Ophthalmology</i> , 1998 , 105, 707-16	7.3	55
7	Soluble adhesion molecules in CSF are increased in children with severe head injury. <i>Journal of Neurotrauma</i> , 1998 , 15, 777-87	5.4	69
6	Interstitial adenosine, inosine, and hypoxanthine are increased after experimental traumatic brain injury in the rat. <i>Journal of Neurotrauma</i> , 1998 , 15, 163-70	5.4	84
5	Cerebrovascular response in infants and young children following severe traumatic brain injury: a preliminary report. <i>Pediatric Neurosurgery</i> , 1997 , 26, 200-7	0.9	187
4	Treatment of traumatic brain injury with moderate hypothermia. <i>New England Journal of Medicine</i> , 1997 , 336, 540-6	59.2	1137
3	Assessment of cerebral blood flow and CO2 reactivity after controlled cortical impact by perfusion magnetic resonance imaging using arterial spin-labeling in rats. <i>Journal of Cerebral Blood Flow and Metabolism</i> , 1997 , 17, 865-74	7.3	68
2	Cerebrospinal fluid and plasma nitrite and nitrate concentrations after head injury in humans. <i>Critical Care Medicine</i> , 1996 , 24, 1243-51	1.4	96
1	Quantitative computer-assisted image analysis of suction biopsy in pediatric gastroesophageal reflux. <i>Pediatric Pathology</i> , 1994 , 14, 653-64		13