

Yung-Ming Jeng

List of Publications by Citations

Source: <https://exaly.com/author-pdf/4252767/yung-ming-jeng-publications-by-citations.pdf>
Version: 2024-04-09

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

256 papers	11,285 citations	58 h-index	93 g-index
267 ext. papers	13,024 ext. citations	6.8 avg, IF	6 L-index

#	Paper	IF	Citations
256	B cells are associated with survival and immunotherapy response in sarcoma. <i>Nature</i> , 2020 , 577, 556-560	50.4	538
255	Elastic modulus measurements of human liver and correlation with pathology. <i>Ultrasound in Medicine and Biology</i> , 2002 , 28, 467-74	3.5	314
254	H3K9 histone methyltransferase G9a promotes lung cancer invasion and metastasis by silencing the cell adhesion molecule Ep-CAM. <i>Cancer Research</i> , 2010 , 70, 7830-40	10.1	274
253	The VEGF-C/Flt-4 axis promotes invasion and metastasis of cancer cells. <i>Cancer Cell</i> , 2006 , 9, 209-23	24.3	273
252	Beta-catenin mutations are associated with a subset of low-stage hepatocellular carcinoma negative for hepatitis B virus and with favorable prognosis. <i>American Journal of Pathology</i> , 2000 , 157, 763-70	5.8	256
251	Overexpression and amplification of Aurora-A in hepatocellular carcinoma. <i>Clinical Cancer Research</i> , 2004 , 10, 2065-71	12.9	253
250	TET1 suppresses cancer invasion by activating the tissue inhibitors of metalloproteinases. <i>Cell Reports</i> , 2012 , 2, 568-79	10.6	214
249	Mutation and loss of expression of ARID1A in uterine low-grade endometrioid carcinoma. <i>American Journal of Surgical Pathology</i> , 2011 , 35, 625-32	6.7	206
248	Peginterferon alfa-2a plus ribavirin for the treatment of dual chronic infection with hepatitis B and C viruses. <i>Gastroenterology</i> , 2009 , 136, 496-504.e3	13.3	183
247	Clinical relevance of hepatitis B virus genotype in children with chronic infection and hepatocellular carcinoma. <i>Gastroenterology</i> , 2004 , 127, 1733-8	13.3	173
246	Cyclooxygenase-2 induces EP1- and HER-2/Neu-dependent vascular endothelial growth factor-C up-regulation: a novel mechanism of lymphangiogenesis in lung adenocarcinoma. <i>Cancer Research</i> , 2004 , 64, 554-64	10.1	172
245	Multiple lineages of human breast cancer stem/progenitor cells identified by profiling with stem cell markers. <i>PLoS ONE</i> , 2009 , 4, e8377	3.7	169
244	MicroRNA-519c suppresses hypoxia-inducible factor-1alpha expression and tumor angiogenesis. <i>Cancer Research</i> , 2010 , 70, 2675-85	10.1	163
243	Expression of the c-kit protein is associated with certain subtypes of salivary gland carcinoma. <i>Cancer Letters</i> , 2000 , 154, 107-11	9.9	161
242	Morphological subclassification of intrahepatic cholangiocarcinoma: etiological, clinicopathological, and molecular features. <i>Modern Pathology</i> , 2014 , 27, 1163-73	9.8	146
241	Connective tissue growth factor and its role in lung adenocarcinoma invasion and metastasis. <i>Journal of the National Cancer Institute</i> , 2004 , 96, 364-75	9.7	138
240	Oncologic results of laparoscopic versus conventional open surgery for stage II or III left-sided colon cancers: a randomized controlled trial. <i>Annals of Surgical Oncology</i> , 2007 , 14, 109-17	3.1	134

239	Sinonasal undifferentiated carcinoma and nasopharyngeal-type undifferentiated carcinoma: two clinically, biologically, and histopathologically distinct entities. <i>American Journal of Surgical Pathology</i> , 2002 , 26, 371-6	6.7	124
238	RNA-binding protein insulin-like growth factor II mRNA-binding protein 3 expression promotes tumor invasion and predicts early recurrence and poor prognosis in hepatocellular carcinoma. <i>Hepatology</i> , 2008 , 48, 1118-27	11.2	121
237	SIRT1 promotes tumorigenesis and resistance to chemotherapy in hepatocellular carcinoma and its expression predicts poor prognosis. <i>Annals of Surgical Oncology</i> , 2012 , 19, 2011-9	3.1	116
236	Histone demethylase RBP2 promotes lung tumorigenesis and cancer metastasis. <i>Cancer Research</i> , 2013 , 73, 4711-21	10.1	113
235	Connective tissue growth factor inhibits metastasis and acts as an independent prognostic marker in colorectal cancer. <i>Gastroenterology</i> , 2005 , 128, 9-23	13.3	108
234	IL-6 induces AGS gastric cancer cell invasion via activation of the c-Src/RhoA/ROCK signaling pathway. <i>International Journal of Cancer</i> , 2007 , 120, 2600-8	7.5	107
233	Loss of the oxidative stress sensor NPGPx compromises GRP78 chaperone activity and induces systemic disease. <i>Molecular Cell</i> , 2012 , 48, 747-59	17.6	106
232	Management of hepatic angiolipoma. <i>Journal of Gastrointestinal Surgery</i> , 2007 , 11, 452-7	3.3	105
231	P53 gene and Wnt signaling in benign neoplasms: beta-catenin mutations in hepatic adenoma but not in focal nodular hyperplasia. <i>Hepatology</i> , 2002 , 36, 927-35	11.2	103
230	The H3K9 methyltransferase G9a is a marker of aggressive ovarian cancer that promotes peritoneal metastasis. <i>Molecular Cancer</i> , 2014 , 13, 189	42.1	102
229	Activation of Robo1 signaling of breast cancer cells by Slit2 from stromal fibroblast restrains tumorigenesis via blocking PI3K/Akt/E-catenin pathway. <i>Cancer Research</i> , 2012 , 72, 4652-61	10.1	100
228	Somatic mutations of beta-catenin play a crucial role in the tumorigenesis of sporadic hepatoblastoma. <i>Cancer Letters</i> , 2000 , 152, 45-51	9.9	95
227	Ribavirin and interferon is effective for hepatitis C virus clearance in hepatitis B and C dually infected patients. <i>Hepatology</i> , 2003 , 37, 568-76	11.2	94
226	Expression of mutant nuclear beta-catenin correlates with non-invasive hepatocellular carcinoma, absence of portal vein spread, and good prognosis. <i>Journal of Pathology</i> , 2001 , 193, 95-101	9.4	94
225	ASPM is a novel marker for vascular invasion, early recurrence, and poor prognosis of hepatocellular carcinoma. <i>Clinical Cancer Research</i> , 2008 , 14, 4814-20	12.9	92
224	Loss of corepressor PER2 under hypoxia up-regulates OCT1-mediated EMT gene expression and enhances tumor malignancy. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013 , 110, 12331-6	11.5	91
223	Identification of a novel FN1-FGFR1 genetic fusion as a frequent event in phosphaturic mesenchymal tumour. <i>Journal of Pathology</i> , 2015 , 235, 539-45	9.4	90
222	Identification of calreticulin as a prognosis marker and angiogenic regulator in human gastric cancer. <i>Annals of Surgical Oncology</i> , 2009 , 16, 524-33	3.1	90

221	Targeting IL-17B-IL-17RB signaling with an anti-IL-17RB antibody blocks pancreatic cancer metastasis by silencing multiple chemokines. <i>Journal of Experimental Medicine</i> , 2015 , 212, 333-49	16.6	89
220	Cadherin-17 is a useful diagnostic marker for adenocarcinomas of the digestive system. <i>Modern Pathology</i> , 2008 , 21, 1379-86	9.8	87
219	KLF6, a putative tumor suppressor gene, is mutated in astrocytic gliomas. <i>International Journal of Cancer</i> , 2003 , 105, 625-9	7.5	87
218	Lin-28B expression promotes transformation and invasion in human hepatocellular carcinoma. <i>Carcinogenesis</i> , 2010 , 31, 1516-22	4.6	85
217	Removal of BRCA1/CtIP/ZBRK1 repressor complex on ANG1 promoter leads to accelerated mammary tumor growth contributed by prominent vasculature. <i>Cancer Cell</i> , 2006 , 10, 13-24	24.3	84
216	Significance of Aurora B overexpression in hepatocellular carcinoma. Aurora B Overexpression in HCC. <i>BMC Cancer</i> , 2010 , 10, 461	4.8	83
215	Cyr61 induces gastric cancer cell motility/invasion via activation of the integrin/nuclear factor-kappaB/cyclooxygenase-2 signaling pathway. <i>Clinical Cancer Research</i> , 2005 , 11, 5809-20	12.9	82
214	Gas6/Axl pathway promotes tumor invasion through the transcriptional activation of Slug in hepatocellular carcinoma. <i>Carcinogenesis</i> , 2014 , 35, 769-75	4.6	81
213	FIC1 and BSEP defects in Taiwanese patients with chronic intrahepatic cholestasis with low gamma-glutamyltranspeptidase levels. <i>Journal of Pediatrics</i> , 2002 , 140, 119-24	3.6	77
212	IL-25 causes apoptosis of IL-25R-expressing breast cancer cells without toxicity to nonmalignant cells. <i>Science Translational Medicine</i> , 2011 , 3, 78ra31	17.5	75
211	Clinical Significance of Circulating Tumor Microemboli as a Prognostic Marker in Patients with Pancreatic Ductal Adenocarcinoma. <i>Clinical Chemistry</i> , 2016 , 62, 505-13	5.5	69
210	Expression of hepatocyte transporters and nuclear receptors in children with early and late-stage biliary atresia. <i>Pediatric Research</i> , 2008 , 63, 667-73	3.2	68
209	Lymphoepithelioma-like cholangiocarcinoma: an Epstein-Barr virus-associated tumor. <i>American Journal of Surgical Pathology</i> , 2001 , 25, 516-20	6.7	67
208	Down-regulation of annexin A10 in hepatocellular carcinoma is associated with vascular invasion, early recurrence, and poor prognosis in synergy with p53 mutation. <i>American Journal of Pathology</i> , 2002 , 160, 1831-7	5.8	66
207	Effect of connective tissue growth factor on hypoxia-inducible factor 1alpha degradation and tumor angiogenesis. <i>Journal of the National Cancer Institute</i> , 2006 , 98, 984-95	9.7	65
206	Angiopoietin-like protein 1 suppresses SLUG to inhibit cancer cell motility. <i>Journal of Clinical Investigation</i> , 2013 , 123, 1082-95	15.9	65
205	Leiomyosarcoma with alternative lengthening of telomeres is associated with aggressive histologic features, loss of ATRX expression, and poor clinical outcome. <i>American Journal of Surgical Pathology</i> , 2015 , 39, 236-44	6.7	64
204	Differential DNA methylation associated with hepatitis B virus infection in hepatocellular carcinoma. <i>International Journal of Cancer</i> , 2007 , 121, 1257-64	7.5	63

203	Overexpression of KIAA0101 predicts high stage, early tumor recurrence, and poor prognosis of hepatocellular carcinoma. <i>Clinical Cancer Research</i> , 2007 , 13, 5368-76	12.9	63
202	Overexpression of CTHRC1 in hepatocellular carcinoma promotes tumor invasion and predicts poor prognosis. <i>PLoS ONE</i> , 2013 , 8, e70324	3.7	62
201	Comparison of medial-to-lateral versus traditional lateral-to-medial laparoscopic dissection sequences for resection of rectosigmoid cancers: randomized controlled clinical trial. <i>World Journal of Surgery</i> , 2003 , 27, 190-6	3.3	60
200	Serum adiponectin correlates with viral characteristics but not histologic features in patients with chronic hepatitis C. <i>Journal of Hepatology</i> , 2005 , 43, 235-42	13.4	59
199	Adenylate kinase-4 is a marker of poor clinical outcomes that promotes metastasis of lung cancer by downregulating the transcription factor ATF3. <i>Cancer Research</i> , 2012 , 72, 5119-29	10.1	58
198	Knockdown of contactin-1 expression suppresses invasion and metastasis of lung adenocarcinoma. <i>Cancer Research</i> , 2006 , 66, 2553-61	10.1	57
197	Traditional serrated adenoma has two pathways of neoplastic progression that are distinct from the sessile serrated pathway of colorectal carcinogenesis. <i>Modern Pathology</i> , 2014 , 27, 1375-85	9.8	55
196	Role of p53 and Eatenin mutations in conjunction with CK19 expression on early tumor recurrence and prognosis of hepatocellular carcinoma. <i>Journal of Gastrointestinal Surgery</i> , 2011 , 15, 321-9	3.3	53
195	Solid pseudopapillary neoplasms of the pancreas: is there a pathologic basis for the observed gender differences in incidence?. <i>Surgery</i> , 2005 , 137, 591-6	3.6	53
194	Adipocytes promote malignant growth of breast tumours with monocarboxylate transporter 2 expression via hydroxybutyrate. <i>Nature Communications</i> , 2017 , 8, 14706	17.4	50
193	Cytoplasmic CD24 expression is a novel prognostic factor in diffuse-type gastric adenocarcinoma. <i>Annals of Surgical Oncology</i> , 2007 , 14, 2748-58	3.1	50
192	Sorafenib and its derivative SC-1 exhibit antifibrotic effects through signal transducer and activator of transcription 3 inhibition. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2015 , 112, 7243-8	11.5	49
191	Viral hepatitis is associated with intrahepatic cholangiocarcinoma with cholangiolar differentiation and N-cadherin expression. <i>Modern Pathology</i> , 2011 , 24, 810-9	9.8	49
190	TERT promoter mutation in resectable hepatocellular carcinomas: a strong association with hepatitis C infection and absence of hepatitis B infection. <i>International Journal of Surgery</i> , 2014 , 12, 659-65	7.5	47
189	Post-transplantation lymphoproliferative disorders localizing to the gastrointestinal tract after liver transplantation: report of five pediatric cases. <i>Pediatric Transplantation</i> , 2006 , 10, 390-4	1.8	47
188	CD24 expression is a prognostic factor in intrahepatic cholangiocarcinoma. <i>Cancer Letters</i> , 2006 , 235, 34-9	9.9	46
187	Comprehensive screening of alternative lengthening of telomeres phenotype and loss of ATRX expression in sarcomas. <i>Modern Pathology</i> , 2015 , 28, 1545-54	9.8	45
186	Transient elastography is useful in diagnosing biliary atresia and predicting prognosis after hepatoportoenterostomy. <i>Hepatology</i> , 2018 , 68, 616-624	11.2	45

185	HSD3B1 as a novel trophoblast-associated marker that assists in the differential diagnosis of trophoblastic tumors and tumorlike lesions. <i>American Journal of Surgical Pathology</i> , 2008 , 32, 236-42	6.7	45
184	Absence of epidermal growth factor receptor exon 18-21 mutation in hepatocellular carcinoma. <i>Cancer Letters</i> , 2005 , 224, 117-21	9.9	45
183	Diffuse expression of RNA-binding protein IMP3 predicts high-stage lymph node metastasis and poor prognosis in colorectal adenocarcinoma. <i>Annals of Surgical Oncology</i> , 2009 , 16, 1711-9	3.1	44
182	Nuclear localization of annexin A1 is a prognostic factor in oral squamous cell carcinoma. <i>Journal of Surgical Oncology</i> , 2008 , 97, 544-50	2.8	44
181	HMB-45 may be a more sensitive maker than S-100 or Melan-A for immunohistochemical diagnosis of primary oral and nasal mucosal melanomas. <i>Journal of Oral Pathology and Medicine</i> , 2005 , 34, 540-5	3.3	44
180	GRP78 expression correlates with histologic differentiation and favorable prognosis in neuroblastic tumors. <i>International Journal of Cancer</i> , 2005 , 113, 920-7	7.5	42
179	High Glucose Triggers Nucleotide Imbalance through O-GlcNAcylation of Key Enzymes and Induces KRAS Mutation in Pancreatic Cells. <i>Cell Metabolism</i> , 2019 , 29, 1334-1349.e10	24.6	40
178	RBMV, a novel inhibitor of glycogen synthase kinase 3 β increases tumor stemness and predicts poor prognosis of hepatocellular carcinoma. <i>Hepatology</i> , 2015 , 62, 1480-96	11.2	40
177	Hepatocyte growth factor activates Wnt pathway by transcriptional activation of LEF1 to facilitate tumor invasion. <i>Carcinogenesis</i> , 2012 , 33, 1142-8	4.6	40
176	Targeted next-generation sequencing of cancer genes identified frequent TP53 and ATRX mutations in leiomyosarcoma. <i>American Journal of Translational Research (discontinued)</i> , 2015 , 7, 2072-81		40
175	The Aurora kinase inhibitor VE-465 has anticancer effects in pre-clinical studies of human hepatocellular carcinoma. <i>Journal of Hepatology</i> , 2009 , 50, 518-27	13.4	39
174	Frequent alterations of HER2 through mutation, amplification, or overexpression in pleomorphic lobular carcinoma of the breast. <i>Breast Cancer Research and Treatment</i> , 2015 , 150, 447-55	4.4	38
173	Clinical Manifestations and Outcomes of Patients with Sarcomatoid Hepatocellular Carcinoma. <i>Hepatology</i> , 2019 , 69, 209-221	11.2	38
172	Clinical features and treatment responses of children with eosinophilic gastroenteritis. <i>Pediatrics and Neonatology</i> , 2011 , 52, 272-8	1.8	38
171	Alternative lengthening of telomeres phenotype in malignant vascular tumors is highly associated with loss of ATRX expression and is frequently observed in hepatic angiosarcomas. <i>Human Pathology</i> , 2015 , 46, 1360-6	3.7	37
170	Glutaminase 2 stabilizes Dicer to repress Snail and metastasis in hepatocellular carcinoma cells. <i>Cancer Letters</i> , 2016 , 383, 282-294	9.9	37
169	Resolution of diabetes after pancreaticoduodenectomy in patients with and without pancreatic ductal cell adenocarcinoma. <i>Annals of Surgical Oncology</i> , 2013 , 20, 242-9	3.1	37
168	Desmoplastic small round cell tumor of the kidney. <i>American Journal of Surgical Pathology</i> , 2004 , 28, 1379-83	6.7	37

167	Opposite roles of human pancreatitis-associated protein and REG1A expression in hepatocellular carcinoma: association of pancreatitis-associated protein expression with low-stage hepatocellular carcinoma, beta-catenin mutation, and favorable prognosis. <i>Clinical Cancer Research</i> , 2005 , 11, 2568-75	12.9	37
166	Alternative lengthening of telomeres and loss of ATRX are frequent events in pleomorphic and dedifferentiated liposarcomas. <i>Modern Pathology</i> , 2015 , 28, 1064-73	9.8	36
165	Leukocyte cell-derived chemotaxin 2 antagonizes MET receptor activation to suppress hepatocellular carcinoma vascular invasion by protein tyrosine phosphatase 1B recruitment. <i>Hepatology</i> , 2014 , 59, 974-85	11.2	36
164	GNAS Is frequently mutated in a specific subgroup of intraductal papillary neoplasms of the bile duct. <i>American Journal of Surgical Pathology</i> , 2013 , 37, 1862-70	6.7	36
163	Unexpected rapid progression of metastatic adenoid cystic carcinoma during treatment with imatinib mesylate. <i>Head and Neck</i> , 2005 , 27, 1022-7	4.2	36
162	TERT promoter mutation is uncommon in acral lentiginous melanoma. <i>Journal of Cutaneous Pathology</i> , 2014 , 41, 504-8	1.7	35
161	Notch1 expression predicts an unfavorable prognosis and serves as a therapeutic target of patients with neuroblastoma. <i>Clinical Cancer Research</i> , 2010 , 16, 4411-20	12.9	35
160	Multiple intrahepatic biliary cysts in children with biliary atresia. <i>Journal of Pediatric Surgery</i> , 2002 , 37, 1183-7	2.6	35
159	β1,4-Galactosyltransferase III enhances invasive phenotypes via β1-integrin and predicts poor prognosis in neuroblastoma. <i>Clinical Cancer Research</i> , 2013 , 19, 1705-16	12.9	34
158	RBMY, a male germ cell-specific RNA-binding protein, activated in human liver cancers and transforms rodent fibroblasts. <i>Oncogene</i> , 2004 , 23, 5815-22	9.2	33
157	CTP synthase forms the cytoophidium in human hepatocellular carcinoma. <i>Experimental Cell Research</i> , 2017 , 361, 292-299	4.2	32
156	Hepatic steatosis and neonatal intrahepatic cholestasis caused by citrin deficiency (NICCD) in Taiwanese infants. <i>Journal of Pediatrics</i> , 2006 , 148, 642-6	3.6	32
155	Utility of hepatocyte nuclear factor-1β as a diagnostic marker in ovarian carcinomas with clear cells. <i>Histopathology</i> , 2012 , 61, 760-8	7.3	31
154	p15(PAF) is an Rb/E2F-regulated S-phase protein essential for DNA synthesis and cell cycle progression. <i>PLoS ONE</i> , 2013 , 8, e61196	3.7	31
153	Depletion of β-catenin from mature hepatocytes of mice promotes expansion of hepatic progenitor cells and tumor development. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011 , 108, 18384-9	11.5	31
152	Progressive familial intrahepatic cholestasis with high gamma-glutamyltranspeptidase levels in Taiwanese infants: role of MDR3 gene defect?. <i>Pediatric Research</i> , 2001 , 50, 50-5	3.2	31
151	Differential expression of hyaluronan synthase 2 in breast carcinoma and its biological significance. <i>Histopathology</i> , 2014 , 65, 328-39	7.3	29
150	S100P immunostaining identifies a subset of peripheral-type intrahepatic cholangiocarcinomas with morphological and molecular features similar to those of perihilar and extrahepatic cholangiocarcinomas. <i>Histopathology</i> , 2012 , 61, 1106-16	7.3	29

149	MicroRNA 3Q and nucleotide modification patterns and arm selection preference in liver tissues. <i>BMC Systems Biology</i> , 2012 , 6 Suppl 2, S14	3.5	29
148	B3GNT3 expression suppresses cell migration and invasion and predicts favorable outcomes in neuroblastoma. <i>Cancer Science</i> , 2013 , 104, 1600-8	6.9	29
147	Assessing hepatic fibrosis: comparing the intravoxel incoherent motion in MRI with acoustic radiation force impulse imaging in US. <i>European Radiology</i> , 2015 , 25, 3552-9	8	28
146	IMP-3 promotes migration and invasion of melanoma cells by modulating the expression of HMGA2 and predicts poor prognosis in melanoma. <i>Journal of Investigative Dermatology</i> , 2015 , 135, 1065-1073	4.3	28
145	Annexin A10 is an immunohistochemical marker for adenocarcinoma of the upper gastrointestinal tract and pancreatobiliary system. <i>Histopathology</i> , 2013 , 63, 640-8	7.3	28
144	Mutation of the DR5/TRAIL receptor 2 gene is infrequent in hepatocellular carcinoma. <i>Cancer Letters</i> , 2002 , 181, 205-8	9.9	28
143	Congenital interstitial cell of cajal hyperplasia with neuronal intestinal dysplasia. <i>American Journal of Surgical Pathology</i> , 2000 , 24, 1568-72	6.7	28
142	Immune-related adverse reactions in the hepatobiliary system: second-generation check-point inhibitors highlight diverse histological changes. <i>Histopathology</i> , 2020 , 76, 470-480	7.3	28
141	TGF- β secreted by Tregs in lymph nodes promotes breast cancer malignancy via up-regulation of IL-17RB. <i>EMBO Molecular Medicine</i> , 2017 , 9, 1660-1680	12	27
140	Aberrant expression of annexin A10 is closely related to gastric phenotype in serrated pathway to colorectal carcinoma. <i>Modern Pathology</i> , 2015 , 28, 268-78	9.8	27
139	The stem cell E3-ligase Lin-41 promotes liver cancer progression through inhibition of microRNA-mediated gene silencing. <i>Journal of Pathology</i> , 2013 , 229, 486-96	9.4	27
138	Connective tissue growth factor acts as a therapeutic agent and predictor for peritoneal carcinomatosis of colorectal cancer. <i>Clinical Cancer Research</i> , 2011 , 17, 3077-88	12.9	27
137	GALNT2 suppresses malignant phenotypes through IGF-1 receptor and predicts favorable prognosis in neuroblastoma. <i>Oncotarget</i> , 2014 , 5, 12247-59	3.3	27
136	Incorporation of the factor IX Padua mutation into FIX-Triple improves clotting activity in vitro and in vivo. <i>Thrombosis and Haemostasis</i> , 2013 , 110, 244-56	7	26
135	B4GALNT3 expression predicts a favorable prognosis and suppresses cell migration and invasion via Integrin signaling in neuroblastoma. <i>American Journal of Pathology</i> , 2011 , 179, 1394-404	5.8	26
134	Male germ cell-specific RNA binding protein RBMY: a new oncogene explaining male predominance in liver cancer. <i>PLoS ONE</i> , 2011 , 6, e26948	3.7	26
133	Identification of GRP75 as an independent favorable prognostic marker of neuroblastoma by a proteomics analysis. <i>Clinical Cancer Research</i> , 2008 , 14, 6237-45	12.9	26
132	Relationship between ultrasound backscattered statistics and the concentration of fatty droplets in livers: an animal study. <i>PLoS ONE</i> , 2013 , 8, e63543	3.7	26

131	Prevalence and rapid identification of clarithromycin-resistant <i>Helicobacter pylori</i> isolates in children. <i>Pediatric Infectious Disease Journal</i> , 2001 , 20, 662-6	3.4	26
130	RNF43 Is an Early and Specific Mutated Gene in the Serrated Pathway, With Increased Frequency in Traditional Serrated Adenoma and Its Associated Malignancy. <i>American Journal of Surgical Pathology</i> , 2016 , 40, 1352-9	6.7	26
129	Quantification of hepatic steatosis: a comparison of the accuracy among multiple magnetic resonance techniques. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2014 , 29, 807-13	4	25
128	Hepatitis B virus basal core promoter mutation and DNA load correlate with expression of hepatitis B core antigen in patients with chronic hepatitis B. <i>Journal of Infectious Diseases</i> , 2009 , 199, 742-9	7	25
127	Ex vivo imaging and quantification of liver fibrosis using second-harmonic generation microscopy. <i>Journal of Biomedical Optics</i> , 2010 , 15, 036002	3.5	24
126	Prognostic implication of MET overexpression in myxofibrosarcomas: an integrative array comparative genomic hybridization, real-time quantitative PCR, immunoblotting, and immunohistochemical analysis. <i>Modern Pathology</i> , 2010 , 23, 1379-92	9.8	24
125	Histone deacetylase 6 regulates estrogen receptor alpha in uterine leiomyoma. <i>Reproductive Sciences</i> , 2011 , 18, 755-62	3	24
124	Nuclear translocation of beta-catenin protein but absence of beta-catenin and APC mutation in gastrointestinal carcinoid tumor. <i>Annals of Surgical Oncology</i> , 2006 , 13, 1604-9	3.1	24
123	Synergistic effect of radiation and interleukin-6 on hepatitis B virus reactivation in liver through STAT3 signaling pathway. <i>International Journal of Radiation Oncology Biology Physics</i> , 2009 , 75, 1545-52	4	23
122	Quantification of Serum Matrix Metalloproteinase 7 Levels May Assist in the Diagnosis and Predict the Outcome for Patients with Biliary Atresia. <i>Journal of Pediatrics</i> , 2019 , 208, 30-37.e1	3.6	22
121	Expression of hypoxic marker carbonic anhydrase IX predicts poor prognosis in resectable hepatocellular carcinoma. <i>PLoS ONE</i> , 2015 , 10, e0119181	3.7	22
120	Aryl hydrocarbon receptor downregulates MYCN expression and promotes cell differentiation of neuroblastoma. <i>PLoS ONE</i> , 2014 , 9, e88795	3.7	22
119	Long-term follow-up of children with postnatal immunoprophylaxis failure who were infected with hepatitis B virus surface antigen gene mutant. <i>Journal of Infectious Diseases</i> , 2013 , 207, 1047-57	7	22
118	Nuclear overexpression of mitotic regulatory proteins in biliary tract cancer: correlation with clinicopathologic features and patient survival. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2009 , 18, 417-23	4	22
117	S100P expression is a novel prognostic factor in hepatocellular carcinoma and predicts survival in patients with high tumor stage or early recurrent tumors. <i>PLoS ONE</i> , 2013 , 8, e65501	3.7	22
116	SPOCK1 Is a Novel Transforming Growth Factor-Induced Myoepithelial Marker That Enhances Invasion and Correlates with Poor Prognosis in Breast Cancer. <i>PLoS ONE</i> , 2016 , 11, e0162933	3.7	22
115	The Diagnostic Utility of PAX8 for Neuroendocrine Tumors: An Immunohistochemical Reappraisal. <i>Applied Immunohistochemistry and Molecular Morphology</i> , 2016 , 24, 57-63	1.9	22
114	Clinicopathologic Characterization of GREB1-rearranged Uterine Sarcomas With Variable Sex-Cord Differentiation. <i>American Journal of Surgical Pathology</i> , 2019 , 43, 928-942	6.7	22

113	Human pancreatic afferent and efferent nerves: mapping and 3-D illustration of exocrine, endocrine, and adipose innervation. <i>American Journal of Physiology - Renal Physiology</i> , 2019 , 317, G694-G706	5.1	21
112	Human cationic trypsinogen but not serine peptidase inhibitor, Kazal type 1 variants increase the risk of type 1 autoimmune pancreatitis. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2014 , 29, 2038-42	4	21
111	Insulin-like growth factor II mRNA-binding protein 3 expression predicts unfavorable prognosis in patients with neuroblastoma. <i>Cancer Science</i> , 2011 , 102, 2191-8	6.9	21
110	Downregulation of alpha-fetoprotein expression by LHX4: a critical role in hepatocarcinogenesis. <i>Carcinogenesis</i> , 2011 , 32, 1815-23	4.6	21
109	Expression of the caudal-type homeodomain transcription factor CDX2 is related to clinical outcome in biliary tract carcinoma. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2007 , 22, 389-94	4.4	21
108	Proline-directed protein kinase FA is a powerful and independent prognostic predictor for progression and patient survival of hepatocellular carcinoma. <i>Journal of Clinical Oncology</i> , 2006 , 24, 3780-8	2.3	21
107	Diagnostic FDG and FDOPA positron emission tomography scans distinguish the genomic type and treatment outcome of neuroblastoma. <i>Oncotarget</i> , 2016 , 7, 18774-86	3.3	21
106	Tumor size is a major determinant of prognosis of resected stage I hepatocellular carcinoma. <i>Langenbeck's Archives of Surgery</i> , 2015 , 400, 725-34	3.4	20
105	High neuroendocrine component is a factor for poor prognosis in gastrointestinal high-grade malignant mixed adenoneuroendocrine neoplasms. <i>Journal of the Chinese Medical Association</i> , 2015 , 78, 454-9	2.8	20
104	RNF43 mutation frequently occurs with GNAS mutation and mucin hypersecretion in intraductal papillary neoplasms of the bile duct. <i>Histopathology</i> , 2017 , 70, 756-765	7.3	20
103	Insulin-like growth factor II mRNA-binding protein 3 expression promotes tumor formation and invasion and predicts poor prognosis in oral squamous cell carcinoma. <i>Journal of Oral Pathology and Medicine</i> , 2011 , 40, 699-705	3.3	20
102	Liver steatosis classification using high-frequency ultrasound. <i>Ultrasound in Medicine and Biology</i> , 2005 , 31, 599-605	3.5	20
101	Nuclear GRP75 binds retinoic acid receptors to promote neuronal differentiation of neuroblastoma. <i>PLoS ONE</i> , 2011 , 6, e26236	3.7	20
100	Mutational analysis of MED12 exon 2 in a spectrum of fibroepithelial tumours of the breast: implications for pathogenesis and histogenesis. <i>Histopathology</i> , 2016 , 68, 433-41	7.3	20
99	Connective tissue growth factor inhibits gastric cancer peritoneal metastasis by blocking integrin $\beta 1$ -dependent adhesion. <i>Gastric Cancer</i> , 2015 , 18, 504-15	7.6	19
98	Biological significance of GATA3, cytokeratin 20, cytokeratin 5/6 and p53 expression in muscle-invasive bladder cancer. <i>PLoS ONE</i> , 2019 , 14, e0221785	3.7	18
97	Targeted next-generation sequencing identifies distinct clinicopathologic and molecular entities of intraductal papillary neoplasms of the bile duct. <i>Modern Pathology</i> , 2019 , 32, 1637-1645	9.8	18
96	PRRX2 as a novel TGF- β -induced factor enhances invasion and migration in mammary epithelial cell and correlates with poor prognosis in breast cancer. <i>Molecular Carcinogenesis</i> , 2016 , 55, 2247-2259	5	18

95	Expression of insulin-like growth factor II mRNA-binding protein 3 predicts early recurrence and poor prognosis in intrahepatic cholangiocarcinoma. <i>International Journal of Surgery</i> , 2013 , 11, 85-91	7.5	18
94	NKX6-1 Is a Novel Immunohistochemical Marker for Pancreatic and Duodenal Neuroendocrine Tumors. <i>American Journal of Surgical Pathology</i> , 2015 , 39, 850-7	6.7	18
93	Phenytoin-induced pseudolymphoma: reevaluation using modern molecular biology techniques. <i>Epilepsia</i> , 1996 , 37, 104-7	6.4	18
92	Reliability of a single-region sample to evaluate tumor immune microenvironment in hepatocellular carcinoma. <i>Journal of Hepatology</i> , 2020 , 72, 489-497	13.4	18
91	Hepatic ADC map as an adjunct to conventional abdominal MRI to evaluate hepatic fibrotic and clinical cirrhotic severity in biliary atresia patients. <i>European Radiology</i> , 2015 , 25, 2992-3002	8	17
90	Decreased expressions of hepsin in human hepatocellular carcinomas. <i>Liver International</i> , 2006 , 26, 774-809	8.9	17
89	Concordant body stalk anomaly in monozygotic twinning--early embryo cleavage disorder. <i>Prenatal Diagnosis</i> , 1996 , 16, 467-70	3.2	16
88	Input of microenvironmental regulation on colorectal cancer: role of the CCN family. <i>World Journal of Gastroenterology</i> , 2014 , 20, 6826-31	5.6	16
87	Carbonic anhydrase VI: a novel marker for salivary serous acinar differentiation and its application to discriminate acinic cell carcinoma from mammary analogue secretory carcinoma of the salivary gland. <i>Histopathology</i> , 2016 , 68, 641-7	7.3	16
86	Clinical course of de novo hepatitis B infection after pediatric liver transplantation. <i>Liver Transplantation</i> , 2010 , 16, 215-21	4.5	15
85	Lamivudine treatment in maternally transmitted chronic hepatitis B virus infection patients. <i>Pediatrics International</i> , 2005 , 47, 372-7	1.2	15
84	IL-18-induced interaction between IMP3 and HuR contributes to COX-2 mRNA stabilization in acute myeloid leukemia. <i>Journal of Leukocyte Biology</i> , 2016 , 99, 131-41	6.5	14
83	Klotho-beta and fibroblast growth factor 19 expression correlates with early recurrence of resectable hepatocellular carcinoma. <i>Liver International</i> , 2019 , 39, 1682-1691	7.9	14
82	Biological significance of TERT promoter mutation in papillary urothelial neoplasm of low malignant potential. <i>Histopathology</i> , 2018 , 72, 795-803	7.3	14
81	Prognostic significance of TWEAK expression in colorectal cancer and effect of its inhibition on invasion. <i>Annals of Surgical Oncology</i> , 2012 , 19 Suppl 3, S385-94	3.1	14
80	Development of autoantibodies after pediatric liver transplantation. <i>Pediatric Transplantation</i> , 2013 , 17, 144-8	1.8	14
79	Underexpression of hepatocyte nuclear factor-1 in chromophobe renal cell carcinoma. <i>Histopathology</i> , 2013 , 62, 589-94	7.3	14
78	Phosphorylation of NuSAP by Cdk1 regulates its interaction with microtubules in mitosis. <i>Cell Cycle</i> , 2011 , 10, 4083-9	4.7	14

77	Expression of LEF1 is an independent prognostic factor for patients with oral squamous cell carcinoma. <i>Journal of the Formosan Medical Association</i> , 2014 , 113, 934-9	3.2	13
76	Clinical significance and evolution of hepatic HBsAg expression in HBeAg-positive patients receiving interferon therapy. <i>Journal of Gastroenterology</i> , 2014 , 49, 356-62	6.9	13
75	Calreticulin mediates nerve growth factor-induced neuronal differentiation. <i>Journal of Molecular Neuroscience</i> , 2012 , 47, 571-81	3.3	13
74	Upregulation of endocan by Epstein-Barr virus latent membrane protein 1 and its clinical significance in nasopharyngeal carcinoma. <i>PLoS ONE</i> , 2013 , 8, e82254	3.7	13
73	Survival after pancreaticoduodenectomy for ampullary cancer is not affected by age. <i>World Journal of Surgery</i> , 2010 , 34, 2945-52	3.3	13
72	The Role of Organelle Stresses in Diabetes Mellitus and Obesity: Implication for Treatment. <i>Analytical Cellular Pathology</i> , 2015 , 2015, 972891	3.4	12
71	HSD3B1 is a specific trophoblast-associated marker not expressed in a wide spectrum of tumors. <i>International Journal of Gynecological Cancer</i> , 2013 , 23, 343-7	3.5	12
70	Intravasation-related metastatic factors in colorectal cancer. <i>Tumor Biology</i> , 2004 , 25, 48-55	2.9	12
69	A multidisciplinary team care approach improves outcomes in high-risk pediatric neuroblastoma patients. <i>Oncotarget</i> , 2017 , 8, 4360-4372	3.3	12
68	Pancreatic cancer screening in different risk individuals with family history of pancreatic cancer-a prospective cohort study in Taiwan. <i>American Journal of Cancer Research</i> , 2017 , 7, 357-369	4.4	12
67	Pancreatic cancer-derived small extracellular vesical Ezrin regulates macrophage polarization and promotes metastasis. <i>American Journal of Cancer Research</i> , 2020 , 10, 12-37	4.4	12
66	Exploring Markers of Exhausted CD8 T Cells to Predict Response to Immune Checkpoint Inhibitor Therapy for Hepatocellular Carcinoma. <i>Liver Cancer</i> , 2021 , 10, 346-359	9.1	12
65	Lymphatic vessel remodeling and invasion in pancreatic cancer progression. <i>EBioMedicine</i> , 2019 , 47, 98-118	11.8	11
64	Prognostic Significance of CDCP1 Expression in Colorectal Cancer and Effect of Its Inhibition on Invasion and Migration. <i>Annals of Surgical Oncology</i> , 2015 , 22, 4335-43	3.1	11
63	The prognostic roles of and correlation between and protein expression in neuroblastoma. <i>Journal of Clinical Pathology</i> , 2020 , 73, 154-161	3.9	11
62	Microinsertions in PRKACA cause activation of the protein kinase A pathway in cardiac myxoma. <i>Journal of Pathology</i> , 2017 , 242, 134-139	9.4	10
61	Changes in backscattered ultrasonic envelope statistics as a function of thrombus age: an in vitro study. <i>Ultrasound in Medicine and Biology</i> , 2015 , 41, 498-508	3.5	10
60	Expression of bile duct transcription factor HNF1 α predicts early tumor recurrence and is a stage-independent prognostic factor in hepatocellular carcinoma. <i>Journal of Gastrointestinal Surgery</i> , 2014 , 18, 1784-94	3.3	10

59	Potential synergistic anti-tumor activity between lenalidomide and sorafenib in hepatocellular carcinoma. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2014 , 29, 2021-31	4	10
58	Validation of Indications for Surgery of European Evidence-Based Guidelines for Patients with Pancreatic Intraductal Papillary Mucinous Neoplasms. <i>Journal of Gastrointestinal Surgery</i> , 2020 , 24, 2536-2543	3.3	10
57	Roles of Epstein-Barr virus viral load monitoring in the prediction of posttransplant lymphoproliferative disorder in pediatric liver transplantation. <i>Journal of the Formosan Medical Association</i> , 2019 , 118, 1362-1368	3.2	10
56	Interplay between desmoglein2 and hypoxia controls metastasis in breast cancer. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021 , 118,	11.5	10
55	Persistent elevation of hepatocyte growth factor activator inhibitors in cholangiopathies affects liver fibrosis and differentiation. <i>Hepatology</i> , 2012 , 55, 161-72	11.2	9
54	Elevated preoperative serum CA19-9 levels in patients with hepatocellular carcinoma is associated with poor prognosis after resection. <i>Scientific World Journal, The</i> , 2013 , 2013, 380797	2.2	9
53	Paradoxical overexpression of MBNL2 in hepatocellular carcinoma inhibits tumor growth and invasion. <i>Oncotarget</i> , 2016 , 7, 65589-65601	3.3	9
52	MLN4924, a Novel NEDD8-activating enzyme inhibitor, exhibits antitumor activity and enhances cisplatin-induced cytotoxicity in human cervical carcinoma: in vitro and in vivo study. <i>American Journal of Cancer Research</i> , 2015 , 5, 3350-62	4.4	9
51	MED12 exon 2 mutation as a highly sensitive and specific marker in distinguishing phyllodes tumours from other spindle neoplasms of the breast. <i>Apmis</i> , 2016 , 124, 356-64	3.4	9
50	Intestinal Behçet disease presenting as neonatal onset chronic diarrhea in an 11-month-old male baby. <i>European Journal of Pediatrics</i> , 2005 , 164, 523-5	4.1	8
49	Increased Trimethylation of histone H3K36 associates with biliary differentiation and predicts poor prognosis in resectable hepatocellular carcinoma. <i>PLoS ONE</i> , 2018 , 13, e0206261	3.7	8
48	Aberrant expressions of annexin A10 short isoform, osteopontin and alpha-fetoprotein at chromosome 4q cooperatively contribute to progression and poor prognosis of hepatocellular carcinoma. <i>International Journal of Oncology</i> , 2005 , 26, 1053-61	1	8
47	Complementary roles of E-cadherin and glutamine synthetase immunostaining in diagnosis of chemotherapy-treated and untreated hepatoblastoma. <i>Journal of the Formosan Medical Association</i> , 2017 , 116, 549-553	3.2	7
46	Mist1: a novel nuclear marker for acinic cell carcinoma of the salivary gland. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2019 , 475, 617-624	5.1	7
45	Correlation of molecular and morphological features of appendiceal epithelial neoplasms. <i>Histopathology</i> , 2019 , 75, 468-477	7.3	7
44	Plectin Mutations in Progressive Familial Intrahepatic Cholestasis. <i>Hepatology</i> , 2019 , 70, 2221-2224	11.2	7
43	Overweight increases the risk of malignancy in patients with pancreatic mucinous cystic neoplasms. <i>Medicine (United States)</i> , 2015 , 94, e797	1.8	7
42	Claudin-18 as a Marker for Identifying the Stomach and Pancreatobiliary Tract as the Primary Sites of Metastatic Adenocarcinoma. <i>American Journal of Surgical Pathology</i> , 2020 , 44, 1643-1648	6.7	7

41	Cystic fibrosis transmembrane conductance regulator gene variants are associated with autoimmune pancreatitis and slow response to steroid treatment. <i>Journal of Cystic Fibrosis</i> , 2015 , 14, 661-7	4.1	6
40	Emerging importance of autoimmune hepatitis in children in Taiwan, an endemic area for viral hepatitis. <i>Pediatrics and Neonatology</i> , 2009 , 50, 65-9	1.8	6
39	X-linked liver glycogenosis in a Taiwanese family: transmission from undiagnosed males. <i>Pediatrics and Neonatology</i> , 2009 , 50, 230-3	1.8	6
38	Acute Q fever presenting as fever of unknown origin with rapidly progressive hepatic failure in a patient with alcoholism. <i>Journal of the Formosan Medical Association</i> , 2008 , 107, 896-901	3.2	6
37	Bone marrow transplantation results in donor-derived hepatocytes in an animal model of inherited cholestatic liver disease. <i>Journal of Biomedical Science</i> , 2008 , 15, 615-22	13.3	6
36	p53 gene and Wnt signaling in benign neoplasms: E-catenin mutations in hepatic adenoma but not in focal nodular hyperplasia. <i>Hepatology</i> , 2002 , 36, 927-935	11.2	6
35	MYCN RNA levels determined by quantitative in situ hybridization is better than MYCN gene dosages in predicting the prognosis of neuroblastoma patients. <i>Modern Pathology</i> , 2020 , 33, 531-540	9.8	6
34	Differential integration rates of hepatitis B virus DNA in the liver of children with chronic hepatitis B virus infection and hepatocellular carcinoma. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2005 , 20, 1206-14	4	5
33	Glypican-3 induces oncogenicity by preventing IGF-1R degradation, a process that can be blocked by Grb10. <i>Oncotarget</i> , 2017 , 8, 80429-80442	3.3	5
32	Treatment outcomes regarding the addition of targeted agents in the therapeutic portfolio for stage II-III rectal cancer undergoing neoadjuvant chemoradiation. <i>Oncotarget</i> , 2017 , 8, 101832-101846	3.3	5
31	Hepatocellular adenoma in Taiwan: Distinct ensemble of male predominance, overweight/obesity, and inflammatory subtype. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2020 , 35, 680-688	4	5
30	Characterization of initial key steps of IL-17 receptor B oncogenic signaling for targeted therapy of pancreatic cancer. <i>Science Translational Medicine</i> , 2021 , 13,	17.5	5
29	Traditional Serrated Pathway-associated Colorectal Carcinoma: Morphologic Reappraisal of Serrated Morphology, Tumor Budding, and Identification of Frequent PTEN Alterations. <i>American Journal of Surgical Pathology</i> , 2019 , 43, 1042-1051	6.7	5
28	Autophagy restricts mitochondrial DNA damage-induced release of ENDOG (endonuclease G) to regulate genome stability. <i>Autophagy</i> , 2021 , 17, 3444-3460	10.2	5
27	Complement C1q mediates the expansion of periportal hepatic progenitor cells in senescence-associated inflammatory liver. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020 , 117, 6717-6725	11.5	4
26	Deletion of cadherin-17 enhances intestinal permeability and susceptibility to intestinal tumour formation. <i>Journal of Pathology</i> , 2018 , 246, 289-299	9.4	4
25	The overexpression of MYST4 in human solid tumors is associated with increased aggressiveness and decreased overall survival. <i>International Journal of Clinical and Experimental Pathology</i> , 2019 , 12, 431-442	1.4	4
24	Frequent BRAF mutation in early-onset colorectal cancer in Taiwan: association with distinct clinicopathological and molecular features and poor clinical outcome. <i>Journal of Clinical Pathology</i> , 2016 , 69, 319-25	3.9	3

23	Overexpression of Her-2/neu in Epithelial Ovarian Carcinoma Induces Vascular Endothelial Growth Factor C by Activating NF- κ B: Implications for Malignant Ascites Formation and Tumor Lymphangiogenesis. <i>Journal of Biomedical Science</i> , 2004 , 11, 249-259	13.3	3
22	Biliary hamartomas with delayed 99mTc-diisopropyl iminodiacetic acid clearance. <i>Journal of Gastroenterology</i> , 2005 , 40, 540-4	6.9	3
21	Cellular 5-hydroxymethylcytosine content determines tumorigenic potential and prognosis of pancreatic ductal adenocarcinoma. <i>American Journal of Cancer Research</i> , 2018 , 8, 2548-2563	4.4	3
20	Characterization of metastatic tumor antigen 1 and its interaction with hepatitis B virus X protein in NF-B signaling and tumor progression in a woodchuck hepatocellular carcinoma model. <i>Oncotarget</i> , 2016 , 7, 47173-47185	3.3	3
19	Radical endoscopic polypectomy combined with double-balloon enteroscopy and colonoscopy for Peutz-Jeghers syndrome. <i>Journal of Pediatric Gastroenterology and Nutrition</i> , 2010 , 51, 370-2	2.8	3
18	Lymphoepithelioma-like Intrahepatic Cholangiocarcinoma Is a Distinct Entity With Frequent pTERT/TP53 Mutations and Comprises 2 Subgroups Based on Epstein-Barr Virus Infection. <i>American Journal of Surgical Pathology</i> , 2021 , 45, 1409-1418	6.7	3
17	Clostridium difficile infection in a patient with Crohn disease. <i>Journal of the Formosan Medical Association</i> , 2012 , 111, 347-9	3.2	2
16	Aberrant expressions of annexin A10 short isoform, osteopontin and Fetoprotein at chromosome 4q cooperatively contribute to progression and poor prognosis of hepatocellular carcinoma 2005 , 26, 1053		2
15	YoungQ modulus measurements of human liver and correlation with pathological findings		2
14	Clinical manifestations and outcomes of patients with scirrhous hepatocellular carcinoma. <i>Hepatology International</i> , 2021 , 15, 472-481	8.8	2
13	The occurrence of primary hepatic adenoma in deceased donor renal transplant recipient. <i>International Braz J Urol: Official Journal of the Brazilian Society of Urology</i> , 2014 , 40, 118-22	2	1
12	Unique Phenotypes With Corresponding Pathology in Late-Onset Hereditary Transthyretin Amyloidosis of A97S vs. V30M.. <i>Frontiers in Aging Neuroscience</i> , 2021 , 13, 786322	5.3	1
11	S100P as a marker for poor survival and advanced stage in gallbladder carcinoma. <i>Annals of Diagnostic Pathology</i> , 2021 , 52, 151736	2.2	1
10	CONCORDANT BODY STALK ANOMALY IN MONOZYGOTIC TWINNINGEARLY EMBRYO CLEAVAGE DISORDER 1996 , 16, 467		1
9	Corneal transparency and scleral opacity arises from the nanoarchitecture of the constituent collagen fibrils.. <i>Biomedical Optics Express</i> , 2022 , 13, 1485-1496	3.5	0
8	Rapid Trio Exome Sequencing for Autosomal Recessive Renal Tubular Dysgenesis in Recurrent Oligohydramnios. <i>Frontiers in Genetics</i> , 2021 , 12, 606970	4.5	0
7	Local islet remodelling associated with duct lesion-islet complex in adult human pancreas. <i>Diabetologia</i> , 2021 , 64, 2266-2278	10.3	0
6	Association of annexin A10 expression with poor prognosis of intrahepatic cholangiocarcinoma.. <i>BMC Cancer</i> , 2022 , 22, 219	4.8	0

- 5 2 Role of immunohistochemical expression of p53 in colorectal carcinoma. *Handbook of Immunohistochemistry and in Situ Hybridization of Human Carcinomas*, **2002**, 139-147
- 4 The study of clinicopathologic correlation with the expression level of 5-hydroxymethylcytosine in GIST.. *Journal of Clinical Oncology*, **2018**, 36, e23520-e23520 2.2
- 3 Retraction: Knockdown of Contactin-1 Expression Suppresses Invasion and Metastasis of Lung Adenocarcinoma. *Cancer Research*, **2019**, 79, 3793 10.1
- 2 Immunoglobulin G4-related sclerosing cholangitis in a 3 years of age boy. *Advances in Digestive Medicine*, **2021**, 8, 59-63 0.1
- 1 Differential effects of glucose and N-acetylglucosamine on genome instability.. *American Journal of Cancer Research*, **2022**, 12, 1556-1576 4.4