

Yung-Ming Jeng

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/4252767/yung-ming-jeng-publications-by-year.pdf>

Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

256 papers	11,285 citations	58 h-index	93 g-index
267 ext. papers	13,024 ext. citations	6.8 avg, IF	6 L-index

#	Paper	IF	Citations
256	Corneal transparency and scleral opacity arises from the nanoarchitecture of the constituent collagen fibrils.. <i>Biomedical Optics Express</i> , 2022 , 13, 1485-1496	3.5	0
255	Association of annexin A10 expression with poor prognosis of intrahepatic cholangiocarcinoma.. <i>BMC Cancer</i> , 2022 , 22, 219	4.8	0
254	Differential effects of glucose and N-acetylglucosamine on genome instability.. <i>American Journal of Cancer Research</i> , 2022 , 12, 1556-1576	4.4	
253	Unique Phenotypes With Corresponding Pathology in Late-Onset Hereditary Transthyretin Amyloidosis of A97S vs. V30M.. <i>Frontiers in Aging Neuroscience</i> , 2021 , 13, 786322	5.3	1
252	Characterization of initial key steps of IL-17 receptor B oncogenic signaling for targeted therapy of pancreatic cancer. <i>Science Translational Medicine</i> , 2021 , 13,	17.5	5
251	Lymphoepithelioma-like Intrahepatic Cholangiocarcinoma Is a Distinct Entity With Frequent pTERT/TP53 Mutations and Comprises 2 Subgroups Based on Epstein-Barr Virus Infection. <i>American Journal of Surgical Pathology</i> , 2021 , 45, 1409-1418	6.7	3
250	Exploring Markers of Exhausted CD8 T Cells to Predict Response to Immune Checkpoint Inhibitor Therapy for Hepatocellular Carcinoma. <i>Liver Cancer</i> , 2021 , 10, 346-359	9.1	12
249	S100P as a marker for poor survival and advanced stage in gallbladder carcinoma. <i>Annals of Diagnostic Pathology</i> , 2021 , 52, 151736	2.2	1
248	Rapid Trio Exome Sequencing for Autosomal Recessive Renal Tubular Dysgenesis in Recurrent Oligohydramnios. <i>Frontiers in Genetics</i> , 2021 , 12, 606970	4.5	0
247	Local islet remodelling associated with duct lesion-islet complex in adult human pancreas. <i>Diabetologia</i> , 2021 , 64, 2266-2278	10.3	0
246	Immunoglobulin G4-related sclerosing cholangitis in a 3 years of age boy. <i>Advances in Digestive Medicine</i> , 2021 , 8, 59-63	0.1	
245	Autophagy restricts mitochondrial DNA damage-induced release of ENDOG (endonuclease G) to regulate genome stability. <i>Autophagy</i> , 2021 , 17, 3444-3460	10.2	5
244	Interplay between desmoglein2 and hypoxia controls metastasis in breast cancer. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021 , 118,	11.5	10
243	Clinical manifestations and outcomes of patients with scirrhous hepatocellular carcinoma. <i>Hepatology International</i> , 2021 , 15, 472-481	8.8	2
242	Complement C1q mediates the expansion of periportal hepatic progenitor cells in senescence-associated inflammatory liver. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020 , 117, 6717-6725	11.5	4
241	B cells are associated with survival and immunotherapy response in sarcoma. <i>Nature</i> , 2020 , 577, 556-560	50.4	538
240	Pancreatic cancer-derived small extracellular vesical Ezrin regulates macrophage polarization and promotes metastasis. <i>American Journal of Cancer Research</i> , 2020 , 10, 12-37	4.4	12

239	Claudin-18 as a Marker for Identifying the Stomach and Pancreatobiliary Tract as the Primary Sites of Metastatic Adenocarcinoma. <i>American Journal of Surgical Pathology</i> , 2020 , 44, 1643-1648	6.7	7
238	Hepatocellular adenoma in Taiwan: Distinct ensemble of male predominance, overweight/obesity, and inflammatory subtype. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2020 , 35, 680-688	4	5
237	MYCN RNA levels determined by quantitative in situ hybridization is better than MYCN gene dosages in predicting the prognosis of neuroblastoma patients. <i>Modern Pathology</i> , 2020 , 33, 531-540	9.8	6
236	Immune-related adverse reactions in the hepatobiliary system: second-generation check-point inhibitors highlight diverse histological changes. <i>Histopathology</i> , 2020 , 76, 470-480	7.3	28
235	Reliability of a single-region sample to evaluate tumor immune microenvironment in hepatocellular carcinoma. <i>Journal of Hepatology</i> , 2020 , 72, 489-497	13.4	18
234	Validation of Indications for Surgery of European Evidence-Based Guidelines for Patients with Pancreatic Intraductal Papillary Mucinous Neoplasms. <i>Journal of Gastrointestinal Surgery</i> , 2020 , 24, 2536-2543	3.3	10
233	The prognostic roles of and correlation between and protein expression in neuroblastoma. <i>Journal of Clinical Pathology</i> , 2020 , 73, 154-161	3.9	11
232	Biological significance of GATA3, cytokeratin 20, cytokeratin 5/6 and p53 expression in muscle-invasive bladder cancer. <i>PLoS ONE</i> , 2019 , 14, e0221785	3.7	18
231	Lymphatic vessel remodeling and invasion in pancreatic cancer progression. <i>EBioMedicine</i> , 2019 , 47, 98-113	11.8	11
230	Human pancreatic afferent and efferent nerves: mapping and 3-D illustration of exocrine, endocrine, and adipose innervation. <i>American Journal of Physiology - Renal Physiology</i> , 2019 , 317, G694-G706	5.1	21
229	Klotho-beta and fibroblast growth factor 19 expression correlates with early recurrence of resectable hepatocellular carcinoma. <i>Liver International</i> , 2019 , 39, 1682-1691	7.9	14
228	Targeted next-generation sequencing identifies distinct clinicopathologic and molecular entities of intraductal papillary neoplasms of the bile duct. <i>Modern Pathology</i> , 2019 , 32, 1637-1645	9.8	18
227	Mist1: a novel nuclear marker for acinic cell carcinoma of the salivary gland. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2019 , 475, 617-624	5.1	7
226	Correlation of molecular and morphological features of appendiceal epithelial neoplasms. <i>Histopathology</i> , 2019 , 75, 468-477	7.3	7
225	High Glucose Triggers Nucleotide Imbalance through O-GlcNAcylation of Key Enzymes and Induces KRAS Mutation in Pancreatic Cells. <i>Cell Metabolism</i> , 2019 , 29, 1334-1349.e10	24.6	40
224	Quantification of Serum Matrix Metalloproteinase 7 Levels May Assist in the Diagnosis and Predict the Outcome for Patients with Biliary Atresia. <i>Journal of Pediatrics</i> , 2019 , 208, 30-37.e1	3.6	22
223	Clinical Manifestations and Outcomes of Patients with Sarcomatoid Hepatocellular Carcinoma. <i>Hepatology</i> , 2019 , 69, 209-221	11.2	38
222	Plectin Mutations in Progressive Familial Intrahepatic Cholestasis. <i>Hepatology</i> , 2019 , 70, 2221-2224	11.2	7

221	The overexpression of MYST4 in human solid tumors is associated with increased aggressiveness and decreased overall survival. <i>International Journal of Clinical and Experimental Pathology</i> , 2019 , 12, 431-442	1.4	4
220	Retraction: Knockdown of Contactin-1 Expression Suppresses Invasion and Metastasis of Lung Adenocarcinoma. <i>Cancer Research</i> , 2019 , 79, 3793	10.1	
219	Clinicopathologic Characterization of GREB1-rearranged Uterine Sarcomas With Variable Sex-Cord Differentiation. <i>American Journal of Surgical Pathology</i> , 2019 , 43, 928-942	6.7	22
218	Traditional Serrated Pathway-associated Colorectal Carcinoma: Morphologic Reappraisal of Serrated Morphology, Tumor Budding, and Identification of Frequent PTEN Alterations. <i>American Journal of Surgical Pathology</i> , 2019 , 43, 1042-1051	6.7	5
217	Roles of Epstein-Barr virus viral load monitoring in the prediction of posttransplant lymphoproliferative disorder in pediatric liver transplantation. <i>Journal of the Formosan Medical Association</i> , 2019 , 118, 1362-1368	3.2	10
216	Transient elastography is useful in diagnosing biliary atresia and predicting prognosis after hepatoportoenterostomy. <i>Hepatology</i> , 2018 , 68, 616-624	11.2	45
215	Biological significance of TERT promoter mutation in papillary urothelial neoplasm of low malignant potential. <i>Histopathology</i> , 2018 , 72, 795-803	7.3	14
214	Deletion of cadherin-17 enhances intestinal permeability and susceptibility to intestinal tumour formation. <i>Journal of Pathology</i> , 2018 , 246, 289-299	9.4	4
213	Cellular 5-hydroxymethylcytosine content determines tumorigenic potential and prognosis of pancreatic ductal adenocarcinoma. <i>American Journal of Cancer Research</i> , 2018 , 8, 2548-2563	4.4	3
212	The study of clinicopathologic correlation with the expression level of 5-hydroxymethylcytosine in GIST.. <i>Journal of Clinical Oncology</i> , 2018 , 36, e23520-e23520	2.2	
211	Increased Trimethylation of histone H3K36 associates with biliary differentiation and predicts poor prognosis in resectable hepatocellular carcinoma. <i>PLoS ONE</i> , 2018 , 13, e0206261	3.7	8
210	Complementary roles of E-catenin and glutamine synthetase immunostaining in diagnosis of chemotherapy-treated and untreated hepatoblastoma. <i>Journal of the Formosan Medical Association</i> , 2017 , 116, 549-553	3.2	7
209	Adipocytes promote malignant growth of breast tumours with monocarboxylate transporter 2 expression via E-hydroxybutyrate. <i>Nature Communications</i> , 2017 , 8, 14706	17.4	50
208	Microinsertions in PRKACA cause activation of the protein kinase A pathway in cardiac myxoma. <i>Journal of Pathology</i> , 2017 , 242, 134-139	9.4	10
207	CTP synthase forms the cytoophidium in human hepatocellular carcinoma. <i>Experimental Cell Research</i> , 2017 , 361, 292-299	4.2	32
206	TGF- β secreted by Tregs in lymph nodes promotes breast cancer malignancy via up-regulation of IL-17RB. <i>EMBO Molecular Medicine</i> , 2017 , 9, 1660-1680	12	27
205	RNF43 mutation frequently occurs with GNAS mutation and mucin hypersecretion in intraductal papillary neoplasms of the bile duct. <i>Histopathology</i> , 2017 , 70, 756-765	7.3	20
204	A multidisciplinary team care approach improves outcomes in high-risk pediatric neuroblastoma patients. <i>Oncotarget</i> , 2017 , 8, 4360-4372	3.3	12

203	Glypican-3 induces oncogenicity by preventing IGF-1R degradation, a process that can be blocked by Grb10. <i>Oncotarget</i> , 2017 , 8, 80429-80442	3.3	5
202	Treatment outcomes regarding the addition of targeted agents in the therapeutic portfolio for stage II-III rectal cancer undergoing neoadjuvant chemoradiation. <i>Oncotarget</i> , 2017 , 8, 101832-101846	3.3	5
201	Pancreatic cancer screening in different risk individuals with family history of pancreatic cancer-a prospective cohort study in Taiwan. <i>American Journal of Cancer Research</i> , 2017 , 7, 357-369	4.4	12
200	IL-18-induced interaction between IMP3 and HuR contributes to COX-2 mRNA stabilization in acute myeloid leukemia. <i>Journal of Leukocyte Biology</i> , 2016 , 99, 131-41	6.5	14
199	Glutaminase 2 stabilizes Dicer to repress Snail and metastasis in hepatocellular carcinoma cells. <i>Cancer Letters</i> , 2016 , 383, 282-294	9.9	37
198	PRRX2 as a novel TGF- β -induced factor enhances invasion and migration in mammary epithelial cell and correlates with poor prognosis in breast cancer. <i>Molecular Carcinogenesis</i> , 2016 , 55, 2247-2259	5	18
197	Clinical Significance of Circulating Tumor Microemboli as a Prognostic Marker in Patients with Pancreatic Ductal Adenocarcinoma. <i>Clinical Chemistry</i> , 2016 , 62, 505-13	5.5	69
196	Frequent BRAF mutation in early-onset colorectal cancer in Taiwan: association with distinct clinicopathological and molecular features and poor clinical outcome. <i>Journal of Clinical Pathology</i> , 2016 , 69, 319-25	3.9	3
195	Paradoxical overexpression of MBNL2 in hepatocellular carcinoma inhibits tumor growth and invasion. <i>Oncotarget</i> , 2016 , 7, 65589-65601	3.3	9
194	Diagnostic FDG and FDOPA positron emission tomography scans distinguish the genomic type and treatment outcome of neuroblastoma. <i>Oncotarget</i> , 2016 , 7, 18774-86	3.3	21
193	Characterization of metastatic tumor antigen 1 and its interaction with hepatitis B virus X protein in NF- κ B signaling and tumor progression in a woodchuck hepatocellular carcinoma model. <i>Oncotarget</i> , 2016 , 7, 47173-47185	3.3	3
192	SPOCK1 Is a Novel Transforming Growth Factor- β -Induced Myoepithelial Marker That Enhances Invasion and Correlates with Poor Prognosis in Breast Cancer. <i>PLoS ONE</i> , 2016 , 11, e0162933	3.7	22
191	MED12 exon 2 mutation as a highly sensitive and specific marker in distinguishing phyllodes tumours from other spindle neoplasms of the breast. <i>Apmis</i> , 2016 , 124, 356-64	3.4	9
190	Mutational analysis of MED12 exon 2 in a spectrum of fibroepithelial tumours of the breast: implications for pathogenesis and histogenesis. <i>Histopathology</i> , 2016 , 68, 433-41	7.3	20
189	Carbonic anhydrase VI: a novel marker for salivary serous acinar differentiation and its application to discriminate acinic cell carcinoma from mammary analogue secretory carcinoma of the salivary gland. <i>Histopathology</i> , 2016 , 68, 641-7	7.3	16
188	The Diagnostic Utility of PAX8 for Neuroendocrine Tumors: An Immunohistochemical Reappraisal. <i>Applied Immunohistochemistry and Molecular Morphology</i> , 2016 , 24, 57-63	1.9	22
187	RNF43 Is an Early and Specific Mutated Gene in the Serrated Pathway, With Increased Frequency in Traditional Serrated Adenoma and Its Associated Malignancy. <i>American Journal of Surgical Pathology</i> , 2016 , 40, 1352-9	6.7	26
186	Targeting IL-17B-IL-17RB signaling with an anti-IL-17RB antibody blocks pancreatic cancer metastasis by silencing multiple chemokines. <i>Journal of Experimental Medicine</i> , 2015 , 212, 333-49	16.6	89

185	Sorafenib and its derivative SC-1 exhibit antifibrotic effects through signal transducer and activator of transcription 3 inhibition. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2015 , 112, 7243-8	11.5	49
184	Alternative lengthening of telomeres phenotype in malignant vascular tumors is highly associated with loss of ATRX expression and is frequently observed in hepatic angiosarcomas. <i>Human Pathology</i> , 2015 , 46, 1360-6	3.7	37
183	Cystic fibrosis transmembrane conductance regulator gene variants are associated with autoimmune pancreatitis and slow response to steroid treatment. <i>Journal of Cystic Fibrosis</i> , 2015 , 14, 661-7	4.1	6
182	Frequent alterations of HER2 through mutation, amplification, or overexpression in pleomorphic lobular carcinoma of the breast. <i>Breast Cancer Research and Treatment</i> , 2015 , 150, 447-55	4.4	38
181	Prognostic Significance of CDCP1 Expression in Colorectal Cancer and Effect of Its Inhibition on Invasion and Migration. <i>Annals of Surgical Oncology</i> , 2015 , 22, 4335-43	3.1	11
180	Hepatic ADC map as an adjunct to conventional abdominal MRI to evaluate hepatic fibrotic and clinical cirrhotic severity in biliary atresia patients. <i>European Radiology</i> , 2015 , 25, 2992-3002	8	17
179	Tumor size is a major determinant of prognosis of resected stage I hepatocellular carcinoma. <i>Langenbeck's Archives of Surgery</i> , 2015 , 400, 725-34	3.4	20
178	Comprehensive screening of alternative lengthening of telomeres phenotype and loss of ATRX expression in sarcomas. <i>Modern Pathology</i> , 2015 , 28, 1545-54	9.8	45
177	Aberrant expression of annexin A10 is closely related to gastric phenotype in serrated pathway to colorectal carcinoma. <i>Modern Pathology</i> , 2015 , 28, 268-78	9.8	27
176	High neuroendocrine component is a factor for poor prognosis in gastrointestinal high-grade malignant mixed adenoneuroendocrine neoplasms. <i>Journal of the Chinese Medical Association</i> , 2015 , 78, 454-9	2.8	20
175	Changes in backscattered ultrasonic envelope statistics as a function of thrombus age: an in vitro study. <i>Ultrasound in Medicine and Biology</i> , 2015 , 41, 498-508	3.5	10
174	Identification of a novel FN1-FGFR1 genetic fusion as a frequent event in phosphaturic mesenchymal tumour. <i>Journal of Pathology</i> , 2015 , 235, 539-45	9.4	90
173	Overweight increases the risk of malignancy in patients with pancreatic mucinous cystic neoplasms. <i>Medicine (United States)</i> , 2015 , 94, e797	1.8	7
172	NKX6-1 Is a Novel Immunohistochemical Marker for Pancreatic and Duodenal Neuroendocrine Tumors. <i>American Journal of Surgical Pathology</i> , 2015 , 39, 850-7	6.7	18
171	RBMV, a novel inhibitor of glycogen synthase kinase 3 β increases tumor stemness and predicts poor prognosis of hepatocellular carcinoma. <i>Hepatology</i> , 2015 , 62, 1480-96	11.2	40
170	The Role of Organelle Stresses in Diabetes Mellitus and Obesity: Implication for Treatment. <i>Analytical Cellular Pathology</i> , 2015 , 2015, 972891	3.4	12
169	Expression of hypoxic marker carbonic anhydrase IX predicts poor prognosis in resectable hepatocellular carcinoma. <i>PLoS ONE</i> , 2015 , 10, e0119181	3.7	22
168	Leiomyosarcoma with alternative lengthening of telomeres is associated with aggressive histologic features, loss of ATRX expression, and poor clinical outcome. <i>American Journal of Surgical Pathology</i> , 2015 , 39, 236-44	6.7	64

167	Connective tissue growth factor inhibits gastric cancer peritoneal metastasis by blocking integrin $\beta 1$ -dependent adhesion. <i>Gastric Cancer</i> , 2015 , 18, 504-15	7.6	19
166	Alternative lengthening of telomeres and loss of ATRX are frequent events in pleomorphic and dedifferentiated liposarcomas. <i>Modern Pathology</i> , 2015 , 28, 1064-73	9.8	36
165	Assessing hepatic fibrosis: comparing the intravoxel incoherent motion in MRI with acoustic radiation force impulse imaging in US. <i>European Radiology</i> , 2015 , 25, 3552-9	8	28
164	IMP-3 promotes migration and invasion of melanoma cells by modulating the expression of HMGA2 and predicts poor prognosis in melanoma. <i>Journal of Investigative Dermatology</i> , 2015 , 135, 1065-1073	4.3	28
163	Targeted next-generation sequencing of cancer genes identified frequent TP53 and ATRX mutations in leiomyosarcoma. <i>American Journal of Translational Research (discontinued)</i> , 2015 , 7, 2072-87		40
162	MLN4924, a Novel NEDD8-activating enzyme inhibitor, exhibits antitumor activity and enhances cisplatin-induced cytotoxicity in human cervical carcinoma: in vitro and in vivo study. <i>American Journal of Cancer Research</i> , 2015 , 5, 3350-62	4.4	9
161	Differential expression of hyaluronan synthase 2 in breast carcinoma and its biological significance. <i>Histopathology</i> , 2014 , 65, 328-39	7.3	29
160	The H3K9 methyltransferase G9a is a marker of aggressive ovarian cancer that promotes peritoneal metastasis. <i>Molecular Cancer</i> , 2014 , 13, 189	42.1	102
159	Morphological subclassification of intrahepatic cholangiocarcinoma: etiological, clinicopathological, and molecular features. <i>Modern Pathology</i> , 2014 , 27, 1163-73	9.8	146
158	TERT promoter mutation in resectable hepatocellular carcinomas: a strong association with hepatitis C infection and absence of hepatitis B infection. <i>International Journal of Surgery</i> , 2014 , 12, 659-65	7.5	47
157	The occurrence of primary hepatic adenoma in deceased donor renal transplant recipient. <i>International Braz J Urol: Official Journal of the Brazilian Society of Urology</i> , 2014 , 40, 118-22	2	1
156	Aryl hydrocarbon receptor downregulates MYCN expression and promotes cell differentiation of neuroblastoma. <i>PLoS ONE</i> , 2014 , 9, e88795	3.7	22
155	Gas6/Axl pathway promotes tumor invasion through the transcriptional activation of Slug in hepatocellular carcinoma. <i>Carcinogenesis</i> , 2014 , 35, 769-75	4.6	81
154	Traditional serrated adenoma has two pathways of neoplastic progression that are distinct from the sessile serrated pathway of colorectal carcinogenesis. <i>Modern Pathology</i> , 2014 , 27, 1375-85	9.8	55
153	Leukocyte cell-derived chemotaxin 2 antagonizes MET receptor activation to suppress hepatocellular carcinoma vascular invasion by protein tyrosine phosphatase 1B recruitment. <i>Hepatology</i> , 2014 , 59, 974-85	11.2	36
152	TERT promoter mutation is uncommon in acral lentiginous melanoma. <i>Journal of Cutaneous Pathology</i> , 2014 , 41, 504-8	1.7	35
151	Quantification of hepatic steatosis: a comparison of the accuracy among multiple magnetic resonance techniques. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2014 , 29, 807-13	4	25
150	Expression of bile duct transcription factor HNF1 α predicts early tumor recurrence and is a stage-independent prognostic factor in hepatocellular carcinoma. <i>Journal of Gastrointestinal Surgery</i> , 2014 , 18, 1784-94	3.3	10

149	Potential synergistic anti-tumor activity between lenalidomide and sorafenib in hepatocellular carcinoma. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2014 , 29, 2021-31	4	10
148	Human cationic trypsinogen but not serine peptidase inhibitor, Kazal type 1 variants increase the risk of type 1 autoimmune pancreatitis. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2014 , 29, 2038-42	4	21
147	Expression of LEF1 is an independent prognostic factor for patients with oral squamous cell carcinoma. <i>Journal of the Formosan Medical Association</i> , 2014 , 113, 934-9	3.2	13
146	Clinical significance and evolution of hepatic HBsAg expression in HBeAg-positive patients receiving interferon therapy. <i>Journal of Gastroenterology</i> , 2014 , 49, 356-62	6.9	13
145	GALNT2 suppresses malignant phenotypes through IGF-1 receptor and predicts favorable prognosis in neuroblastoma. <i>Oncotarget</i> , 2014 , 5, 12247-59	3.3	27
144	Input of microenvironmental regulation on colorectal cancer: role of the CCN family. <i>World Journal of Gastroenterology</i> , 2014 , 20, 6826-31	5.6	16
143	Expression of insulin-like growth factor II mRNA-binding protein 3 predicts early recurrence and poor prognosis in intrahepatic cholangiocarcinoma. <i>International Journal of Surgery</i> , 2013 , 11, 85-91	7.5	18
142	The stem cell E3-ligase Lin-41 promotes liver cancer progression through inhibition of microRNA-mediated gene silencing. <i>Journal of Pathology</i> , 2013 , 229, 486-96	9.4	27
141	Resolution of diabetes after pancreaticoduodenectomy in patients with and without pancreatic ductal cell adenocarcinoma. <i>Annals of Surgical Oncology</i> , 2013 , 20, 242-9	3.1	37
140	β1,4-Galactosyltransferase III enhances invasive phenotypes via β1-integrin and predicts poor prognosis in neuroblastoma. <i>Clinical Cancer Research</i> , 2013 , 19, 1705-16	12.9	34
139	GNAS Is frequently mutated in a specific subgroup of intraductal papillary neoplasms of the bile duct. <i>American Journal of Surgical Pathology</i> , 2013 , 37, 1862-70	6.7	36
138	Histone demethylase RBP2 promotes lung tumorigenesis and cancer metastasis. <i>Cancer Research</i> , 2013 , 73, 4711-21	10.1	113
137	Overexpression of CTHRC1 in hepatocellular carcinoma promotes tumor invasion and predicts poor prognosis. <i>PLoS ONE</i> , 2013 , 8, e70324	3.7	62
136	Development of autoantibodies after pediatric liver transplantation. <i>Pediatric Transplantation</i> , 2013 , 17, 144-8	1.8	14
135	Long-term follow-up of children with postnatal immunoprophylaxis failure who were infected with hepatitis B virus surface antigen gene mutant. <i>Journal of Infectious Diseases</i> , 2013 , 207, 1047-57	7	22
134	B3GNT3 expression suppresses cell migration and invasion and predicts favorable outcomes in neuroblastoma. <i>Cancer Science</i> , 2013 , 104, 1600-8	6.9	29
133	Underexpression of hepatocyte nuclear factor-1α in chromophobe renal cell carcinoma. <i>Histopathology</i> , 2013 , 62, 589-94	7.3	14
132	Annexin A10 is an immunohistochemical marker for adenocarcinoma of the upper gastrointestinal tract and pancreatobiliary system. <i>Histopathology</i> , 2013 , 63, 640-8	7.3	28

131	Loss of corepressor PER2 under hypoxia up-regulates OCT1-mediated EMT gene expression and enhances tumor malignancy. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013 , 110, 12331-6	11.5	91
130	Incorporation of the factor IX Padua mutation into FIX-Triple improves clotting activity in vitro and in vivo. <i>Thrombosis and Haemostasis</i> , 2013 , 110, 244-56	7	26
129	HSD3B1 is a specific trophoblast-associated marker not expressed in a wide spectrum of tumors. <i>International Journal of Gynecological Cancer</i> , 2013 , 23, 343-7	3.5	12
128	p15(PAF) is an Rb/E2F-regulated S-phase protein essential for DNA synthesis and cell cycle progression. <i>PLoS ONE</i> , 2013 , 8, e61196	3.7	31
127	Upregulation of endocan by Epstein-Barr virus latent membrane protein 1 and its clinical significance in nasopharyngeal carcinoma. <i>PLoS ONE</i> , 2013 , 8, e82254	3.7	13
126	Elevated preoperative serum CA19-9 levels in patients with hepatocellular carcinoma is associated with poor prognosis after resection. <i>Scientific World Journal, The</i> , 2013 , 2013, 380797	2.2	9
125	Angiopoietin-like protein 1 suppresses SLUG to inhibit cancer cell motility. <i>Journal of Clinical Investigation</i> , 2013 , 123, 1082-95	15.9	65
124	Relationship between ultrasound backscattered statistics and the concentration of fatty droplets in livers: an animal study. <i>PLoS ONE</i> , 2013 , 8, e63543	3.7	26
123	S100P expression is a novel prognostic factor in hepatocellular carcinoma and predicts survival in patients with high tumor stage or early recurrent tumors. <i>PLoS ONE</i> , 2013 , 8, e65501	3.7	22
122	Utility of hepatocyte nuclear factor-1 β s as a diagnostic marker in ovarian carcinomas with clear cells. <i>Histopathology</i> , 2012 , 61, 760-8	7.3	31
121	SIRT1 promotes tumorigenesis and resistance to chemotherapy in hepatocellular carcinoma and its expression predicts poor prognosis. <i>Annals of Surgical Oncology</i> , 2012 , 19, 2011-9	3.1	116
120	Persistent elevation of hepatocyte growth factor activator inhibitors in cholangiopathies affects liver fibrosis and differentiation. <i>Hepatology</i> , 2012 , 55, 161-72	11.2	9
119	S100P immunostaining identifies a subset of peripheral-type intrahepatic cholangiocarcinomas with morphological and molecular features similar to those of perihilar and extrahepatic cholangiocarcinomas. <i>Histopathology</i> , 2012 , 61, 1106-16	7.3	29
118	Loss of the oxidative stress sensor NPGPx compromises GRP78 chaperone activity and induces systemic disease. <i>Molecular Cell</i> , 2012 , 48, 747-59	17.6	106
117	TET1 suppresses cancer invasion by activating the tissue inhibitors of metalloproteinases. <i>Cell Reports</i> , 2012 , 2, 568-79	10.6	214
116	Prognostic significance of TWEAK expression in colorectal cancer and effect of its inhibition on invasion. <i>Annals of Surgical Oncology</i> , 2012 , 19 Suppl 3, S385-94	3.1	14
115	Clostridium difficile infection in a patient with Crohn disease. <i>Journal of the Formosan Medical Association</i> , 2012 , 111, 347-9	3.2	2
114	MicroRNA 3Q and nucleotide modification patterns and arm selection preference in liver tissues. <i>BMC Systems Biology</i> , 2012 , 6 Suppl 2, S14	3.5	29

113	Calreticulin mediates nerve growth factor-induced neuronal differentiation. <i>Journal of Molecular Neuroscience</i> , 2012 , 47, 571-81	3.3	13
112	Adenylate kinase-4 is a marker of poor clinical outcomes that promotes metastasis of lung cancer by downregulating the transcription factor ATF3. <i>Cancer Research</i> , 2012 , 72, 5119-29	10.1	58
111	Activation of Robo1 signaling of breast cancer cells by Slit2 from stromal fibroblast restrains tumorigenesis via blocking PI3K/Akt/βcatenin pathway. <i>Cancer Research</i> , 2012 , 72, 4652-61	10.1	100
110	Hepatocyte growth factor activates Wnt pathway by transcriptional activation of LEF1 to facilitate tumor invasion. <i>Carcinogenesis</i> , 2012 , 33, 1142-8	4.6	40
109	Clinical features and treatment responses of children with eosinophilic gastroenteritis. <i>Pediatrics and Neonatology</i> , 2011 , 52, 272-8	1.8	38
108	B4GALNT3 expression predicts a favorable prognosis and suppresses cell migration and invasion via Integrin signaling in neuroblastoma. <i>American Journal of Pathology</i> , 2011 , 179, 1394-404	5.8	26
107	Mutation and loss of expression of ARID1A in uterine low-grade endometrioid carcinoma. <i>American Journal of Surgical Pathology</i> , 2011 , 35, 625-32	6.7	206
106	Male germ cell-specific RNA binding protein RBMY: a new oncogene explaining male predominance in liver cancer. <i>PLoS ONE</i> , 2011 , 6, e26948	3.7	26
105	Insulin-like growth factor II mRNA-binding protein 3 expression predicts unfavorable prognosis in patients with neuroblastoma. <i>Cancer Science</i> , 2011 , 102, 2191-8	6.9	21
104	Insulin-like growth factor II mRNA-binding protein 3 expression promotes tumor formation and invasion and predicts poor prognosis in oral squamous cell carcinoma. <i>Journal of Oral Pathology and Medicine</i> , 2011 , 40, 699-705	3.3	20
103	Role of p53 and βcatenin mutations in conjunction with CK19 expression on early tumor recurrence and prognosis of hepatocellular carcinoma. <i>Journal of Gastrointestinal Surgery</i> , 2011 , 15, 321-9	3.3	53
102	Connective tissue growth factor acts as a therapeutic agent and predictor for peritoneal carcinomatosis of colorectal cancer. <i>Clinical Cancer Research</i> , 2011 , 17, 3077-88	12.9	27
101	Depletion of βcatenin from mature hepatocytes of mice promotes expansion of hepatic progenitor cells and tumor development. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011 , 108, 18384-9	11.5	31
100	IL-25 causes apoptosis of IL-25R-expressing breast cancer cells without toxicity to nonmalignant cells. <i>Science Translational Medicine</i> , 2011 , 3, 78ra31	17.5	75
99	Phosphorylation of NuSAP by Cdk1 regulates its interaction with microtubules in mitosis. <i>Cell Cycle</i> , 2011 , 10, 4083-9	4.7	14
98	Viral hepatitis is associated with intrahepatic cholangiocarcinoma with cholangiolar differentiation and N-cadherin expression. <i>Modern Pathology</i> , 2011 , 24, 810-9	9.8	49
97	Histone deacetylase 6 regulates estrogen receptor alpha in uterine leiomyoma. <i>Reproductive Sciences</i> , 2011 , 18, 755-62	3	24
96	Downregulation of alpha-fetoprotein expression by LHX4: a critical role in hepatocarcinogenesis. <i>Carcinogenesis</i> , 2011 , 32, 1815-23	4.6	21

95	Nuclear GRP75 binds retinoic acid receptors to promote neuronal differentiation of neuroblastoma. <i>PLoS ONE</i> , 2011 , 6, e26236	3.7	20
94	Ex vivo imaging and quantification of liver fibrosis using second-harmonic generation microscopy. <i>Journal of Biomedical Optics</i> , 2010 , 15, 036002	3.5	24
93	MicroRNA-519c suppresses hypoxia-inducible factor-1alpha expression and tumor angiogenesis. <i>Cancer Research</i> , 2010 , 70, 2675-85	10.1	163
92	Lin-28B expression promotes transformation and invasion in human hepatocellular carcinoma. <i>Carcinogenesis</i> , 2010 , 31, 1516-22	4.6	85
91	H3K9 histone methyltransferase G9a promotes lung cancer invasion and metastasis by silencing the cell adhesion molecule Ep-CAM. <i>Cancer Research</i> , 2010 , 70, 7830-40	10.1	274
90	Prognostic implication of MET overexpression in myxofibrosarcomas: an integrative array comparative genomic hybridization, real-time quantitative PCR, immunoblotting, and immunohistochemical analysis. <i>Modern Pathology</i> , 2010 , 23, 1379-92	9.8	24
89	Notch1 expression predicts an unfavorable prognosis and serves as a therapeutic target of patients with neuroblastoma. <i>Clinical Cancer Research</i> , 2010 , 16, 4411-20	12.9	35
88	Survival after pancreaticoduodenectomy for ampullary cancer is not affected by age. <i>World Journal of Surgery</i> , 2010 , 34, 2945-52	3.3	13
87	Significance of Aurora B overexpression in hepatocellular carcinoma. Aurora B Overexpression in HCC. <i>BMC Cancer</i> , 2010 , 10, 461	4.8	83
86	Clinical course of de novo hepatitis B infection after pediatric liver transplantation. <i>Liver Transplantation</i> , 2010 , 16, 215-21	4.5	15
85	Radical endoscopic polypectomy combined with double-balloon enteroscopy and colonoscopy for Peutz-Jeghers syndrome. <i>Journal of Pediatric Gastroenterology and Nutrition</i> , 2010 , 51, 370-2	2.8	3
84	Nuclear overexpression of mitotic regulatory proteins in biliary tract cancer: correlation with clinicopathologic features and patient survival. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2009 , 18, 417-23	4	22
83	Synergistic effect of radiation and interleukin-6 on hepatitis B virus reactivation in liver through STAT3 signaling pathway. <i>International Journal of Radiation Oncology Biology Physics</i> , 2009 , 75, 1545-52	4	23
82	Identification of calreticulin as a prognosis marker and angiogenic regulator in human gastric cancer. <i>Annals of Surgical Oncology</i> , 2009 , 16, 524-33	3.1	90
81	Diffuse expression of RNA-binding protein IMP3 predicts high-stage lymph node metastasis and poor prognosis in colorectal adenocarcinoma. <i>Annals of Surgical Oncology</i> , 2009 , 16, 1711-9	3.1	44
80	The Aurora kinase inhibitor VE-465 has anticancer effects in pre-clinical studies of human hepatocellular carcinoma. <i>Journal of Hepatology</i> , 2009 , 50, 518-27	13.4	39
79	Peginterferon alfa-2a plus ribavirin for the treatment of dual chronic infection with hepatitis B and C viruses. <i>Gastroenterology</i> , 2009 , 136, 496-504.e3	13.3	183
78	Emerging importance of autoimmune hepatitis in children in Taiwan, an endemic area for viral hepatitis. <i>Pediatrics and Neonatology</i> , 2009 , 50, 65-9	1.8	6

77	X-linked liver glycogenosis in a Taiwanese family: transmission from undiagnosed males. <i>Pediatrics and Neonatology</i> , 2009 , 50, 230-3	1.8	6
76	Hepatitis B virus basal core promoter mutation and DNA load correlate with expression of hepatitis B core antigen in patients with chronic hepatitis B. <i>Journal of Infectious Diseases</i> , 2009 , 199, 742-9	7	25
75	Multiple lineages of human breast cancer stem/progenitor cells identified by profiling with stem cell markers. <i>PLoS ONE</i> , 2009 , 4, e8377	3.7	169
74	Cadherin-17 is a useful diagnostic marker for adenocarcinomas of the digestive system. <i>Modern Pathology</i> , 2008 , 21, 1379-86	9.8	87
73	Acute Q fever presenting as fever of unknown origin with rapidly progressive hepatic failure in a patient with alcoholism. <i>Journal of the Formosan Medical Association</i> , 2008 , 107, 896-901	3.2	6
72	ASPM is a novel marker for vascular invasion, early recurrence, and poor prognosis of hepatocellular carcinoma. <i>Clinical Cancer Research</i> , 2008 , 14, 4814-20	12.9	92
71	Identification of GRP75 as an independent favorable prognostic marker of neuroblastoma by a proteomics analysis. <i>Clinical Cancer Research</i> , 2008 , 14, 6237-45	12.9	26
70	Expression of hepatocyte transporters and nuclear receptors in children with early and late-stage biliary atresia. <i>Pediatric Research</i> , 2008 , 63, 667-73	3.2	68
69	HSD3B1 as a novel trophoblast-associated marker that assists in the differential diagnosis of trophoblastic tumors and tumorlike lesions. <i>American Journal of Surgical Pathology</i> , 2008 , 32, 236-42	6.7	45
68	Bone marrow transplantation results in donor-derived hepatocytes in an animal model of inherited cholestatic liver disease. <i>Journal of Biomedical Science</i> , 2008 , 15, 615-22	13.3	6
67	Nuclear localization of annexin A1 is a prognostic factor in oral squamous cell carcinoma. <i>Journal of Surgical Oncology</i> , 2008 , 97, 544-50	2.8	44
66	RNA-binding protein insulin-like growth factor II mRNA-binding protein 3 expression promotes tumor invasion and predicts early recurrence and poor prognosis in hepatocellular carcinoma. <i>Hepatology</i> , 2008 , 48, 1118-27	11.2	121
65	IL-6 induces AGS gastric cancer cell invasion via activation of the c-Src/RhoA/ROCK signaling pathway. <i>International Journal of Cancer</i> , 2007 , 120, 2600-8	7.5	107
64	Differential DNA methylation associated with hepatitis B virus infection in hepatocellular carcinoma. <i>International Journal of Cancer</i> , 2007 , 121, 1257-64	7.5	63
63	Expression of the caudal-type homeodomain transcription factor CDX2 is related to clinical outcome in biliary tract carcinoma. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2007 , 22, 389-94	4.4	21
62	Cytoplasmic CD24 expression is a novel prognostic factor in diffuse-type gastric adenocarcinoma. <i>Annals of Surgical Oncology</i> , 2007 , 14, 2748-58	3.1	50
61	Management of hepatic angiomyolipoma. <i>Journal of Gastrointestinal Surgery</i> , 2007 , 11, 452-7	3.3	105
60	Overexpression of KIAA0101 predicts high stage, early tumor recurrence, and poor prognosis of hepatocellular carcinoma. <i>Clinical Cancer Research</i> , 2007 , 13, 5368-76	12.9	63

59	Oncologic results of laparoscopic versus conventional open surgery for stage II or III left-sided colon cancers: a randomized controlled trial. <i>Annals of Surgical Oncology</i> , 2007 , 14, 109-17	3.1	134
58	The VEGF-C/Flt-4 axis promotes invasion and metastasis of cancer cells. <i>Cancer Cell</i> , 2006 , 9, 209-23	24.3	273
57	Removal of BRCA1/CtIP/ZBRK1 repressor complex on ANG1 promoter leads to accelerated mammary tumor growth contributed by prominent vasculature. <i>Cancer Cell</i> , 2006 , 10, 13-24	24.3	84
56	Knockdown of contactin-1 expression suppresses invasion and metastasis of lung adenocarcinoma. <i>Cancer Research</i> , 2006 , 66, 2553-61	10.1	57
55	Effect of connective tissue growth factor on hypoxia-inducible factor 1alpha degradation and tumor angiogenesis. <i>Journal of the National Cancer Institute</i> , 2006 , 98, 984-95	9.7	65
54	Proline-directed protein kinase FA is a powerful and independent prognostic predictor for progression and patient survival of hepatocellular carcinoma. <i>Journal of Clinical Oncology</i> , 2006 , 24, 3780-8	2.3	21
53	CD24 expression is a prognostic factor in intrahepatic cholangiocarcinoma. <i>Cancer Letters</i> , 2006 , 235, 34-9	9.9	46
52	Hepatic steatosis and neonatal intrahepatic cholestasis caused by citrin deficiency (NICCD) in Taiwanese infants. <i>Journal of Pediatrics</i> , 2006 , 148, 642-6	3.6	32
51	Post-transplantation lymphoproliferative disorders localizing to the gastrointestinal tract after liver transplantation: report of five pediatric cases. <i>Pediatric Transplantation</i> , 2006 , 10, 390-4	1.8	47
50	Decreased expressions of hepsin in human hepatocellular carcinomas. <i>Liver International</i> , 2006 , 26, 774-80	8.9	17
49	Nuclear translocation of beta-catenin protein but absence of beta-catenin and APC mutation in gastrointestinal carcinoid tumor. <i>Annals of Surgical Oncology</i> , 2006 , 13, 1604-9	3.1	24
48	Aberrant expressions of annexin A10 short isoform, osteopontin and E-cadherin at chromosome 4q cooperatively contribute to progression and poor prognosis of hepatocellular carcinoma 2005 , 26, 1053		2
47	Serum adiponectin correlates with viral characteristics but not histologic features in patients with chronic hepatitis C. <i>Journal of Hepatology</i> , 2005 , 43, 235-42	13.4	59
46	Absence of epidermal growth factor receptor exon 18-21 mutation in hepatocellular carcinoma. <i>Cancer Letters</i> , 2005 , 224, 117-21	9.9	45
45	Connective tissue growth factor inhibits metastasis and acts as an independent prognostic marker in colorectal cancer. <i>Gastroenterology</i> , 2005 , 128, 9-23	13.3	108
44	Liver steatosis classification using high-frequency ultrasound. <i>Ultrasound in Medicine and Biology</i> , 2005 , 31, 599-605	3.5	20
43	Lamivudine treatment in maternally transmitted chronic hepatitis B virus infection patients. <i>Pediatrics International</i> , 2005 , 47, 372-7	1.2	15
42	Differential integration rates of hepatitis B virus DNA in the liver of children with chronic hepatitis B virus infection and hepatocellular carcinoma. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2005 , 20, 1206-14	4	5

41	HMB-45 may be a more sensitive maker than S-100 or Melan-A for immunohistochemical diagnosis of primary oral and nasal mucosal melanomas. <i>Journal of Oral Pathology and Medicine</i> , 2005 , 34, 540-5	3.3	44
40	Solid pseudopapillary neoplasms of the pancreas: is there a pathologic basis for the observed gender differences in incidence?. <i>Surgery</i> , 2005 , 137, 591-6	3.6	53
39	Unexpected rapid progression of metastatic adenoid cystic carcinoma during treatment with imatinib mesylate. <i>Head and Neck</i> , 2005 , 27, 1022-7	4.2	36
38	GRP78 expression correlates with histologic differentiation and favorable prognosis in neuroblastic tumors. <i>International Journal of Cancer</i> , 2005 , 113, 920-7	7.5	42
37	Biliary hamartomas with delayed 99mTc-diisopropyl iminodiacetic acid clearance. <i>Journal of Gastroenterology</i> , 2005 , 40, 540-4	6.9	3
36	Intestinal Behçet disease presenting as neonatal onset chronic diarrhea in an 11-month-old male baby. <i>European Journal of Pediatrics</i> , 2005 , 164, 523-5	4.1	8
35	Cyr61 induces gastric cancer cell motility/invasion via activation of the integrin/nuclear factor-kappaB/cyclooxygenase-2 signaling pathway. <i>Clinical Cancer Research</i> , 2005 , 11, 5809-20	12.9	82
34	Opposite roles of human pancreatitis-associated protein and REG1A expression in hepatocellular carcinoma: association of pancreatitis-associated protein expression with low-stage hepatocellular carcinoma, beta-catenin mutation, and favorable prognosis. <i>Clinical Cancer Research</i> , 2005 , 11, 2568-75	12.9	37
33	Aberrant expressions of annexin A10 short isoform, osteopontin and alpha-fetoprotein at chromosome 4q cooperatively contribute to progression and poor prognosis of hepatocellular carcinoma. <i>International Journal of Oncology</i> , 2005 , 26, 1053-61	1	8
32	Cyclooxygenase-2 induces EP1- and HER-2/Neu-dependent vascular endothelial growth factor-C up-regulation: a novel mechanism of lymphangiogenesis in lung adenocarcinoma. <i>Cancer Research</i> , 2004 , 64, 554-64	10.1	172
31	Overexpression and amplification of Aurora-A in hepatocellular carcinoma. <i>Clinical Cancer Research</i> , 2004 , 10, 2065-71	12.9	253
30	Intravasation-related metastatic factors in colorectal cancer. <i>Tumor Biology</i> , 2004 , 25, 48-55	2.9	12
29	Connective tissue growth factor and its role in lung adenocarcinoma invasion and metastasis. <i>Journal of the National Cancer Institute</i> , 2004 , 96, 364-75	9.7	138
28	Overexpression of Her-2/neu in Epithelial Ovarian Carcinoma Induces Vascular Endothelial Growth Factor C by Activating NF- κ B: Implications for Malignant Ascites Formation and Tumor Lymphangiogenesis. <i>Journal of Biomedical Science</i> , 2004 , 11, 249-259	13.3	3
27	RBMV, a male germ cell-specific RNA-binding protein, activated in human liver cancers and transforms rodent fibroblasts. <i>Oncogene</i> , 2004 , 23, 5815-22	9.2	33
26	Clinical relevance of hepatitis B virus genotype in children with chronic infection and hepatocellular carcinoma. <i>Gastroenterology</i> , 2004 , 127, 1733-8	13.3	173
25	Desmoplastic small round cell tumor of the kidney. <i>American Journal of Surgical Pathology</i> , 2004 , 28, 1379-83	6.7	37
24	Ribavirin and interferon is effective for hepatitis C virus clearance in hepatitis B and C dually infected patients. <i>Hepatology</i> , 2003 , 37, 568-76	11.2	94

23	Comparison of medial-to-lateral versus traditional lateral-to-medial laparoscopic dissection sequences for resection of rectosigmoid cancers: randomized controlled clinical trial. <i>World Journal of Surgery</i> , 2003 , 27, 190-6	3.3	60
22	KLF6, a putative tumor suppressor gene, is mutated in astrocytic gliomas. <i>International Journal of Cancer</i> , 2003 , 105, 625-9	7.5	87
21	Elastic modulus measurements of human liver and correlation with pathology. <i>Ultrasound in Medicine and Biology</i> , 2002 , 28, 467-74	3.5	314
20	P53 gene and Wnt signaling in benign neoplasms: beta-catenin mutations in hepatic adenoma but not in focal nodular hyperplasia. <i>Hepatology</i> , 2002 , 36, 927-35	11.2	103
19	Sinonasal undifferentiated carcinoma and nasopharyngeal-type undifferentiated carcinoma: two clinically, biologically, and histopathologically distinct entities. <i>American Journal of Surgical Pathology</i> , 2002 , 26, 371-6	6.7	124
18	2 Role of immunohistochemical expression of p53 in colorectal carcinoma. <i>Handbook of Immunohistochemistry and in Situ Hybridization of Human Carcinomas</i> , 2002 , 139-147		
17	Down-regulation of annexin A10 in hepatocellular carcinoma is associated with vascular invasion, early recurrence, and poor prognosis in synergy with p53 mutation. <i>American Journal of Pathology</i> , 2002 , 160, 1831-7	5.8	66
16	Multiple intrahepatic biliary cysts in children with biliary atresia. <i>Journal of Pediatric Surgery</i> , 2002 , 37, 1183-7	2.6	35
15	FIC1 and BSEP defects in Taiwanese patients with chronic intrahepatic cholestasis with low gamma-glutamyltranspeptidase levels. <i>Journal of Pediatrics</i> , 2002 , 140, 119-24	3.6	77
14	Mutation of the DR5/TRAIL receptor 2 gene is infrequent in hepatocellular carcinoma. <i>Cancer Letters</i> , 2002 , 181, 205-8	9.9	28
13	p53 gene and Wnt signaling in benign neoplasms: Eatenin mutations in hepatic adenoma but not in focal nodular hyperplasia. <i>Hepatology</i> , 2002 , 36, 927-935	11.2	6
12	Expression of mutant nuclear beta-catenin correlates with non-invasive hepatocellular carcinoma, absence of portal vein spread, and good prognosis. <i>Journal of Pathology</i> , 2001 , 193, 95-101	9.4	94
11	Progressive familial intrahepatic cholestasis with high gamma-glutamyltranspeptidase levels in Taiwanese infants: role of MDR3 gene defect?. <i>Pediatric Research</i> , 2001 , 50, 50-5	3.2	31
10	Lymphoepithelioma-like cholangiocarcinoma: an Epstein-Barr virus-associated tumor. <i>American Journal of Surgical Pathology</i> , 2001 , 25, 516-20	6.7	67
9	Prevalence and rapid identification of clarithromycin-resistant <i>Helicobacter pylori</i> isolates in children. <i>Pediatric Infectious Disease Journal</i> , 2001 , 20, 662-6	3.4	26
8	Congenital interstitial cell of cajal hyperplasia with neuronal intestinal dysplasia. <i>American Journal of Surgical Pathology</i> , 2000 , 24, 1568-72	6.7	28
7	Beta-catenin mutations are associated with a subset of low-stage hepatocellular carcinoma negative for hepatitis B virus and with favorable prognosis. <i>American Journal of Pathology</i> , 2000 , 157, 763-70	5.8	256
6	Expression of the c-kit protein is associated with certain subtypes of salivary gland carcinoma. <i>Cancer Letters</i> , 2000 , 154, 107-11	9.9	161

5	Somatic mutations of beta-catenin play a crucial role in the tumorigenesis of sporadic hepatoblastoma. <i>Cancer Letters</i> , 2000 , 152, 45-51	9.9	95
4	Phenytoin-induced pseudolymphoma: reevaluation using modern molecular biology techniques. <i>Epilepsia</i> , 1996 , 37, 104-7	6.4	18
3	Concordant body stalk anomaly in monozygotic twinning--early embryo cleavage disorder. <i>Prenatal Diagnosis</i> , 1996 , 16, 467-70	3.2	16
2	CONCORDANT BODY STALK ANOMALY IN MONOZYGOTIC TWINNING--EARLY EMBRYO CLEAVAGE DISORDER 1996 , 16, 467		1
1	Young's modulus measurements of human liver and correlation with pathological findings		2