Akio Kihara

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/425530/akio-kihara-publications-by-year.pdf

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

151	15,421	51	123
papers	citations	h-index	g-index
164	17,244 ext. citations	5.2	6.32
ext. papers		avg, IF	L-index

#	Paper	IF	Citations
151	Formation of fatty alcohols-components of meibum lipids-by the fatty acyl-CoA reductase FAR2 is essential for dry eye prevention <i>FASEB Journal</i> , 2022 , 36, e22216	0.9	2
150	Hypomyelinating spastic dyskinesia and ichthyosis caused by a homozygous splice site mutation leading to exon skipping in ELOVL1 <i>Brain and Development</i> , 2022 ,	2.2	2
149	Whole picture of human stratum corneum ceramides, including the chain-length diversity of long-chain bases. <i>Journal of Lipid Research</i> , 2022 , 100235	6.3	2
148	Improvement of Evaporative Dry Eye With Meibomian Gland Dysfunction in Model Mice by Treatment With Ophthalmic Solution Containing Mineral Oil. <i>Translational Vision Science and Technology</i> , 2021 , 10, 21	3.3	2
147	Diverse meibum lipids produced by Awat1 and Awat2 are important for stabilizing tear film and protecting the ocular surface. <i>IScience</i> , 2021 , 24, 102478	6.1	6
146	Amlexanox enhances the antitumor effect of anti-PD-1 antibody. <i>Biochemical and Biophysical Research Communications</i> , 2021 , 560, 1-6	3.4	1
145	Production of branched-chain very-long-chain fatty acids by fatty acid elongases and their tissue distribution in mammals. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2021 , 1866, 158842	5	7
144	Impaired production of the skin barrier lipid acylceramide by CYP4F22 ichthyosis mutations. <i>Journal of Dermatological Science</i> , 2021 , 101, 69-71	4.3	1
143	Direct uptake of sphingosine-1-phosphate independent of phospholipid phosphatases. <i>Journal of Biological Chemistry</i> , 2021 , 296, 100605	5.4	3
142	Comprehensive stratum corneum ceramide profiling reveals reduced acylceramides in ichthyosis patient with CERS3 mutations. <i>Journal of Dermatology</i> , 2021 , 48, 447-456	1.6	6
141	Protein-bound ceramide levels in the epidermis of transglutaminase 1-deficient mice. <i>Journal of Dermatology</i> , 2021 , 48, 1799-1801	1.6	
140	Impaired Skin Barrier Function Due to Reduced & Acylceramide Levels in a Mouse Model of Sjigren-Larsson Syndrome. <i>Molecular and Cellular Biology</i> , 2021 , 41, e0035221	4.8	1
139	Comparative profiling and comprehensive quantification of stratum corneum ceramides in humans and mice by LC/MS/MS. <i>Journal of Lipid Research</i> , 2020 , 61, 884-895	6.3	26
138	Catalytic residues, substrate specificity, and role in carbon starvation of the 2-hydroxy FA dioxygenase Mpo1 in yeast. <i>Journal of Lipid Research</i> , 2020 , 61, 1104-1114	6.3	0
137	FTY720 Protects Against Ischemia-Reperfusion Injury by Preventing the Redistribution of Tight Junction Proteins and Decreases Inflammation in the Subacute Phase in an Experimental Stroke Model. <i>Translational Stroke Research</i> , 2020 , 11, 1103-1116	7.8	15
136	Biosynthesis of the anti-lipid-microdomain sphingoid base 4,14-sphingadiene by the ceramide desaturase FADS3. <i>FASEB Journal</i> , 2020 , 34, 3318-3335	0.9	18
135	Skin permeability barrier formation by the ichthyosis-causative gene through formation of the barrier lipid ⊞acylceramide. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020 , 117, 2914-2922	11.5	28

(2018-2020)

134	Lipid polarity gradient formed by Ehydroxy lipids in tear film prevents dry eye disease. <i>ELife</i> , 2020 , 9,	8.9	27
133	N-glycosylation of Rim21 at an Unconventional Site Fine-tunes Its Behavior in the Plasma Membrane. <i>Cell Structure and Function</i> , 2020 , 45, 1-8	2.2	3
132	Novel biallelic FA2H mutations in a Japanese boy with fatty acid hydroxylase-associated neurodegeneration. <i>Brain and Development</i> , 2020 , 42, 217-221	2.2	4
131	Severe Skin Permeability Barrier Dysfunction in Knockout Mice Deficient in a Fatty Acid Hydroxylase Crucial to Acylceramide Production. <i>Journal of Investigative Dermatology</i> , 2020 , 140, 319-3	3 2 6.e4	12
130	The very-long-chain fatty acid elongase Elo2 rescues lethal defects associated with loss of the nuclear barrier function in fission yeast cells. <i>Journal of Cell Science</i> , 2019 , 132,	5.3	15
129	Neural symptoms in a gene knockout mouse model of Sjgren-Larsson syndrome are associated with a decrease in 2-hydroxygalactosylceramide. <i>FASEB Journal</i> , 2019 , 33, 928-941	0.9	14
128	Reduced chain length in myelin sphingolipids and poorer motor coordination in mice deficient in the fatty acid elongase. <i>FASEB BioAdvances</i> , 2019 , 1, 747-759	2.8	11
127	Yeast Mpo1 Is a Novel Dioxygenase That Catalyzes the EOxidation of a 2-Hydroxy Fatty Acid in an Fe-Dependent Manner. <i>Molecular and Cellular Biology</i> , 2019 , 39,	4.8	4
126	De novo mutation in causes ichthyosis, , hypomyelination, spastic paraplegia, high frequency deafness and optic atrophy. <i>Journal of Medical Genetics</i> , 2019 , 56, 164-175	5.8	35
125	The role of PNPLA1 in EO-acylceramide synthesis and skin barrier function. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2019 , 1864, 869-879	5	24
124	Decreased Skin Barrier Lipid Acylceramide and Differentiation-Dependent Gene Expression in Ichthyosis Gene Nipal4-Knockout Mice. <i>Journal of Investigative Dermatology</i> , 2018 , 138, 741-749	4.3	10
123	Sphingolipids activate the endoplasmic reticulum stress surveillance pathway. <i>Journal of Cell Biology</i> , 2018 , 217, 495-505	7.3	21
122	Very long-chain tear film lipids produced by fatty acid elongase ELOVL1 prevent dry eye disease in mice. <i>FASEB Journal</i> , 2018 , 32, 2966-2978	0.9	40
121	Sphingosine 1-phosphate receptor modulator ONO-4641 stimulates CD11bGr-1 cell expansion and inhibits lymphocyte infiltration in the lungs to ameliorate murine pulmonary emphysema. <i>Mucosal Immunology</i> , 2018 , 11, 1606-1620	9.2	11
120	Metabolism of long-chain bases of sphingolipids and fatty acid Ebxidation. <i>Plant Morphology</i> , 2018 , 30, 5-14	О	
119	Structure-inspired design of a sphingolipid mimic sphingosine-1-phosphate receptor agonist from a naturally occurring sphingomyelin synthase inhibitor. <i>Chemical Communications</i> , 2018 , 54, 12758-12761	5.8	6
118	Molecular mechanism of the ichthyosis pathology of Chanarin-Dorfman syndrome: Stimulation of PNPLA1-catalyzed EO-acylceramide production by ABHD5. <i>Journal of Dermatological Science</i> , 2018 , 92, 245-253	4.3	23
117	Widespread tissue distribution and synthetic pathway of polyunsaturated C24:2 sphingolipids in mammals. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2018 , 1863, 1441-1448	5	7

116	PNPLA1 is a transacylase essential for the generation of the skin barrier lipid ED-acylceramide. <i>Nature Communications</i> , 2017 , 8, 14610	17.4	61
115	Systematic analysis of Ca homeostasis in based on chemical-genetic interaction profiles. <i>Molecular Biology of the Cell</i> , 2017 , 28, 3415-3427	3.5	7
114	Phytosphingosine degradation pathway includes fatty acid Ebxidation reactions in the endoplasmic reticulum. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017 , 114, E2616-E2623	11.5	27
113	Decreases in 15-lipoxygenase metabolites in Olmsted syndrome model rats. <i>Journal of Dermatological Science</i> , 2017 , 85, 186-196	4.3	4
112	The Rim101 pathway contributes to ER stress adaptation through sensing the state of plasma membrane. <i>Biochemical Journal</i> , 2017 , 474, 51-63	3.8	11
111	Loop 5 region is important for the activity of the long-chain base transporter Rsb1. <i>Journal of Biochemistry</i> , 2017 , 161, 207-213	3.1	O
110	Biallelic Mutations in KDSR Disrupt Ceramide Synthesis and Result in alspectrum of Keratinization Disorders Associated with Thrombocytopenia. <i>Journal of Investigative Dermatology</i> , 2017 , 137, 2344-23	5 ⁴ 3·3	35
109	The 3-hydroxyacyl-CoA dehydratases HACD1 and HACD2 exhibit functional redundancy and are active in a wide range of fatty acid elongation pathways. <i>Journal of Biological Chemistry</i> , 2017 , 292, 155	3 § :4155	55 ¹ 1 ⁵
108	??????????????????????????????????????	0	
107	Guidelines for the use and interpretation of assays for monitoring autophagy (3rd edition). <i>Autophagy</i> , 2016 , 12, 1-222	10.2	3838
106	A role of the sphingosine-1-phosphate (S1P)-S1P receptor 2 pathway in epithelial defense against cancer (EDAC). <i>Molecular Biology of the Cell</i> , 2016 , 27, 491-9	3.5	29
105	AM251 Suppresses Epithelial-Mesenchymal Transition of Renal Tubular Epithelial Cells. <i>PLoS ONE</i> , 2016 , 11, e0167848	3.7	13
104	Disruption of the Sjgren-Larsson Syndrome Gene Aldh3a2 in Mice Increases Keratinocyte Growth and Retards Skin Barrier Recovery. <i>Journal of Biological Chemistry</i> , 2016 , 291, 11676-88	5.4	24
103	Long-chain bases of sphingolipids are transported into cells via the acyl-CoA synthetases. <i>Scientific Reports</i> , 2016 , 6, 25469	4.9	17
102	Enzyme Activities of the Ceramide Synthases CERS2-6 Are Regulated by Phosphorylation in the C-terminal Region. <i>Journal of Biological Chemistry</i> , 2016 , 291, 7477-87	5.4	37
101	Synthesis and degradation pathways, functions, and pathology of ceramides and epidermal acylceramides. <i>Progress in Lipid Research</i> , 2016 , 63, 50-69	14.3	108
100	Mechanistic Details of Early Steps in Coenzyme Q Biosynthesis Pathway in Yeast. <i>Cell Chemical</i>	8.2	51
	Biology, 2016 , 23, 1241-1250		

(2013-2015)

98	Histological analyses by matrix-assisted laser desorption/ionization-imaging mass spectrometry reveal differential localization of sphingomyelin molecular species regulated by particular ceramide synthase in mouse brains. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2015 ,	5	17
97	1851, 1554-65 HACD1, a regulator of membrane composition and fluidity, promotes myoblast fusion and skeletal muscle growth. <i>Journal of Molecular Cell Biology</i> , 2015 , 7, 429-40	6.3	27
96	The C-terminal Cytosolic Region of Rim21 Senses Alterations in Plasma Membrane Lipid Composition: INSIGHTS INTO SENSING MECHANISMS FOR PLASMA MEMBRANE LIPID ASYMMETRY. Journal of Biological Chemistry, 2015 , 290, 30797-805	5.4	19
95	Essential role of the cytochrome P450 CYP4F22 in the production of acylceramide, the key lipid for skin permeability barrier formation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2015 , 112, 7707-12	11.5	102
94	Sphingolipid Metabolism via Sphingosine 1-Phosphate and Its Role in Physiology, Pathology, and Nutrition 2015 , 127-138		
93	Opt2 mediates the exposure of phospholipids during cellular adaptation to altered lipid asymmetry. <i>Journal of Cell Science</i> , 2015 , 128, 61-9	5.3	9
92	Identification of the phytosphingosine metabolic pathway leading to odd-numbered fatty acids. <i>Nature Communications</i> , 2014 , 5, 5338	17.4	57
91	Lorenzoß oil inhibits ELOVL1 and lowers the level of sphingomyelin with a saturated very long-chain fatty acid. <i>Journal of Lipid Research</i> , 2014 , 55, 524-30	6.3	36
90	Sphingosine 1-phosphate is a key metabolite linking sphingolipids to glycerophospholipids. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2014 , 1841, 766-72	5	42
89	Metabolism of very long-chain Fatty acids: genes and pathophysiology. <i>Biomolecules and Therapeutics</i> , 2014 , 22, 83-92	4.2	137
88	Signaling events of the Rim101 pathway occur at the plasma membrane in a ubiquitination-dependent manner. <i>Molecular and Cellular Biology</i> , 2014 , 34, 3525-34	4.8	36
87	Dual functions of the trans-2-enoyl-CoA reductase TER in the sphingosine 1-phosphate metabolic pathway and in fatty acid elongation. <i>Journal of Biological Chemistry</i> , 2014 , 289, 24736-48	5.4	28
86	Integrin 9 on lymphatic endothelial cells regulates lymphocyte egress. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014 , 111, 3080-5	11.5	24
85	Two modes of regulation of the fatty acid elongase ELOVL6 by the 3-ketoacyl-CoA reductase KAR in the fatty acid elongation cycle. <i>PLoS ONE</i> , 2014 , 9, e101823	3.7	13
84	Phs1 and the synthesis of very long chain Fatty acids are required for ballistospore formation. <i>PLoS ONE</i> , 2014 , 9, e105147	3.7	6
83	Identification of acyl-CoA synthetases involved in the mammalian sphingosine 1-phosphate metabolic pathway. <i>Biochemical and Biophysical Research Communications</i> , 2013 , 442, 195-201	3.4	38
82	Identification of residues important for the catalysis, structure maintenance, and substrate specificity of yeast 3-hydroxyacyl-CoA dehydratase Phs1. <i>FEBS Letters</i> , 2013 , 587, 804-9	3.8	3
81	Unperverted synthesis of complex sphingolipids is essential for cell survival under nitrogen starvation. <i>Genes To Cells</i> , 2013 , 18, 650-9	2.3	17

80	Substrate specificity, plasma membrane localization, and lipid modification of the aldehyde dehydrogenase ALDH3B1. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2013 , 1831, 1395-401	5	21
79	Mutation for nonsyndromic mental retardation in the trans-2-enoyl-CoA reductase TER gene involved in fatty acid elongation impairs the enzyme activity and stability, leading to change in sphingolipid profile. <i>Journal of Biological Chemistry</i> , 2013 , 288, 36741-9	5.4	22
78	Impaired epidermal permeability barrier in mice lacking elovl1, the gene responsible for very-long-chain fatty acid production. <i>Molecular and Cellular Biology</i> , 2013 , 33, 2787-96	4.8	97
77	Congenital myopathy is caused by mutation of HACD1. Human Molecular Genetics, 2013, 22, 5229-36	5.6	40
76	Effects on vesicular transport pathways at the late endosome in cells with limited very long-chain fatty acids. <i>Journal of Lipid Research</i> , 2013 , 54, 831-842	6.3	23
75	Cooperative Synthesis of Ultra Long-Chain Fatty Acid and Ceramide during Keratinocyte Differentiation. <i>PLoS ONE</i> , 2013 , 8, e67317	3.7	35
74	Analysis of substrate specificity of human DHHC protein acyltransferases using a yeast expression system. <i>Molecular Biology of the Cell</i> , 2012 , 23, 4543-51	3.5	66
73	The Sjgren-Larsson syndrome gene encodes a hexadecenal dehydrogenase of the sphingosine 1-phosphate degradation pathway. <i>Molecular Cell</i> , 2012 , 46, 461-71	17.6	119
72	A shift in sphingolipid composition from C24 to C16 increases susceptibility to apoptosis in HeLa cells. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2012 , 1821, 1031-7	5	64
71	Guidelines for the use and interpretation of assays for monitoring autophagy. <i>Autophagy</i> , 2012 , 8, 445	-5 44 .2	2783
71 70		-5 4∂ .2 5·4	2783 46
•	Guidelines for the use and interpretation of assays for monitoring autophagy. <i>Autophagy</i> , 2012 , 8, 445 Membrane protein Rim21 plays a central role in sensing ambient pH in Saccharomyces cerevisiae.		
70	Guidelines for the use and interpretation of assays for monitoring autophagy. <i>Autophagy</i> , 2012 , 8, 445 Membrane protein Rim21 plays a central role in sensing ambient pH in Saccharomyces cerevisiae. <i>Journal of Biological Chemistry</i> , 2012 , 287, 38473-81 Palmitoylated calnexin is a key component of the ribosome-translocon complex. <i>EMBO Journal</i> ,	5.4	46
70 69	Guidelines for the use and interpretation of assays for monitoring autophagy. <i>Autophagy</i> , 2012 , 8, 445 Membrane protein Rim21 plays a central role in sensing ambient pH in Saccharomyces cerevisiae. <i>Journal of Biological Chemistry</i> , 2012 , 287, 38473-81 Palmitoylated calnexin is a key component of the ribosome-translocon complex. <i>EMBO Journal</i> , 2012 , 31, 1823-35 Degradation of long-chain base 1-phosphate (LCBP) in Arabidopsis: functional characterization of LCBP phosphatase involved in the dehydration stress response. <i>Journal of Plant Research</i> , 2012 ,	5.4	46 116
7° 69 68	Guidelines for the use and interpretation of assays for monitoring autophagy. <i>Autophagy</i> , 2012 , 8, 445 Membrane protein Rim21 plays a central role in sensing ambient pH in Saccharomyces cerevisiae. <i>Journal of Biological Chemistry</i> , 2012 , 287, 38473-81 Palmitoylated calnexin is a key component of the ribosome-translocon complex. <i>EMBO Journal</i> , 2012 , 31, 1823-35 Degradation of long-chain base 1-phosphate (LCBP) in Arabidopsis: functional characterization of LCBP phosphatase involved in the dehydration stress response. <i>Journal of Plant Research</i> , 2012 , 125, 439-49 Very long-chain fatty acids: elongation, physiology and related disorders. <i>Journal of Biochemistry</i> ,	5·4 13 2.6	46 116 25
7° 69 68 67	Guidelines for the use and interpretation of assays for monitoring autophagy. <i>Autophagy</i> , 2012 , 8, 445 Membrane protein Rim21 plays a central role in sensing ambient pH in Saccharomyces cerevisiae. <i>Journal of Biological Chemistry</i> , 2012 , 287, 38473-81 Palmitoylated calnexin is a key component of the ribosome-translocon complex. <i>EMBO Journal</i> , 2012 , 31, 1823-35 Degradation of long-chain base 1-phosphate (LCBP) in Arabidopsis: functional characterization of LCBP phosphatase involved in the dehydration stress response. <i>Journal of Plant Research</i> , 2012 , 125, 439-49 Very long-chain fatty acids: elongation, physiology and related disorders. <i>Journal of Biochemistry</i> , 2012 , 152, 387-95 Sphingolipids regulate the yeast high-osmolarity glycerol response pathway. <i>Molecular and Cellular</i>	5·4 13 2.6	46 116 25 227
70 69 68 67 66	Guidelines for the use and interpretation of assays for monitoring autophagy. <i>Autophagy</i> , 2012 , 8, 445 Membrane protein Rim21 plays a central role in sensing ambient pH in Saccharomyces cerevisiae. <i>Journal of Biological Chemistry</i> , 2012 , 287, 38473-81 Palmitoylated calnexin is a key component of the ribosome-translocon complex. <i>EMBO Journal</i> , 2012 , 31, 1823-35 Degradation of long-chain base 1-phosphate (LCBP) in Arabidopsis: functional characterization of LCBP phosphatase involved in the dehydration stress response. <i>Journal of Plant Research</i> , 2012 , 125, 439-49 Very long-chain fatty acids: elongation, physiology and related disorders. <i>Journal of Biochemistry</i> , 2012 , 152, 387-95 Sphingolipids regulate the yeast high-osmolarity glycerol response pathway. <i>Molecular and Cellular Biology</i> , 2012 , 32, 2861-70 Sphingolipid synthesis is involved in autophagy in Saccharomyces cerevisiae. <i>Biochemical and</i>	5.4 13 2.6 3.1 4.8	46 116 25 227 43

(2007-2010)

62	ELOVL1 production of C24 acyl-CoAs is linked to C24 sphingolipid synthesis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010 , 107, 18439-44	11.5	229
61	Hetero-oligomeric interactions of an ELOVL4 mutant protein: implications in the molecular mechanism of Stargardt-3 macular dystrophy. <i>Molecular Vision</i> , 2010 , 16, 2438-45	2.3	17
60	Lysophosphatidic acid 2 receptor-mediated supramolecular complex formation regulates its antiapoptotic effect. <i>Journal of Biological Chemistry</i> , 2009 , 284, 14558-71	5.4	57
59	Feedback inactivation of D-serine synthesis by NMDA receptor-elicited translocation of serine racemase to the membrane. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009 , 106, 7589-94	11.5	65
58	Palmitoylation of the sphingosine 1-phosphate receptor S1P is involved in its signaling functions and internalization. <i>Genes To Cells</i> , 2009 , 14, 911-23	2.3	13
57	Ceramide biosynthesis in keratinocyte and its role in skin function. <i>Biochimie</i> , 2009 , 91, 784-90	4.6	189
56	Characterization of four mammalian 3-hydroxyacyl-CoA dehydratases involved in very long-chain fatty acid synthesis. <i>FEBS Letters</i> , 2008 , 582, 2435-40	3.8	70
55	Production and release of sphingosine 1-phosphate and the phosphorylated form of the immunomodulator FTY720. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2008 , 1781, 496-502	5	69
54	A splicing isoform of LPP1, LPP1a, exhibits high phosphatase activity toward FTY720 phosphate. Biochemical and Biophysical Research Communications, 2008 , 375, 675-9	3.4	15
53	The Rim101 pathway is involved in Rsb1 expression induced by altered lipid asymmetry. <i>Molecular Biology of the Cell</i> , 2008 , 19, 1922-31	3.5	41
52	2-Hydroxy-ceramide synthesis by ceramide synthase family: enzymatic basis for the preference of FA chain length. <i>Journal of Lipid Research</i> , 2008 , 49, 2356-64	6.3	78
51	Membrane topology and essential amino acid residues of Phs1, a 3-hydroxyacyl-CoA dehydratase involved in very long-chain fatty acid elongation. <i>Journal of Biological Chemistry</i> , 2008 , 283, 11199-209	5.4	46
50	A sphingosine kinase activity assay using direct infusion electrospray ionization tandem mass spectrometry. <i>Analytical Biochemistry</i> , 2008 , 380, 35-40	3.1	7
49	Rapid trafficking of c-Src, a non-palmitoylated Src-family kinase, between the plasma membrane and late endosomes/lysosomes. <i>Experimental Cell Research</i> , 2007 , 313, 2651-66	4.2	68
48	Regulation of the transport and protein levels of the inositol phosphorylceramide mannosyltransferases Csg1 and Csh1 by the Ca2+-binding protein Csg2. <i>Journal of Biological Chemistry</i> , 2007 , 282, 8613-21	5.4	29
47	Intracellular trafficking pathway of yeast long-chain base kinase Lcb4, from its synthesis to its degradation. <i>Journal of Biological Chemistry</i> , 2007 , 282, 28485-28492	5.4	6
46	Metabolism and biological functions of two phosphorylated sphingolipids, sphingosine 1-phosphate and ceramide 1-phosphate. <i>Progress in Lipid Research</i> , 2007 , 46, 126-44	14.3	145
45	Lack of sphingosine 1-phosphate-degrading enzymes in erythrocytes. <i>Biochemical and Biophysical Research Communications</i> , 2007 , 357, 212-7	3.4	153

44	The immunomodulator FTY720 is phosphorylated and released from platelets. <i>European Journal of Pharmacology</i> , 2007 , 568, 106-11	5.3	30
43	Sphingosine kinase assay system with fluorescent detection in high performance liquid chromatography. <i>Archives of Pharmacal Research</i> , 2006 , 29, 1049-54	6.1	10
42	Sphingosine 1-phosphate is released from the cytosol of rat platelets in a carrier-mediated manner. <i>Journal of Lipid Research</i> , 2006 , 47, 614-21	6.3	136
41	Mouse sphingosine kinase isoforms SPHK1a and SPHK1b differ in enzymatic traits including stability, localization, modification, and oligomerization. <i>Journal of Biological Chemistry</i> , 2006 , 281, 4532	<u>2</u> 594	72
40	Intracellular localization and tissue-specific distribution of human and yeast DHHC cysteine-rich domain-containing proteins. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2006 , 1761, 474-83	5	297
39	Changes in S1P1 and S1P2 expression during embryonal development and primitive endoderm differentiation of F9 cells. <i>Biochemical and Biophysical Research Communications</i> , 2006 , 344, 852-8	3.4	6
38	Lipid asymmetry of the eukaryotic plasma membrane: functions and related enzymes. <i>Biological and Pharmaceutical Bulletin</i> , 2006 , 29, 1542-6	2.3	94
37	Rescue of cell growth by sphingosine with disruption of lipid microdomain formation in Saccharomyces cerevisiae deficient in sphingolipid biosynthesis. <i>Biochemical Journal</i> , 2006 , 394, 237-42	3.8	24
36	LASS3 (longevity assurance homologue 3) is a mainly testis-specific (dihydro)ceramide synthase with relatively broad substrate specificity. <i>Biochemical Journal</i> , 2006 , 398, 531-8	3.8	135
35	Synthesis, Metabolism, and Trans-Bilayer Movement of Long-Chain Base 2006 , 95-106		1
34	Sphingolipid-to-glycerophospholipid conversion in SPL-null cells implies the existence of an alternative isozyme. <i>Biochemical and Biophysical Research Communications</i> , 2005 , 329, 474-9	3.4	9
33	Mammalian Lass6 and its related family members regulate synthesis of specific ceramides. <i>Biochemical Journal</i> , 2005 , 390, 263-71	3.8	288
32	Products by the sphingosine kinase/sphingosine 1-phosphate (S1P) lyase pathway but not S1P stimulate mitogenesis. <i>Genes To Cells</i> , 2005 , 10, 605-15	2.3	47
31	Phosphorylation by Pho85 cyclin-dependent kinase acts as a signal for the down-regulation of the yeast sphingoid long-chain base kinase Lcb4 during the stationary phase. <i>Journal of Biological Chemistry</i> , 2005 , 280, 6520-7	5.4	24
30	Long-chain base kinase Lcb4 Is anchored to the membrane through its palmitoylation by Akr1. <i>Molecular and Cellular Biology</i> , 2005 , 25, 9189-97	4.8	38
29	Regulation of the sphingoid long-chain base kinase Lcb4p by ergosterol and heme: studies in phytosphingosine-resistant mutants. <i>Journal of Biological Chemistry</i> , 2005 , 280, 36674-82	5.4	16
28	Cross talk between sphingolipids and glycerophospholipids in the establishment of plasma membrane asymmetry. <i>Molecular Biology of the Cell</i> , 2004 , 15, 4949-59	3.5	78
27	FVT-1 is a mammalian 3-ketodihydrosphingosine reductase with an active site that faces the cytosolic side of the endoplasmic reticulum membrane. <i>Journal of Biological Chemistry</i> , 2004 , 279, 4924	-5-4s	64

26	Asp177 in C4 domain of mouse sphingosine kinase 1a is important for the sphingosine recognition. <i>FEBS Letters</i> , 2004 , 578, 106-10	3.8	39
25	Sphingosine-1-phosphate lyase SPL is an endoplasmic reticulum-resident, integral membrane protein with the pyridoxal 5Rphosphate binding domain exposed to the cytosol. <i>Biochemical and Biophysical Research Communications</i> , 2004 , 325, 338-43	3.4	125
24	Identification of the human sphingolipid C4-hydroxylase, hDES2, and its up-regulation during keratinocyte differentiation. <i>FEBS Letters</i> , 2004 , 563, 93-7	3.8	76
23	Identification and characterization of a novel human sphingosine-1-phosphate phosphohydrolase, hSPP2. <i>Journal of Biological Chemistry</i> , 2003 , 278, 1268-72	5.4	144
22	Transmembrane topology of sphingoid long-chain base-1-phosphate phosphatase, Lcb3p. <i>Genes To Cells</i> , 2003 , 8, 525-35	2.3	35
21	Csg1p and newly identified Csh1p function in mannosylinositol phosphorylceramide synthesis by interacting with Csg2p. <i>Journal of Biological Chemistry</i> , 2003 , 278, 45049-55	5.4	70
20	Distribution of sphingosine kinase activity in mouse tissues: contribution of SPHK1. <i>Biochemical and Biophysical Research Communications</i> , 2003 , 309, 155-60	3.4	103
19	Sphingosine-1-phosphate lyase is involved in the differentiation of F9 embryonal carcinoma cells to primitive endoderm. <i>Journal of Biological Chemistry</i> , 2003 , 278, 14578-85	5.4	69
18	Identification and characterization of a Saccharomyces cerevisiae gene, RSB1, involved in sphingoid long-chain base release. <i>Journal of Biological Chemistry</i> , 2002 , 277, 30048-54	5.4	76
17	Beclin-phosphatidylinositol 3-kinase complex functions at the trans-Golgi network. <i>EMBO Reports</i> , 2001 , 2, 330-5	6.5	716
16	Autophagosome requires specific early Sec proteins for its formation and NSF/SNARE for vacuolar fusion. <i>Molecular Biology of the Cell</i> , 2001 , 12, 3690-702	3.5	296
15	Revisiting the lysogenization control of bacteriophage lambda. Identification and characterization of a new host component, HflD. <i>Journal of Biological Chemistry</i> , 2001 , 276, 13695-700	5.4	26
14	Two distinct Vps34 phosphatidylinositol 3-kinase complexes function in autophagy and carboxypeptidase Y sorting in Saccharomyces cerevisiae. <i>Journal of Cell Biology</i> , 2001 , 152, 519-30	7-3	811
13	Three-dimensional structure of phosphoenolpyruvate carboxylase: a proposed mechanism for allosteric inhibition. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1999 , 96, 823-8	11.5	125
12	Dislocation of membrane proteins in FtsH-mediated proteolysis. <i>EMBO Journal</i> , 1999 , 18, 2970-81	13	95
11	Polypeptide binding of Escherichia coli FtsH (HflB). <i>Molecular Microbiology</i> , 1998 , 28, 803-12	4.1	32
10	Different pathways for protein degradation by the FtsH/HflKC membrane-embedded protease complex: an implication from the interference by a mutant form of a new substrate protein, YccA. <i>Journal of Molecular Biology</i> , 1998 , 279, 175-88	6.5	100
9	Roles of the periplasmic domain of Escherichia coli FtsH (HflB) in protein interactions and activity modulation. <i>Journal of Biological Chemistry</i> , 1998 , 273, 22326-33	5.4	37

8	Translocation, folding, and stability of the HflKC complex with signal anchor topogenic sequences. Journal of Biological Chemistry, 1998 , 273, 29770-5	5.4	29
7	Host regulation of lysogenic decision in bacteriophage lambda: transmembrane modulation of FtsH (HflB), the cII degrading protease, by HflKC (HflA). <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1997 , 94, 5544-9	11.5	94
6	Subunit a of proton ATPase F0 sector is a substrate of the FtsH protease in Escherichia coli. <i>FEBS Letters</i> , 1996 , 399, 26-8	3.8	104
5	FtsH (HflB) is an ATP-dependent protease selectively acting on SecY and some other membrane proteins. <i>Journal of Biological Chemistry</i> , 1996 , 271, 31196-201	5.4	117
4	Phosphoenolpyruvate carboxylase: Alteration of catalytic and regulatory properties by site-directed mutagenesis and isolation of the gene from an extreme thermophile. <i>Energy Conversion and Management</i> , 1995 , 36, 751-754	10.6	
3	Product of a new gene, syd, functionally interacts with SecY when overproduced in Escherichia coli. <i>Journal of Biological Chemistry</i> , 1995 , 270, 5519-26	5.4	55
2	Intracellular stability of alpha fragments of beta-galactosidase: effects of amino-terminally fused polypeptides. <i>Biochemical and Biophysical Research Communications</i> , 1995 , 215, 452-8	3.4	9
1	FtsH is required for proteolytic elimination of uncomplexed forms of SecY, an essential protein translocase subunit. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1995 , 92, 4532-6	11.5	219