

# Janet R Kumita

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/4257083/janet-r-kumita-publications-by-year.pdf>

**Version:** 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

72  
papers

3,538  
citations

33  
h-index

59  
g-index

77  
ext. papers

4,258  
ext. citations

7.2  
avg, IF

4.91  
L-index

#	Paper	IF	Citations
72	Comparative Studies in the A30P and A53T $\beta$ Synuclein Strains to Investigate the Molecular Origins of Parkinson's Disease. <i>Frontiers in Cell and Developmental Biology</i> , <b>2021</b> , 9, 552549	5.7	5
71	Characterisation of the structural, dynamic and aggregation properties of the W64R amyloidogenic variant of human lysozyme. <i>Biophysical Chemistry</i> , <b>2021</b> , 271, 106563	3.5	1
70	Squalamine and Its Derivatives Modulate the Aggregation of Amyloid- $\beta$ and $\beta$ Synuclein and Suppress the Toxicity of Their Oligomers. <i>Frontiers in Neuroscience</i> , <b>2021</b> , 15, 680026	5.1	11
69	Different Folding States from the Same Protein Sequence Determine Reversible vs Irreversible Amyloid Fate. <i>Journal of the American Chemical Society</i> , <b>2021</b> , 143, 11473-11481	16.4	7
68	Engineering mono- and multi-valent inhibitors on a modular scaffold. <i>Chemical Science</i> , <b>2021</b> , 12, 880-895	5.4	4
67	Exogenous misfolded protein oligomers can cross the intestinal barrier and cause a disease phenotype in <i>C. elegans</i> . <i>Scientific Reports</i> , <b>2021</b> , 11, 14391	4.9	1
66	A dopamine metabolite stabilizes neurotoxic amyloid- $\beta$ oligomers. <i>Communications Biology</i> , <b>2021</b> , 4, 19	6.7	6
65	The Influence of Pathogenic Mutations in $\beta$ Synuclein on Biophysical and Structural Characteristics of Amyloid Fibrils. <i>ACS Nano</i> , <b>2020</b> , 14, 5213-5222	16.7	24
64	Rationally Designed Antibodies as Research Tools to Study the Structure-Toxicity Relationship of Amyloid- $\beta$ Oligomers. <i>International Journal of Molecular Sciences</i> , <b>2020</b> , 21,	6.3	7
63	Serum amyloid P component promotes formation of distinct aggregated lysozyme morphologies and reduces toxicity in <i>Drosophila</i> flies expressing F57I lysozyme. <i>PLoS ONE</i> , <b>2020</b> , 15, e0227227	3.7	1
62	Cholesterol-rich naked mole-rat brain lipid membranes are susceptible to amyloid beta-induced damage. <i>Aging</i> , <b>2020</b> , 12, 22266-22290	5.6	6
61	Mapping pathogenic processes contributing to neurodegeneration in <i>Drosophila</i> models of Alzheimer's disease. <i>FEBS Open Bio</i> , <b>2020</b> , 10, 338-350	2.7	3
60	Rapid Structural, Kinetic, and Immunochemical Analysis of Alpha-Synuclein Oligomers in Solution. <i>Nano Letters</i> , <b>2020</b> , 20, 8163-8169	11.5	5
59	Probing the unfolded protein response in long-lived naked mole-rats. <i>Biochemical and Biophysical Research Communications</i> , <b>2020</b> , 529, 1151-1157	3.4	3
58	A prion-like domain in ELF3 functions as a thermosensor in <i>Arabidopsis</i> . <i>Nature</i> , <b>2020</b> , 585, 256-260	50.4	128
57	Trodusquemine displaces protein misfolded oligomers from cell membranes and abrogates their cytotoxicity through a generic mechanism. <i>Communications Biology</i> , <b>2020</b> , 3, 435	6.7	23
56	Using Tetracysteine-Tagged TDP-43 with a Biarsenical Dye To Monitor Real-Time Trafficking in a Cell Model of Amyotrophic Lateral Sclerosis. <i>Biochemistry</i> , <b>2019</b> , 58, 4086-4095	3.2	3

55	Chemical and mechanistic analysis of photodynamic inhibition of Alzheimer's Amyloid aggregation. <i>Chemical Communications</i> , <b>2019</b> , 55, 1152-1155	5.8	11
54	Defining Synuclein species responsible for Parkinson's disease phenotypes in mice. <i>Journal of Biological Chemistry</i> , <b>2019</b> , 294, 10392-10406	5.4	55
53	Secondary nucleation and elongation occur at different sites on Alzheimer's amyloid- $\beta$ aggregates. <i>Science Advances</i> , <b>2019</b> , 5, eaau3112	14.3	74
52	Fabrication and Characterization of Reconstituted Silk Microgels for the Storage and Release of Small Molecules. <i>Macromolecular Rapid Communications</i> , <b>2019</b> , 40, e1800898	4.8	23
51	Human pregnancy zone protein stabilizes misfolded proteins including preeclampsia- and Alzheimer's-associated amyloid beta peptide. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2019</b> , 116, 6101-6110	11.5	32
50	Homage to Chris Dobson. <i>Frontiers in Molecular Biosciences</i> , <b>2019</b> , 6, 137	5.6	
49	Trodusquemine enhances A $\beta$ aggregation but suppresses its toxicity by displacing oligomers from cell membranes. <i>Nature Communications</i> , <b>2019</b> , 10, 225	17.4	69
48	Cholesterol catalyses A $\beta$ 2 aggregation through a heterogeneous nucleation pathway in the presence of lipid membranes. <i>Nature Chemistry</i> , <b>2018</b> , 10, 673-683	17.6	126
47	Single-Molecule Characterization of the Interactions between Extracellular Chaperones and Toxic Synuclein Oligomers. <i>Cell Reports</i> , <b>2018</b> , 23, 3492-3500	10.6	42
46	Multistep Inhibition of Synuclein Aggregation and Toxicity in Vitro and in Vivo by Trodusquemine. <i>ACS Chemical Biology</i> , <b>2018</b> , 13, 2308-2319	4.9	52
45	Inhibition of Synuclein Fibril Elongation by Hsp70 Is Governed by a Kinetic Binding Competition between Synuclein Species. <i>Biochemistry</i> , <b>2017</b> , 56, 1177-1180	3.2	45
44	Flow cytometric measurement of the cellular propagation of TDP-43 aggregation. <i>Prion</i> , <b>2017</b> , 11, 195-204	3.3	16
43	Amyloid-like Fibrils from an Helical Transmembrane Protein. <i>Biochemistry</i> , <b>2017</b> , 56, 3225-3233	3.2	14
42	Systematic development of small molecules to inhibit specific microscopic steps of A $\beta$ 2 aggregation in Alzheimer's disease. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2017</b> , 114, E200-E208	11.5	134
41	Clusterin protects neurons against intracellular proteotoxicity. <i>Acta Neuropathologica Communications</i> , <b>2017</b> , 5, 81	7.3	33
40	The relevance of contact-independent cell-to-cell transfer of TDP-43 and SOD1 in amyotrophic lateral sclerosis. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , <b>2017</b> , 1863, 2762-2771	6.9	14
39	Application of Lysine-specific Labeling to Detect Transient Interactions Present During Human Lysozyme Amyloid Fibril Formation. <i>Scientific Reports</i> , <b>2017</b> , 7, 15018	4.9	5
38	The Significance of the Location of Mutations for the Native-State Dynamics of Human Lysozyme. <i>Biophysical Journal</i> , <b>2016</b> , 111, 2358-2367	2.9	12

37	Structure and dynamics of the integrin LFA-1 I-domain in the inactive state underlie its inside-out/outside-in signaling and allosteric mechanisms. <i>Structure</i> , <b>2015</b> , 23, 745-53	5.2	15
36	The influence of novel gemini surfactants containing cycloalkyl side-chains on the structural phases of DNA in solution. <i>Colloids and Surfaces B: Biointerfaces</i> , <b>2015</b> , 131, 83-92	6	15
35	Solvent exposure of Tyr10 as a probe of structural differences between monomeric and aggregated forms of the amyloid- $\beta$ peptide. <i>Biochemical and Biophysical Research Communications</i> , <b>2015</b> , 468, 696-701	3.4	8
34	Alpha-2-Macroglobulin Is Acutely Sensitive to Freezing and Lyophilization: Implications for Structural and Functional Studies. <i>PLoS ONE</i> , <b>2015</b> , 10, e0130036	3.7	7
33	NMR characterization of the conformational fluctuations of the human lymphocyte function-associated antigen-1 I-domain. <i>Protein Science</i> , <b>2014</b> , 23, 1596-606	6.3	7
32	Hypochlorite-induced structural modifications enhance the chaperone activity of human $\alpha$ -macroglobulin. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2014</b> , 111, E2081-90	11.5	49
31	Intercellular propagated misfolding of wild-type Cu/Zn superoxide dismutase occurs via exosome-dependent and -independent mechanisms. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2014</b> , 111, 3620-5	11.5	293
30	Protease-activated alpha-2-macroglobulin can inhibit amyloid formation via two distinct mechanisms. <i>FEBS Letters</i> , <b>2013</b> , 587, 398-403	3.8	32
29	Protein amyloids develop an intrinsic fluorescence signature during aggregation. <i>Analyst, The</i> , <b>2013</b> , 138, 2156-62	5	140
28	A nanobody binding to non-amyloidogenic regions of the protein human lysozyme enhances partial unfolding but inhibits amyloid fibril formation. <i>Journal of Physical Chemistry B</i> , <b>2013</b> , 117, 13245-13258	3.4	33
27	Analysis of the native structure, stability and aggregation of biotinylated human lysozyme. <i>PLoS ONE</i> , <b>2012</b> , 7, e50192	3.7	20
26	Disease-related amyloidogenic variants of human lysozyme trigger the unfolded protein response and disturb eye development in <i>Drosophila melanogaster</i> . <i>FASEB Journal</i> , <b>2012</b> , 26, 192-202	0.9	15
25	Native-state stability determines the extent of degradation relative to secretion of protein variants from <i>Pichia pastoris</i> . <i>PLoS ONE</i> , <b>2011</b> , 6, e22692	3.7	41
24	Population of nonnative states of lysozyme variants drives amyloid fibril formation. <i>Journal of the American Chemical Society</i> , <b>2011</b> , 133, 7737-7743	16.4	67
23	A FRET sensor for non-invasive imaging of amyloid formation in vivo. <i>ChemPhysChem</i> , <b>2011</b> , 12, 673-680	3.2	76
22	Disulfide Bonds Reduce the Toxicity of the Amyloid Fibrils Formed by an Extracellular Protein. <i>Angewandte Chemie</i> , <b>2011</b> , 123, 7186-7189	3.6	6
21	Disulfide bonds reduce the toxicity of the amyloid fibrils formed by an extracellular protein. <i>Angewandte Chemie - International Edition</i> , <b>2011</b> , 50, 7048-51	16.4	48
20	A non-natural variant of human lysozyme (I59T) mimics the in vitro behaviour of the I56T variant that is responsible for a form of familial amyloidosis. <i>Protein Engineering, Design and Selection</i> , <b>2010</b> , 23, 499-506	1.9	17

19	Local cooperativity in an amyloidogenic state of human lysozyme observed at atomic resolution. <i>Journal of the American Chemical Society</i> , <b>2010</b> , 132, 15580-8	16.4	49
18	ANS binding reveals common features of cytotoxic amyloid species. <i>ACS Chemical Biology</i> , <b>2010</b> , 5, 735-40	4.9	291
17	Towards multiparametric fluorescent imaging of amyloid formation: studies of a YFP model of alpha-synuclein aggregation. <i>Journal of Molecular Biology</i> , <b>2010</b> , 395, 627-42	6.5	62
16	The non-core regions of human lysozyme amyloid fibrils influence cytotoxicity. <i>Journal of Molecular Biology</i> , <b>2010</b> , 402, 783-96	6.5	84
15	Protein Chemistry of Amyloid Fibrils and Chaperones: Implications for Amyloid Formation and Disease. <i>Current Chemical Biology</i> , <b>2010</b> , 4, 89-98	0.4	7
14	alpha2-Macroglobulin and haptoglobin suppress amyloid formation by interacting with prefibrillar protein species. <i>Journal of Biological Chemistry</i> , <b>2009</b> , 284, 4246-54	5.4	72
13	Engineering a camelid antibody fragment that binds to the active site of human lysozyme and inhibits its conversion into amyloid fibrils. <i>Biochemistry</i> , <b>2008</b> , 47, 11041-54	3.2	58
12	Molecular determinants of the aggregation behavior of alpha- and beta-synuclein. <i>Protein Science</i> , <b>2008</b> , 17, 887-98	6.3	84
11	The extracellular chaperone clusterin influences amyloid formation and toxicity by interacting with prefibrillar structures. <i>FASEB Journal</i> , <b>2007</b> , 21, 2312-22	0.9	237
10	The extracellular chaperone clusterin potently inhibits human lysozyme amyloid formation by interacting with prefibrillar species. <i>Journal of Molecular Biology</i> , <b>2007</b> , 369, 157-67	6.5	74
9	Normal and aberrant biological self-assembly: Insights from studies of human lysozyme and its amyloidogenic variants. <i>Accounts of Chemical Research</i> , <b>2006</b> , 39, 603-10	24.3	84
8	Impact of the native-state stability of human lysozyme variants on protein secretion by <i>Pichia pastoris</i> . <i>FEBS Journal</i> , <b>2006</b> , 273, 711-20	5.7	43
7	Alpha-helix formation in a photoswitchable peptide tracked from picoseconds to microseconds by time-resolved IR spectroscopy. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2005</b> , 102, 2379-84	11.5	174
6	Rationalising lysozyme amyloidosis: insights from the structure and solution dynamics of T70N lysozyme. <i>Journal of Molecular Biology</i> , <b>2005</b> , 352, 823-36	6.5	39
5	A water-soluble azobenzene cross-linker for photocontrol of peptide conformation. <i>Bioconjugate Chemistry</i> , <b>2003</b> , 14, 824-9	6.3	69
4	The kinetics of helix unfolding of an azobenzene cross-linked peptide probed by nanosecond time-resolved optical rotatory dispersion. <i>Journal of the American Chemical Society</i> , <b>2003</b> , 125, 12443-9	16.4	60
3	Achieving photo-control of protein conformation and activity: producing a photo-controlled leucine zipper. <i>Faraday Discussions</i> , <b>2003</b> , 122, 89-103; discussion 171-90	3.6	24
2	Using an azobenzene cross-linker to either increase or decrease peptide helix content upon trans-to-cis photoisomerization. <i>Chemistry and Biology</i> , <b>2002</b> , 9, 391-7		130

1 A cysteine-free firefly luciferase retains luminescence activity. *Biochemical and Biophysical Research Communications*, **2000**, 267, 394-7

3.4 13