

Chengwen Sun

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/425904/chengwen-sun-publications-by-year.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

59
papers

2,173
citations

27
h-index

45
g-index

59
ext. papers

2,350
ext. citations

5.7
avg, IF

4.63
L-index

#	Paper	IF	Citations
59	Apelin Does Not Impair Coronary Artery Relaxation Mediated by Nitric Oxide-Induced Activation of BK Channels. <i>Frontiers in Pharmacology</i> , 2021 , 12, 679005	5.6	2
58	Role of PI3-Kinase in Angiotensin II-Induced Cardiac Hypertrophy: Class I Versus Class III. <i>Frontiers in Pharmacology</i> , 2021 , 12, 608523	5.6	1
57	Apelin inhibits an endothelium-derived hyperpolarizing factor-like pathway in rat cerebral arteries. <i>Peptides</i> , 2020 , 132, 170350	3.8	3
56	Biodistribution of TAT or QLPVM coupled to receptor targeted liposomes for delivery of anticancer therapeutics to brain in vitro and in vivo. <i>Nanomedicine: Nanotechnology, Biology, and Medicine</i> , 2020 , 23, 102112	6	23
55	Dual functionalized liposomes for efficient co-delivery of anti-cancer chemotherapeutics for the treatment of glioblastoma. <i>Journal of Controlled Release</i> , 2019 , 307, 247-260	11.7	61
54	Apelin Reduces Nitric Oxide-Induced Relaxation of Cerebral Arteries by Inhibiting Activation of Large-Conductance, Calcium-Activated K Channels. <i>Journal of Cardiovascular Pharmacology</i> , 2018 , 71, 223-232	3.1	13
53	Activation of Large Conductance, Calcium-Activated Potassium Channels by Nitric Oxide Mediates Apelin-Induced Relaxation of Isolated Rat Coronary Arteries. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2018 , 366, 265-273	4.7	16
52	Angiotensin-(1-7) attenuates angiotensin II-induced cardiac hypertrophy via a Sirt3-dependent mechanism. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2017 , 312, H980-H991	5.2	38
51	Adrenergic Receptor-stimulated Cardiac Myocyte Apoptosis: Role of Cytochrome P450 Hydroxylase. <i>Journal of Cardiovascular Pharmacology</i> , 2017 , 70, 94-101	3.1	9
50	20-Hydroxyeicosatetraenoic Acid Is a Key Mediator of Angiotensin II-induced Apoptosis in Cardiac Myocytes. <i>Journal of Cardiovascular Pharmacology</i> , 2015 , 66, 86-95	3.1	18
49	High salt-diet reduces SLC14A1 gene expression in the choroid plexus of Dahl salt sensitive rats. <i>Biochemical and Biophysical Research Communications</i> , 2015 , 461, 254-9	3.4	7
48	Influence of short-chain cell-penetrating peptides on transport of doxorubicin encapsulating receptor-targeted liposomes across brain endothelial barrier. <i>Pharmaceutical Research</i> , 2014 , 31, 1194-209	4.5	51
47	Needleless emulsion electrospinning for scalable fabrication of core-shell nanofibers. <i>Journal of Applied Polymer Science</i> , 2014 , 131, n/a-n/a	2.9	18
46	Maternal nutrient restriction during pregnancy impairs an endothelium-derived hyperpolarizing factor-like pathway in sheep fetal coronary arteries. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2014 , 307, H134-42	5.2	18
45	Cell penetrating peptide tethered bi-ligand liposomes for delivery to brain in vivo: Biodistribution and transfection. <i>Journal of Controlled Release</i> , 2013 , 167, 1-10	11.7	127
44	In vitro assessment of clevidipine using the profilin1 hypertensive mouse model. <i>Pharmaceuticals</i> , 2013 , 6, 623-33	5.2	1
43	GABAB receptor gene transfer into the nucleus tractus solitarii induces chronic blood pressure elevation in normotensive rats. <i>Circulation Journal</i> , 2013 , 77, 2558-66	2.9	6

42	20-Hydroxyeicosatetraenoic acid mediates isolated heart ischemia/reperfusion injury by increasing NADPH oxidase-derived reactive oxygen species production. <i>Circulation Journal</i> , 2013 , 77, 1807-16	2.9	29
41	20-Hydroxyeicosatetraenoic acid contributes to the inhibition of K ⁺ channel activity and vasoconstrictor response to angiotensin II in rat renal microvessels. <i>PLoS ONE</i> , 2013 , 8, e82482	3.7	44
40	Apelin-13 inhibits large-conductance Ca ²⁺ -activated K ⁺ channels in cerebral artery smooth muscle cells via a PI3-kinase dependent mechanism. <i>PLoS ONE</i> , 2013 , 8, e83051	3.7	27
39	Lentil polyphenol extract prevents angiotensin II-induced hypertension, vascular remodelling and perivascular fibrosis. <i>Food and Function</i> , 2012 , 3, 127-33	6.1	14
38	Invertible micellar polymer assemblies for delivery of poorly water-soluble drugs. <i>Biomacromolecules</i> , 2012 , 13, 2537-45	6.9	37
37	Grafting of cell-penetrating peptide to receptor-targeted liposomes improves their transfection efficiency and transport across blood-brain barrier model. <i>Journal of Pharmaceutical Sciences</i> , 2012 , 101, 2468-78	3.9	56
36	Melatonin inhibits nitric oxide signaling by increasing PDE5 phosphorylation in coronary arteries. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2012 , 303, H1418-25	5.2	19
35	Angiotensin-(1-7) attenuates the chronotropic response to angiotensin II via stimulation of PTEN in the spontaneously hypertensive rat neurons. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2012 , 302, H1116-22	5.2	12
34	20-Hydroxyeicosatetraenoic acid induces apoptosis in neonatal rat cardiomyocytes through mitochondrial-dependent pathways. <i>Journal of Cardiovascular Pharmacology</i> , 2011 , 57, 294-301	3.1	94
33	Pressor effect of apelin-13 in the rostral ventrolateral medulla: role of NAD(P)H oxidase-derived superoxide. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2011 , 336, 372-80	4.7	22
32	MT2 receptors mediate the inhibitory effects of melatonin on nitric oxide-induced relaxation of porcine isolated coronary arteries. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2011 , 336, 127-33	4.7	35
31	20-HETE increases NADPH oxidase-derived ROS production and stimulates the L-type Ca ²⁺ channel via a PKC-dependent mechanism in cardiomyocytes. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2010 , 299, H1109-17	5.2	51
30	Morton lentil extract attenuated angiotensin II-induced cardiomyocyte hypertrophy via inhibition of intracellular reactive oxygen species levels in vitro. <i>Journal of Agricultural and Food Chemistry</i> , 2010 , 58, 10382-8	5.7	12
29	Angiotensin II enhances GABA(B) receptor-mediated responses and expression in nucleus tractus solitarii of rats. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2009 , 297, H1837-44	5.2	19
28	Shift to an involvement of phosphatidylinositol 3-kinase in angiotensin II actions on nucleus tractus solitarii neurons of the spontaneously hypertensive rat. <i>Circulation Research</i> , 2009 , 105, 1248-55	15.7	26
27	Apelin gene transfer into the rostral ventrolateral medulla induces chronic blood pressure elevation in normotensive rats. <i>Circulation Research</i> , 2009 , 104, 1421-8	15.7	66
26	Characterization of novel radicals from COX-catalyzed arachidonic acid peroxidation. <i>Free Radical Biology and Medicine</i> , 2009 , 47, 568-76	7.8	24
25	Endothelin-1 induces intracellular [Ca ²⁺] increase via Ca ²⁺ influx through the L-type Ca ²⁺ channel, Ca ²⁺ -induced Ca ²⁺ release and a pathway involving ET A receptors, PKC, PKA and AT1 receptors in cardiomyocytes. <i>Science in China Series C: Life Sciences</i> , 2009 , 52, 360-70		9

24	Mechanical hyperalgesia is attenuated by local administration of octreotide in pristane-induced arthritis in Dark-Agouti rats. <i>Life Sciences</i> , 2008 , 83, 732-8	6.8	8
23	Endothelin-1 regulates cardiac L-type calcium channels via NAD(P)H oxidase-derived superoxide. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2008 , 326, 732-8	4.7	53
22	Angiotensin II increases GABAB receptor expression in nucleus tractus solitarii of rats. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2008 , 294, H2712-20	5.2	33
21	Overexpression of angiotensin-converting enzyme 2 in the rostral ventrolateral medulla causes long-term decrease in blood pressure in the spontaneously hypertensive rats. <i>Hypertension</i> , 2007 , 49, 926-31	8.5	148
20	Lack of macrophage migration inhibitory factor regulation is linked to the increased chronotropic action of angiotensin II in SHR neurons. <i>Hypertension</i> , 2007 , 49, 528-34	8.5	13
19	Interaction between Ang II and GABA systems in NTS: Central resetting of blood pressure regulation. <i>FASEB Journal</i> , 2007 , 21, A879	0.9	
18	Macrophage migration inhibitory factor increases neuronal delayed rectifier K ⁺ current. <i>Journal of Neurophysiology</i> , 2006 , 95, 1042-8	3.2	13
17	Novel mechanism of brain soluble epoxide hydrolase-mediated blood pressure regulation in the spontaneously hypertensive rat. <i>FASEB Journal</i> , 2005 , 19, 626-8	0.9	50
16	NAD(P)H oxidase inhibition attenuates neuronal chronotropic actions of angiotensin II. <i>Circulation Research</i> , 2005 , 96, 659-66	15.7	95
15	Macrophage migration inhibitory factor: an intracellular inhibitor of angiotensin II-induced increases in neuronal activity. <i>Journal of Neuroscience</i> , 2004 , 24, 9944-52	6.6	49
14	20-hydroxyeicosatetraenoic acid (20-HETE): structural determinants for renal vasoconstriction. <i>Bioorganic and Medicinal Chemistry</i> , 2003 , 11, 2803-21	3.4	27
13	Modulation of delayed rectifier potassium current by angiotensin II in CATH.a cells. <i>Biochemical and Biophysical Research Communications</i> , 2003 , 310, 710-4	3.4	20
12	Transduction of a functional domain of the AT1 receptor in neurons by HIV-Tat PTD. <i>Hypertension</i> , 2003 , 41, 751-6	8.5	15
11	PI3-kinase inhibitors abolish the enhanced chronotropic effects of angiotensin II in spontaneously hypertensive rat brain neurons. <i>Journal of Neurophysiology</i> , 2003 , 90, 3155-60	3.2	19
10	Mechanism of cGMP contribution to the vasodilator response to NO in rat middle cerebral arteries. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2002 , 282, H1724-31	5.2	30
9	Chronotropic action of angiotensin II in neurons via protein kinase C and CaMKII. <i>Hypertension</i> , 2002 , 39, 562-6	8.5	43
8	Hypertension-linked decrease in the expression of brain gamma-adducin. <i>Circulation Research</i> , 2002 , 91, 633-9	15.7	18
7	Renal and cardiovascular actions of 20-hydroxyeicosatetraenoic acid and epoxyeicosatrienoic acids. <i>Clinical and Experimental Pharmacology and Physiology</i> , 2000 , 27, 855-65	3	101

6	Role of cGMP versus 20-HETE in the vasodilator response to nitric oxide in rat cerebral arteries. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2000 , 279, H339-50	5.2	69
5	Altered renal hemodynamics and impaired myogenic responses in the fawn-hooded rat. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 1999 , 276, R855-63	3.2	48
4	Role of tyrosine kinase and PKC in the vasoconstrictor response to 20-HETE in renal arterioles. <i>Hypertension</i> , 1999 , 33, 414-8	8.5	105
3	Lovastatin reduces renal vascular reactivity in spontaneously hypertensive rats. <i>American Journal of Hypertension</i> , 1998 , 11, 1222-31	2.3	19
2	Nitric oxide-20-hydroxyeicosatetraenoic acid interaction in the regulation of K ⁺ channel activity and vascular tone in renal arterioles. <i>Circulation Research</i> , 1998 , 83, 1069-79	15.7	146
1	Contribution of 20-HETE to the vasodilator actions of nitric oxide in renal arteries. <i>American Journal of Physiology - Renal Physiology</i> , 1998 , 275, F370-8	4.3	43