

Marinela Maria Dirtu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/426677/publications.pdf>

Version: 2024-02-01

40
papers

1,288
citations

304743

22
h-index

345221

36
g-index

43
all docs

43
docs citations

43
times ranked

1224
citing authors

#	ARTICLE	IF	CITATIONS
1	One-dimensional Looped Chain and Two-dimensional Square Grid Coordination Polymers: Encapsulation of Bis(1,2,4-Triazole)-trans-cyclohexane into the Voids. European Journal of Inorganic Chemistry, 2019, 2019, 543-543.	2.0	0
2	One-Dimensional Looped Chain and Two-Dimensional Square Grid Coordination Polymers: Encapsulation of Bis(1,2,4-Triazole)-trans -cyclohexane into the Voids. European Journal of Inorganic Chemistry, 2019, 2019, 585-591.	2.0	14
3	Selecting the spin crossover profile with controlled crystallization of mononuclear Fe(_{scp}) _{iii} (_{scp}) polymorphs. Dalton Transactions, 2018, 47, 7013-7019.	3.3	9
4	A versatile iron(_{scp}) _{ii} (_{scp})-based colorimetric sensor for the vapor-phase detection of alcohols and toxic gases. Journal of Materials Chemistry C, 2018, 6, 3895-3900.	5.5	45
5	Quantitative Contact Pressure Sensor Based on Spin Crossover Mechanism for Civil Security Applications. Journal of Physical Chemistry C, 2018, 122, 7597-7604.	3.1	58
6	Spin State Crossover, Vibrational, Computational, and Structural Studies of Fe ^{II} -isopropyl-H ₄ Tetrazole Derivatives. European Journal of Inorganic Chemistry, 2018, 2018, 394-413.	2.0	7
7	Spin-Crossover in an Exfoliated 2D Coordination Polymer and Its Implementation in Thermochromic Films. ACS Applied Nano Materials, 2018, 1, 2662-2668.	5.0	22
8	57Fe Mössbauer spectroscopy study of a 2D spin transition coordination polymer built from a tris-1R-tetrazole ligand. Hyperfine Interactions, 2017, 238, 1.	0.5	6
9	Fe ^{II} Spin Transition Materials Including an Amino-Ester 1,2,4-Triazole Derivative, Operating at, below, and above Room Temperature. Inorganic Chemistry, 2016, 55, 4278-4295.	4.0	39
10	Supramolecular homochiral helicity and zigzag hydrogen bonded chains in 1,2,4-triazole derived aminoester and aminoacid. New Journal of Chemistry, 2016, 40, 9025-9029.	2.8	3
11	Heteronuclear nanoparticles supported hydrotalcites containing Ni(II) and Fe(III) stable photocatalysts for Orange II degradation. Applied Clay Science, 2016, 132-133, 641-649.	5.2	16
12	Water effect on the spin-transition behavior of Fe(_{scp}) _{ii} (_{scp}) 1,2,4-triazole 1D chains embedded in pores of MCM-41. Journal of Materials Chemistry C, 2015, 3, 7802-7812.	5.5	46
13	Single-Walled Metal-Organic Nanotube Built from a Simple Synthon. Chemistry - A European Journal, 2015, 21, 4300-4307.	3.3	37
14	Two-Step Spin Transition in a 1D Fe ^{II} 1,2,4-Triazole Chain Compound. Chemistry - A European Journal, 2015, 21, 5843-5855.	3.3	28
15	Pressure Sensor via Optical Detection Based on a 1D Spin Transition Coordination Polymer. Sensors, 2015, 15, 2388-2398.	3.8	50
16	Electronic vs. structural ordering in a manganese(_{scp}) _{iii} (_{scp}) spin crossover complex. Chemical Communications, 2015, 51, 17540-17543.	4.1	77
17	Correction: Electronic vs. structural ordering in a manganese(_{scp}) _{iii} (_{scp}) spin crossover complex. Chemical Communications, 2015, 51, 17630-17630.	4.1	0
18	Spin conversion detected by Mössbauer spectroscopy and $\frac{1}{4}$ SR on a 1D Fell paramagnetic chain. Hyperfine Interactions, 2014, 226, 217-221.	0.5	5

#	ARTICLE	IF	CITATIONS
19	Spin-State Ordering on One Sub-lattice of a Mononuclear Iron(III) Spin Crossover Complex Exhibiting LIESST and TIESST. <i>Chemistry - A European Journal</i> , 2014, 20, 5613-5618.	3.3	83
20	New Mononuclear Cu(II) Complexes and 1D Chains with 4-Amino-4H-1,2,4-triazole. <i>International Journal of Molecular Sciences</i> , 2013, 14, 23597-23613.	4.1	13
21	Weak cooperativity in selected iron(II) 1D coordination polymers. , 2013, , 223-227.		1
22	Neutral and anionic duality of 1,2,4-triazole \pm -amino acid scaffold in 1D coordination polymers. , 2013, , 205-209.		0
23	Iron(<i>scp</i>) <i>ii</i> (<i>scp</i>) spin transition coordination polymers with a zigzag structure. <i>CrystEngComm</i> , 2012, 14, 1223-1231.	2.6	27
24	Impact of ligand spacer and counter-anion in selected 1D iron(II) spin crossover coordination polymers. <i>Hyperfine Interactions</i> , 2012, 205, 69-73.	0.5	8
25	Weak cooperativity in selected iron(II) 1D coordination polymers. <i>Hyperfine Interactions</i> , 2012, 205, 75-79.	0.5	5
26	Neutral and anionic duality of 1,2,4-triazole \pm -amino acid scaffold in 1D coordination polymers. <i>Hyperfine Interactions</i> , 2012, 205, 57-61.	0.5	2
27	Coordination preference and magnetic properties of FeII assemblies with a bis-azole bearing 1,2,4-triazole and tetrazole. <i>Hyperfine Interactions</i> , 2012, 204, 119-123.	0.5	4
28	Superlative Scaffold of 1,2,4-Triazole Derivative of Glycine Steering Linear Chain to a Chiral Helicate. <i>Crystal Growth and Design</i> , 2011, 11, 1375-1384.	3.0	26
29	Influence of Hydrogen Bonding on the Hysteresis Width in Iron(II) Spin-Crossover Complexes. <i>European Journal of Inorganic Chemistry</i> , 2011, 2011, 3193-3206.	2.0	100
30	Zinc complexes with 1,2,4-triazole functionalized amino acid derivatives: Synthesis, structure and β -lactamase assay. <i>Inorganica Chimica Acta</i> , 2011, 368, 21-28.	2.4	29
31	Spin Transition Sensors Based on β -Amino-Acid 1,2,4-Triazole Derivative. <i>International Journal of Molecular Sciences</i> , 2011, 12, 5339-5351.	4.1	24
32	Coordination Polymers and Metal Organic Frameworks Derived from 1,2,4-Triazole Amino Acid Linkers. <i>Polymers</i> , 2011, 3, 1750-1775.	4.5	61
33	Room temperature hysteretic spin transition in 1D iron(II) coordination polymers. <i>Journal of Physics: Conference Series</i> , 2010, 217, 012085.	0.4	20
34	Effect of texture alteration by thin film fabrication on the spin crossover of $[Fe(3-Br-phen)]_{2}(NCS)_{2} \cdot 0.5CH_{3}OH$. <i>Journal of Physics: Conference Series</i> , 2010, 217, 012032.	0.4	8
35	Insights into the Origin of Cooperative Effects in the Spin Transition of $[Fe(NH_{2}trz)_{3}(NO_{3})_{2}]$: the Role of Supramolecular Interactions Evidenced in the Crystal Structure of $[Cu(NH_{2}trz)_{3}(NO_{3})_{2}] \cdot H_{2}O$. <i>Inorganic Chemistry</i> , 2010, 49, 5723-5736.	4.0	131
36	Engineering Three-Dimensional Chains of Porous Nanoballs from a 1,2,4-Triazole-carboxylate Supramolecular Synthon. <i>Crystal Growth and Design</i> , 2010, 10, 1798-1807.	3.0	49

#	ARTICLE	IF	CITATIONS
37	Calorimetric measurements of diluted spin crossover complexes $[Fe_xM_{I\ddot{I}}(btr)_2(NCS)_2] \cdot H_2O$ with $M_{I\ddot{I}}=Zn$ and Ni . <i>Polyhedron</i> , 2009, 28, 2531-2536.	2.2	35
38	Prediction of the Spin Transition Temperature in Fe^{II} One-Dimensional Coordination Polymers: an Anion Based Database. <i>Inorganic Chemistry</i> , 2009, 48, 7838-7852.	4.0	116
39	Rapid Cooling Experiments and Use of an Anionic Nuclear Probe to Sense the Spin Transition of the 1D Coordination Polymers $[Fe(NH_2)_2]_{3-n}SnF_6 \cdot nH_2O$ ($n=4$ amino-1,2,4-triazole). <i>Chemistry - A European Journal</i> , 2008, 14, 3745-3758.	3.3	52
40	Iron(II) spin transition 1,2,4-triazole chain compounds with novel inorganic fluorinated counteranions. <i>Polyhedron</i> , 2007, 26, 2259-2263.	2.2	32