

# Katarzyna M Sawicka

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4270214/publications.pdf>

Version: 2024-02-01

8  
papers

113  
citations

1478458

6  
h-index

1720014

7  
g-index

9  
all docs

9  
docs citations

9  
times ranked

149  
citing authors

#	ARTICLE	IF	CITATIONS
1	Dronedarone (a multichannel blocker) enhances the anticonvulsant potency of lamotrigine, but not that of lacosamide, pregabalin and topiramate in the tonic-clonic seizure model in mice. <i>Epilepsy Research</i> , 2019, 154, 62-68.	1.6	7
2	Influence of dronedarone (a class III antiarrhythmic drug) on the anticonvulsant potency of four classical antiepileptic drugs in the tonic-clonic seizure model in mice. <i>Journal of Neural Transmission</i> , 2019, 126, 115-122.	2.8	4
3	A questionnaire-based, multicenter registry of resistant and pseudo-resistant arterial. <i>Wiadomości Lekarskie</i> , 2019, 72, 1866-1871.	0.3	0
4	Ivabradine attenuates the anticonvulsant potency of lamotrigine, but not that of lacosamide, pregabalin and topiramate in the tonic-clonic seizure model in mice. <i>Epilepsy Research</i> , 2017, 133, 67-70.	1.6	13
5	Influence of Ivabradine on the Anticonvulsant Action of Four Classical Antiepileptic Drugs Against Maximal Electroshock-Induced Seizures in Mice. <i>Neurochemical Research</i> , 2017, 42, 1038-1043.	3.3	15
6	Amiloride enhances the anticonvulsant action of various antiepileptic drugs in the mouse maximal electroshock seizure model. <i>Journal of Neural Transmission</i> , 2009, 116, 57-66.	2.8	30
7	Furosemide potentiates the anticonvulsant action of valproate in the mouse maximal electroshock seizure model. <i>Epilepsy Research</i> , 2007, 76, 66-72.	1.6	22
8	Chronically administered fluoxetine enhances the anticonvulsant activity of conventional antiepileptic drugs in the mouse maximal electroshock model. <i>European Journal of Pharmacology</i> , 2007, 567, 77-82.	3.5	22