

Philipp Schwemmer

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4277409/publications.pdf>

Version: 2024-02-01

33
papers

586
citations

687363

13
h-index

713466

21
g-index

35
all docs

35
docs citations

35
times ranked

930
citing authors

#	ARTICLE	IF	CITATIONS
1	Bird migration in space and time: chain migration by Eurasian curlew <i>Numenius arquata arquata</i> along the East Atlantic Flyway. <i>Journal of Avian Biology</i> , 2022, 2022, .	1.2	5
2	Migrating curlews on schedule: departure and arrival patterns of a long-distance migrant depend on time and breeding location rather than on wind conditions. <i>Movement Ecology</i> , 2021, 9, 9.	2.8	16
3	Analysis of local habitat selection and large-scale attraction/avoidance based on animal tracking data: is there a single best method?. <i>Movement Ecology</i> , 2021, 9, 20.	2.8	5
4	Spatio-temporal movement patterns and habitat choice of red foxes (<i>Vulpes vulpes</i>) and racoon dogs (<i>Nyctereutes procyonoides</i>) along the Wadden Sea coast. <i>European Journal of Wildlife Research</i> , 2021, 67, 1.	1.4	7
5	Timing of spring departure of long distance migrants correlates with previous year's conditions at their breeding site. <i>Biology Letters</i> , 2021, 17, 20210331.	2.3	8
6	Suitability of herring gulls (<i>Larus argentatus</i>) as indicators for detecting intertidal bivalve beds in the Wadden Sea. <i>Ecological Indicators</i> , 2021, 129, 107947.	6.3	1
7	Species composition of foraging birds in association with benthic fauna in four intertidal habitats of the Wadden Sea. <i>Estuarine, Coastal and Shelf Science</i> , 2020, 233, 106537.	2.1	20
8	Ecological insights from three decades of animal movement tracking across a changing Arctic. <i>Science</i> , 2020, 370, 712-715.	12.6	75
9	Modelling and predicting habitats for the neobiotic American razor clam <i>Ensis leei</i> in the Wadden Sea. <i>Estuarine, Coastal and Shelf Science</i> , 2019, 231, 106440.	2.1	6
10	Comparison of bivalve communities between moulting and wintering areas used by Common Scoter <i>Melanitta nigra</i> in the German North Sea. <i>Estuarine, Coastal and Shelf Science</i> , 2019, 229, 106398.	2.1	7
11	Impact of birds on intertidal food webs assessed with ecological network analysis. <i>Estuarine, Coastal and Shelf Science</i> , 2019, 219, 107-119.	2.1	10
12	Modelling distribution of common scoter (<i>Melanitta nigra</i>) by its predominant prey, the American razor clam (<i>Ensis leei</i>) and hydrodynamic parameters. <i>Estuarine, Coastal and Shelf Science</i> , 2019, 225, 106260.	2.1	12
13	A Ship Traffic Disturbance Vulnerability Index for Northwest European Seabirds as a Tool for Marine Spatial Planning. <i>Frontiers in Marine Science</i> , 2019, 6, .	2.5	19
14	An invasive alien bivalve apparently provides a novel food source for moulting and wintering benthic feeding sea ducks. <i>Helgoland Marine Research</i> , 2019, 73, .	1.3	3
15	Operational offshore wind farms and associated ship traffic cause profound changes in distribution patterns of Loons (<i>Gavia</i> spp.). <i>Journal of Environmental Management</i> , 2019, 231, 429-438.	7.8	48
16	Seasonal dynamics and functioning of the Sylt-RÄmÃ, Bight, northern Wadden Sea. <i>Estuarine, Coastal and Shelf Science</i> , 2018, 203, 100-118.	2.1	12
17	Decreasing $\delta^{13}C$ and $\delta^{15}N$ values in four coastal species at different trophic levels indicate a fundamental food-web shift in the southern North and Baltic Seas between 1988 and 2016. <i>Environmental Monitoring and Assessment</i> , 2018, 190, 461.	2.7	9
18	Intercolony variations in movement patterns and foraging behaviors among herring gulls (<i>Larus</i>) Tj ETQq0 0 0 rgBT/Qverlock_10 Tf 50 6.	1.9	29

#	ARTICLE	IF	CITATIONS
19	A fundamental study revisited: Quantitative evidence for territory quality in oystercatchers (<i>Haematopus ostralegus</i>) using GPS data loggers. <i>Ecology and Evolution</i> , 2017, 7, 285-294.	1.9	5
20	Interaction between birds and macrofauna within food webs of six intertidal habitats of the Wadden Sea. <i>PLoS ONE</i> , 2017, 12, e0176381.	2.5	17
21	Body mass change and diet switch tracked by stable isotopes indicate time spent at a stopover site during autumn migration in dunlins <i>Calidris alpina alpina</i> . <i>Journal of Avian Biology</i> , 2016, 47, 806-814.	1.2	8
22	Modelling small-scale foraging habitat use in breeding Eurasian oystercatchers (<i>Haematopus</i>) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 627 2016, 320, 322-333.	2.5	22
23	Migration routes of Eurasian Curlews (<i>Numenius arquata</i>) resting in the eastern Wadden Sea based on GPS telemetry. <i>Journal of Ornithology</i> , 2016, 157, 901-905.	1.1	10
24	Terrestrial and Marine Foraging Strategies of an Opportunistic Seabird Species Breeding in the Wadden Sea. <i>PLoS ONE</i> , 2016, 11, e0159630.	2.5	35
25	Assessment of contaminant levels and trophic relations at a World Heritage Site by measurements in a characteristic shorebird species. <i>Environmental Research</i> , 2015, 136, 163-172.	7.5	2
26	Weather-Related Winter Mortality of Eurasian Oystercatchers (<i>Haematopus ostralegus</i>) in the Northeastern Wadden Sea. <i>Waterbirds</i> , 2014, 37, 319-330.	0.3	15
27	Lesser black-backed gulls (<i>Larus fuscus</i>) consuming swimming crabs: An important link in the food web of the southern North Sea. <i>Estuarine, Coastal and Shelf Science</i> , 2013, 119, 71-78.	2.1	10
28	Effects of ship traffic on seabirds in offshore waters: implications for marine conservation and spatial planning. , 2011, 21, 1851-1860.		69
29	Spatial and temporal patterns of habitat use by Eurasian oystercatchers (<i>Haematopus ostralegus</i>) in the eastern Wadden Sea revealed using GPS data loggers. <i>Marine Biology</i> , 2011, 158, 541-550.	1.5	20
30	Influence of water flow velocity, water depth and colony distance on distribution and foraging patterns of terns in the Wadden Sea. <i>Fisheries Oceanography</i> , 2009, 18, 161-172.	1.7	18
31	Area utilization of gulls in a coastal farmland landscape: habitat mosaic supports niche segregation of opportunistic species. <i>Landscape Ecology</i> , 2008, 23, 355-367.	4.2	30
32	Regular habitat switch as an important feeding strategy of an opportunistic seabird species at the interface between land and sea. <i>Estuarine, Coastal and Shelf Science</i> , 2008, 77, 12-22.	2.1	30
33	Spatial patterns in at-sea behaviour during spring migration by little gulls (<i>Larus minutus</i>) in the southeastern North Sea. <i>Journal of Ornithology</i> , 2006, 147, 354-366.	1.1	2