

# L Belo

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/4282963/l-belo-publications-by-year.pdf>

**Version:** 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

124  
papers

2,555  
citations

28  
h-index

44  
g-index

138  
ext. papers

2,850  
ext. citations

3.6  
avg, IF

4.28  
L-index

#	Paper	IF	Citations
124	Subpopulations of High-Density Lipoprotein: Friends or Foes in Cardiovascular Disease Risk in Chronic Kidney Disease?. <i>Biomedicines</i> , <b>2021</b> , 9,	4.8	1
123	Interleukin 6 (rs1800795) and pentraxin 3 (rs2305619) polymorphisms-association with inflammation and all-cause mortality in end-stage-renal disease patients on dialysis. <i>Scientific Reports</i> , <b>2021</b> , 11, 14768	4.9	2
122	Cell-free DNA as a marker for the outcome of end-stage renal disease patients on haemodialysis. <i>CKJ: Clinical Kidney Journal</i> , <b>2021</b> , 14, 1371-1378	4.5	1
121	New Potential Biomarkers for Chronic Kidney Disease Management-A Review of the Literature. <i>International Journal of Molecular Sciences</i> , <b>2020</b> , 22,	6.3	15
120	Linkage of typically cytosolic peroxidases to erythrocyte membrane - A possible mechanism of protection in Hereditary Spherocytosis. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , <b>2020</b> , 1862, 183172	3.8	6
119	School-based soccer practice is an effective strategy to improve cardiovascular and metabolic risk factors in overweight children. <i>Progress in Cardiovascular Diseases</i> , <b>2020</b> , 63, 807-812	8.5	3
118	The Protective Role of Adiponectin for Lipoproteins in End-Stage Renal Disease Patients: Relationship with Diabetes and Body Mass Index. <i>Oxidative Medicine and Cellular Longevity</i> , <b>2019</b> , 2019, 3021785	6.7	10
117	Hepcidin in chronic kidney disease anemia. <i>Vitamins and Hormones</i> , <b>2019</b> , 110, 243-264	2.5	10
116	Long Pentraxin 3 as a Broader Biomarker for Multiple Risk Factors in End-Stage Renal Disease: Association with All-Cause Mortality. <i>Mediators of Inflammation</i> , <b>2019</b> , 2019, 3295725	4.3	6
115	Hepcidin and diabetes are independently related with soluble transferrin receptor levels in chronic dialysis patients. <i>Renal Failure</i> , <b>2019</b> , 41, 662-672	2.9	3
114	Weight loss achieved by bariatric surgery modifies high-density lipoprotein subfractions and low-density lipoprotein oxidation towards atheroprotection. <i>Clinical Biochemistry</i> , <b>2019</b> , 63, 46-53	3.5	11
113	Influence of the 6-month physical activity programs on renal function in obese boys. <i>Pediatric Research</i> , <b>2018</b> , 83, 1011-1015	3.2	2
112	Impact of weight loss on inflammation and red blood cell biomarkers after laparoscopic gastric banding surgery. <i>Journal of Investigative Medicine</i> , <b>2018</b> , 66, 304-308	2.9	6
111	SP342HEPCIDIN-25 AND TREATMENT WITH ERYTHROPOIESIS STIMULATING AGENTS ARE INDEPENDENTLY RELATED WITH ERYTHROPOIESIS IN CHRONIC HEMODIALYSIS PATIENTS. <i>Nephrology Dialysis Transplantation</i> , <b>2018</b> , 33, i460-i460	4.3	1
110	Physical exercise intervention at school improved hepcidin, inflammation, and iron metabolism in overweight and obese children and adolescents. <i>Pediatric Research</i> , <b>2017</b> , 82, 781-788	3.2	10
109	Adiponectin, Inflammation and Cardiometabolic Risk Factors in Paediatric Obese Patients: Impact of Interventional Studies <b>2017</b> ,		1
108	Renal cell carcinoma: a critical analysis of metabolomic biomarkers emerging from current model systems. <i>Translational Research</i> , <b>2017</b> , 180, 1-11	11	24

107	Iron therapy in chronic kidney disease: Recent changes, benefits and risks. <i>Blood Reviews</i> , <b>2016</b> , 30, 65-72	1.1	23
106	Exercise intervention and cardiovascular risk factors in obese children. Comparison between obese youngsters taking part in a physical activity school-based programme with and without individualised diet counselling: the ACORDA project. <i>Annals of Human Biology</i> , <b>2016</b> , 43, 183-90	1.7	10
105	Impact of a School-Based Intervention Protocol - ACORDA Project - On Adipokines in An Overweight and Obese Pediatric Population. <i>Pediatric Exercise Science</i> , <b>2016</b> , 28, 407-16	2	16
104	Recombinant human erythropoietin-induced erythropoiesis regulates hepcidin expression over iron status in the rat. <i>Blood Cells, Molecules, and Diseases</i> , <b>2016</b> , 59, 63-70	2.1	5
103	Effects of 6-month soccer and traditional physical activity programmes on body composition, cardiometabolic risk factors, inflammatory, oxidative stress markers and cardiorespiratory fitness in obese boys. <i>Journal of Sports Sciences</i> , <b>2016</b> , 34, 1822-9	3.6	29
102	SP313LIVER IRON IS A MAJOR REGULATOR OF HEPCIDIN GENE EXPRESSION VIA BMP/SMAD PATHWAY IN A RAT MODEL OF CHRONIC RENAL FAILURE UNDER TREATMENT WITH HIGH rHuEPO DOSES. <i>Nephrology Dialysis Transplantation</i> , <b>2016</b> , 31, i194-i194	4.3	1
101	Biomarkers in bladder cancer: A metabolomic approach using in vitro and ex vivo model systems. <i>International Journal of Cancer</i> , <b>2016</b> , 139, 256-68	7.5	49
100	Renal risk-benefit determinants of recombinant human erythropoietin therapy in the remnant kidney rat model - hypertension, anaemia, inflammation and drug dose. <i>Clinical and Experimental Pharmacology and Physiology</i> , <b>2016</b> , 43, 343-54	3	4
99	Adipokine Gene Single-Nucleotide Polymorphisms in Portuguese Obese Adolescents: Associations with Plasma Concentrations of Adiponectin, Resistin, IL-6, IL-1 $\beta$ and TNF- $\alpha$ <i>Childhood Obesity</i> , <b>2016</b> , 12, 300-13	2.5	11
98	Pathological and molecular mechanisms underlying resistance to recombinant human erythropoietin therapy in the remnant kidney rat model of chronic kidney disease associated anemia. <i>Biochimie</i> , <b>2016</b> , 125, 150-62	4.6	7
97	Impaired renal endothelial nitric oxide synthase and reticulocyte production as modulators of hypertension induced by rHuEPO in the rat. <i>Life Sciences</i> , <b>2016</b> , 151, 147-156	6.8	4
96	Liver iron is a major regulator of hepcidin gene expression via BMP/SMAD pathway in a rat model of chronic renal failure under treatment with high rHuEPO doses. <i>BioFactors</i> , <b>2016</b> , 42, 296-306	6.1	8
95	Comparison between CDC and WHO BMI z-score and their relation with metabolic risk markers in Northern Portuguese obese adolescents. <i>Diabetology and Metabolic Syndrome</i> , <b>2015</b> , 7, 32	5.6	5
94	Iron-hepcidin dysmetabolism, anemia and renal hypoxia, inflammation and fibrosis in the remnant kidney rat model. <i>PLoS ONE</i> , <b>2015</b> , 10, e0124048	3.7	23
93	Bilirubin is independently associated with oxidized LDL levels in young obese patients. <i>Diabetology and Metabolic Syndrome</i> , <b>2015</b> , 7, 4	5.6	11
92	Protective effect of <i>C. sativa</i> leaf extract against UV mediated-DNA damage in a human keratinocyte cell line. <i>Journal of Photochemistry and Photobiology B: Biology</i> , <b>2015</b> , 144, 28-34	6.7	28
91	Resistance to Recombinant Human Erythropoietin Therapy in a Rat Model of Chronic Kidney Disease Associated Anemia. <i>International Journal of Molecular Sciences</i> , <b>2015</b> , 17,	6.3	6
90	Adiponectin and markers of metabolic syndrome in obese children and adolescents: impact of 8-month regular physical exercise program. <i>Pediatric Research</i> , <b>2014</b> , 76, 159-65	3.2	24

89	The effect of olive leaf supplementation on the constituents of blood and oxidative stability of red blood cells. <i>Journal of Functional Foods</i> , <b>2014</b> , 9, 271-279	5.1	17
88	Haptoglobin 2-2 phenotype is associated with decreased serum iron levels in endstage renal disease patients resistant to rhEPO therapy. <i>British Journal of Biomedical Science</i> , <b>2014</b> , 71, 79-81	1.6	
87	Body fat percentage is a major determinant of total bilirubin independently of UGT1A1*28 polymorphism in young obese. <i>PLoS ONE</i> , <b>2014</b> , 9, e98467	3.7	16
86	Waldenström's macroglobulinemia - a review. <i>Revista Da Associação Médica Brasileira</i> , <b>2014</b> , 60, 490-499	1.4	1
85	Potential cardiovascular risk protection of bilirubin in end-stage renal disease patients under hemodialysis. <i>BioMed Research International</i> , <b>2014</b> , 2014, 175286	3	11
84	Iron as the key modulator of hepcidin expression in erythroid antibody-mediated hypoplasia. <i>BioMed Research International</i> , <b>2014</b> , 2014, 421304	3	4
83	TLR4 and TLR9 Polymorphisms Effect on Inflammatory Response in End-Stage Renal Disease Patients. <i>European Journal of Inflammation</i> , <b>2014</b> , 12, 521-529	0.3	2
82	Aging is Associated with Impaired Renal Function, INF-gamma Induced Inflammation and with Alterations in Iron Regulatory Proteins Gene Expression <b>2014</b> , 5, 356-65		10
81	Letter to the editor: A potential mechanism for the pathogenesis of psoriasis vulgaris. <i>International Journal of Dermatology</i> , <b>2013</b> , 52, 1429-32	1.7	
80	Principal determinants of the length of remission of psoriasis vulgaris after topical, NB-UVB, and PUVA therapy: a follow-up study. <i>American Journal of Clinical Dermatology</i> , <b>2013</b> , 14, 49-53	7.1	15
79	Protective activity of hydroxytyrosol metabolites on erythrocyte oxidative-induced hemolysis. <i>Journal of Agricultural and Food Chemistry</i> , <b>2013</b> , 61, 6636-42	5.7	30
78	Risk factors for mortality in hemodialysis patients: two-year follow-up study. <i>Disease Markers</i> , <b>2013</b> , 35, 791-8	3.2	31
77	Circulating cell-free DNA levels in hemodialysis patients and its association with inflammation, iron metabolism, and rhEPO doses. <i>Hemodialysis International</i> , <b>2013</b> , 17, 664-7	1.7	10
76	Body mass index and resistance to recombinant human erythropoietin therapy in maintenance hemodialysis patients. <i>Renal Failure</i> , <b>2013</b> , 35, 1392-8	2.9	6
75	Vascular access versus the effect of statins on inflammation and fibrinolysis in renal dialysis patients. <i>Journal of Vascular Access</i> , <b>2013</b> , 14, 335-41	1.8	3
74	Oxidized low-density lipoprotein and lipoprotein(a) levels in chronic kidney disease patients under hemodialysis: influence of adiponectin and of a polymorphism in the apolipoprotein(a) gene. <i>Hemodialysis International</i> , <b>2012</b> , 16, 481-90	1.7	17
73	Adiponectin is an independent predictor of tissue plasminogen activator levels in patients under haemodialysis. <i>Scandinavian Journal of Urology and Nephrology</i> , <b>2012</b> , 46, 461-5		
72	Erythroid disturbances before and after treatment of Portuguese psoriasis vulgaris patients: a cross-sectional and longitudinal study. <i>American Journal of Clinical Dermatology</i> , <b>2012</b> , 13, 37-47	7.1	14

71	Cardiac antiapoptotic and proproliferative effect of recombinant human erythropoietin in a moderate stage of chronic renal failure in the rat. <i>Journal of Pharmacy and Bioallied Sciences</i> , <b>2012</b> , 4, 76-83	1.1	15
70	Major determinants of BMP-2 serum levels in hemodialysis patients. <i>Renal Failure</i> , <b>2012</b> , 34, 1355-8	2.9	2
69	Inflammatory disturbances in preeclampsia: relationship between maternal and umbilical cord blood. <i>Journal of Pregnancy</i> , <b>2012</b> , 2012, 684384	2.5	51
68	Main determinants of PON1 activity in hemodialysis patients. <i>American Journal of Nephrology</i> , <b>2012</b> , 36, 317-23	4.6	13
67	Cardiovascular risk factors in portuguese obese children and adolescents: impact of small reductions in body mass index imposed by lifestyle modifications. <i>The Open Biochemistry Journal</i> , <b>2012</b> , 6, 43-50	0.9	15
66	Complementary markers for the clinical severity classification of hereditary spherocytosis in unsplenectomized patients. <i>Blood Cells, Molecules, and Diseases</i> , <b>2011</b> , 46, 166-70	2.1	12
65	Erythropoiesis versus inflammation in Hereditary Spherocytosis clinical outcome. <i>Clinical Biochemistry</i> , <b>2011</b> , 44, 1137-1143	3.5	4
64	Health-related quality of life in Portuguese psoriatic patients: relation with Psoriasis Area and Severity Index and different types of classical psoriatic treatment. <i>Journal of Dermatology</i> , <b>2011</b> , 38, 816-9	1.6	6
63	Elastase release during the hemodialysis procedure seems to induce changes in red blood cell membrane proteins. <i>Hemodialysis International</i> , <b>2011</b> , 15, 429-31	1.7	3
62	Apoptosis of peripheral CD4(+) T-lymphocytes in end-stage renal disease patients under hemodialysis and rhEPO therapies. <i>Renal Failure</i> , <b>2011</b> , 33, 138-43	2.9	21
61	Comparison of Bio-Plex measurements with standard techniques. <i>Clinical Chemistry and Laboratory Medicine</i> , <b>2011</b> , 50, 399-402	5.9	1
60	Circulating adipokine levels in Portuguese patients with psoriasis vulgaris according to body mass index, severity and therapy. <i>Journal of the European Academy of Dermatology and Venereology</i> , <b>2010</b> , 24, 1386-94	4.6	90
59	Neutrophil and monocyte activation in chronic kidney disease patients under hemodialysis and its relationship with resistance to recombinant human erythropoietin and to the hemodialysis procedure. <i>Hemodialysis International</i> , <b>2010</b> , 14, 295-301	1.7	15
58	Interleukin (IL)-22, IL-17, IL-23, IL-8, vascular endothelial growth factor and tumour necrosis factor- $\beta$ levels in patients with psoriasis before, during and after psoralen-ultraviolet A and narrowband ultraviolet B therapy. <i>British Journal of Dermatology</i> , <b>2010</b> , 163, 1282-90	4	100
57	Erythrocyte membrane protein destabilization versus clinical outcome in 160 Portuguese Hereditary Spherocytosis patients. <i>British Journal of Haematology</i> , <b>2010</b> , 149, 785-94	4.5	19
56	Recombinant human erythropoietin treatment protects the cardio-renal axis in a model of moderate chronic renal failure. <i>Renal Failure</i> , <b>2010</b> , 32, 1073-80	2.9	7
55	Lipid levels including oxidized LDL in women with history of preeclampsia. <i>Hypertension in Pregnancy</i> , <b>2010</b> , 29, 93-100	2	14
54	Psoriasis therapy and cardiovascular risk factors: a 12-week follow-up study. <i>American Journal of Clinical Dermatology</i> , <b>2010</b> , 11, 423-32	7.1	30

53	Powerful protective role of 3,4-dihydroxyphenylethanol-elenolic acid dialdehyde against erythrocyte oxidative-induced hemolysis. <i>Journal of Agricultural and Food Chemistry</i> , <b>2010</b> , 58, 135-40	5.7	44
52	Plant aqueous extracts: Antioxidant capacity via haemolysis and bacteriophage P22 protection. <i>Food Control</i> , <b>2010</b> , 21, 633-638	6.2	13
51	Hereditary spherocytosis and the (TA)nTAA polymorphism of UGT1A1 gene promoter region--a comparison of the bilirubin plasmatic levels in the different clinical forms. <i>Blood Cells, Molecules, and Diseases</i> , <b>2010</b> , 44, 117-9	2.1	6
50	C-reactive protein and leucocyte activation in psoriasis vulgaris according to severity and therapy. <i>Journal of the European Academy of Dermatology and Venereology</i> , <b>2010</b> , 24, 789-96	4.6	85
49	In vitro studies with β-catalasemic-like β-erythrocytes and hydrogen peroxide: attention to the formation of lysis resistant erythrocytes. <i>International Journal of Laboratory Hematology</i> , <b>2010</b> , 32, 127-31	2.5	2
48	Antioxidant activity of chitoooligosaccharides upon two biological systems: Erythrocytes and bacteriophages. <i>Carbohydrate Polymers</i> , <b>2010</b> , 79, 1101-1106	10.3	58
47	Leukocyte Count versus C-Reactive Protein Levels in Obese Portuguese Patients Aged 6-12 Years Old. <i>The Open Biochemistry Journal</i> , <b>2010</b> , 4, 72-6	0.9	15
46	Lipoprotein(a) levels in obese Portuguese children and adolescents: contribution of the pentanucleotide repeat (TTTTA)n polymorphism in the apolipoprotein(a) gene. <i>JAMA Pediatrics</i> , <b>2009</b> , 163, 393-4		2
45	Hepcidin serum levels and resistance to recombinant human erythropoietin therapy in haemodialysis patients. <i>Acta Haematologica</i> , <b>2009</b> , 122, 226-9	2.7	33
44	Lipid profile in Portuguese obese children and adolescents: interaction of apolipoprotein E polymorphism with adiponectin levels. <i>JAMA Pediatrics</i> , <b>2009</b> , 163, 1030-6		9
43	Fetal and maternal angiogenic/anti-angiogenic factors in normal and preeclamptic pregnancy. <i>Growth Factors</i> , <b>2009</b> , 27, 345-51	1.6	25
42	Characterization of a rat model of moderate chronic renal failure--focus on hematological, biochemical, and cardio-renal profiles. <i>Renal Failure</i> , <b>2009</b> , 31, 833-42	2.9	8
41	Effects of olive oil polyphenols on erythrocyte oxidative damage. <i>Molecular Nutrition and Food Research</i> , <b>2009</b> , 53, 609-16	5.9	82
40	Erythropoietin promotes deleterious cardiovascular effects and mortality risk in a rat model of chronic sports doping. <i>Cardiovascular Toxicology</i> , <b>2009</b> , 9, 201-10	3.4	18
39	Linkage of cytosolic peroxiredoxin 2 to erythrocyte membrane imposed by hydrogen peroxide-induced oxidative stress. <i>Blood Cells, Molecules, and Diseases</i> , <b>2009</b> , 43, 68-73	2.1	24
38	Haemostatic factors in women with history of preeclampsia. <i>Thrombosis Research</i> , <b>2009</b> , 124, 52-6	8.2	13
37	Circulating levels of adiponectin, oxidized LDL and C-reactive protein in Portuguese patients with psoriasis vulgaris, according to body mass index, severity and duration of the disease. <i>Journal of Dermatological Science</i> , <b>2009</b> , 55, 202-4	4.3	48
36	Erythrocyte changes in preeclampsia: relationship between maternal and cord blood erythrocyte damage. <i>Journal of Perinatal Medicine</i> , <b>2009</b> , 37, 19-27	2.7	12

35	Altered alanine aminotransferase and gamma-glutamyl transpeptidase in women with history of preeclampsia: association with waist-to-hip ratio and body mass index. <i>European Journal of Gastroenterology and Hepatology</i> , <b>2009</b> , 21, 196-200	2.2	3
34	Effects of chitoooligosaccharides on human red blood cell morphology and membrane protein structure. <i>Biomacromolecules</i> , <b>2008</b> , 9, 3346-52	6.9	46
33	Changes in red blood cells membrane protein composition during hemodialysis procedure. <i>Renal Failure</i> , <b>2008</b> , 30, 971-5	2.9	12
32	Relationship between maternal and cord blood hemostatic disturbances in preeclamptic pregnancies. <i>Thrombosis Research</i> , <b>2008</b> , 123, 219-24	8.2	18
31	IL-7 serum levels and lymphopenia in hemodialysis patients, non-responders to recombinant human erythropoietin therapy. <i>Blood Cells, Molecules, and Diseases</i> , <b>2008</b> , 41, 134-5	2.1	2
30	Presence of cytosolic peroxiredoxin 2 in the erythrocyte membrane of patients with hereditary spherocytosis. <i>Blood Cells, Molecules, and Diseases</i> , <b>2008</b> , 41, 5-9	2.1	31
29	Role of prohepcidin, inflammatory markers and iron status in resistance to rhEPO therapy in hemodialysis patients. <i>American Journal of Nephrology</i> , <b>2008</b> , 28, 677-83	4.6	31
28	Neutrophil activation and resistance to recombinant human erythropoietin therapy in hemodialysis patients. <i>American Journal of Nephrology</i> , <b>2008</b> , 28, 935-40	4.6	27
27	Altered erythrocyte membrane protein composition in chronic kidney disease stage 5 patients under haemodialysis and recombinant human erythropoietin therapy. <i>Blood Purification</i> , <b>2008</b> , 26, 267-73	3.1	13
26	DMT1 (NRAMP2/DCT1) genetic variability and resistance to recombinant human erythropoietin therapy in chronic kidney disease patients under haemodialysis. <i>Acta Haematologica</i> , <b>2008</b> , 120, 11-3	2.7	4
25	Similarities between pre-eclampsia and atherosclerosis: a protective effect of physical exercise?. <i>Current Medicinal Chemistry</i> , <b>2008</b> , 15, 2223-9	4.3	15
24	Cross-Talk between Inflammation, Coagulation/Fibrinolysis and Vascular access in Hemodialysis Patients. <i>Journal of Vascular Access</i> , <b>2008</b> , 9, 248-253	1.8	25
23	Inflammation, T-cell phenotype, and inflammatory cytokines in chronic kidney disease patients under hemodialysis and its relationship to resistance to recombinant human erythropoietin therapy. <i>Journal of Clinical Immunology</i> , <b>2008</b> , 28, 268-75	5.7	64
22	Adhesion molecules (VCAM-1 and ICAM-1) and C-reactive protein in women with history of preeclampsia. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , <b>2008</b> , 87, 969-71	3.8	14
21	Fetal lipoprotein changes in pre-eclampsia. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , <b>2008</b> , 87, 628-34	3.8	27
20	Band 3 Profile as a Marker of Erythrocyte Changes in Chronic Kidney Disease Patients. <i>The Open Clinical Chemistry Journal</i> , <b>2008</b> , 1, 57-63		6
19	Cross-talk between inflammation,coagulation/fibrinolysis and vascular access in hemodialysis patients. <i>Journal of Vascular Access</i> , <b>2008</b> , 9, 248-53	1.8	8
18	Enhanced thrombotic risk in pregnancy and postpartum: association with CRP elevation?. <i>American Journal of Obstetrics and Gynecology</i> , <b>2007</b> , 196, e24	6.4	

17	Pre-eclampsia Versus Cardiovascular Disease Versus CRP. <i>Current Hypertension Reviews</i> , <b>2006</b> , 2, 317-323.	3.3	
16	Serum levels of VEGF and TNF-alpha and their association with C-reactive protein in patients with endometriosis. <i>Archives of Gynecology and Obstetrics</i> , <b>2006</b> , 273, 227-31	2.5	95
15	High-density lipoprotein particles may regulate hemostasis in human pregnancy. <i>Fertility and Sterility</i> , <b>2005</b> , 84, 1021-2	4.8	2
14	Oxidized-LDL levels in normal and pre-eclamptic pregnancies: contribution of LDL particle size. <i>Atherosclerosis</i> , <b>2005</b> , 183, 185-6	3.1	6
13	Fluctuations in C-reactive protein concentration and neutrophil activation during normal human pregnancy. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , <b>2005</b> , 123, 46-51	2.4	103
12	Are we employing the most effective CA 125 and CA 19-9 cut-off values to detect endometriosis?. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , <b>2005</b> , 123, 254-5	2.4	25
11	Erythropoietin levels in the different clinical forms of hereditary spherocytosis. <i>British Journal of Haematology</i> , <b>2005</b> , 131, 534-42	4.5	13
10	Protein deficiency balance as a predictor of clinical outcome in hereditary spherocytosis. <i>European Journal of Haematology</i> , <b>2005</b> , 74, 374-80	3.8	19
9	LDL size, total antioxidant status and oxidised LDL in normal human pregnancy: a longitudinal study. <i>Atherosclerosis</i> , <b>2004</b> , 177, 391-9	3.1	72
8	Apolipoprotein E and cholesteryl ester transfer protein polymorphisms in normal and preeclamptic pregnancies. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , <b>2004</b> , 112, 9-15	2.4	34
7	Neutrophil activation and C-reactive protein concentration in preeclampsia. <i>Hypertension in Pregnancy</i> , <b>2003</b> , 22, 129-41	2	56
6	Band 3 as a marker of erythrocyte changes in pregnancy. <i>European Journal of Haematology</i> , <b>2002</b> , 69, 145-51	3.8	18
5	Elevated tissue plasminogen activator as a potential marker of endothelial dysfunction in pre-eclampsia: correlation with proteinuria. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , <b>2002</b> , 109, 1250-5	3.7	38
4	Changes in LDL size and HDL concentration in normal and preeclamptic pregnancies. <i>Atherosclerosis</i> , <b>2002</b> , 162, 425-32	3.1	117
3	Lipoprotein(a): a longitudinal versus a cross-sectional study in normal pregnancy and its levels in preeclampsia. <i>Atherosclerosis</i> , <b>2002</b> , 165, 393-5	3.1	9
2	Erythrocyte damage and leukocyte activation in ischemic stroke. <i>Clinica Chimica Acta</i> , <b>2002</b> , 320, 29-35	6.2	36
1	Leukocyte activation, erythrocyte damage, lipid profile and oxidative stress imposed by high competition physical exercise in adolescents. <i>Clinica Chimica Acta</i> , <b>2001</b> , 306, 119-26	6.2	86