

# Slavica Marinkovic

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4286222/publications.pdf>

Version: 2024-02-01

13

papers

53

citations

1937685

4

h-index

1720034

7

g-index

13

all docs

13

docs citations

13

times ranked

86

citing authors

#	ARTICLE	IF	CITATIONS
1	Diversity of phytoplasmas identified in the polyphagous leafhopper <i>Euscelis incisus</i> (Cicadellidae,) Tj ETQq1 1 0.784314 rgBT /Overlock European Journal of Plant Pathology, 2020, 156, 201-221.	1.7	14
2	Geometric morphometric study of geographic and host-related variability in <i>Aceria</i> spp. (Acarci:) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 702 321-335.	1.6	9
3	Genetic Diversity of Flavescence DorÃ©e Phytoplasmas in Vineyards of Serbia: From the Widespread Occurrence of Autochthonous Map-M51 to the Emergence of Endemic Map-FD2 (Vectotype II) and New Map-FD3 (Vectotype III) Epidemic Genotypes. Agronomy, 2022, 12, 448.	3.0	6
4	<p class="HeadingRunIn"><strong>Eriophyid mite <em> <i>Aceria artemisiifoliae</i> </em>sp.nov. (Acarci:) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 artemisiifolia </em> L. (Asteraceae) in Serbia</strong></p>. Systematic and Applied Acarology, 2016, 21, 919.	0.5	4
5	Symptomatology, (Co)occurrence and Differential Diagnostic PCR Identification of â€“Ca. <i>Phytoplasma solaniâ€™</i> and â€“Ca. <i>Phytoplasma convolvuliâ€™</i> in Field Bindweed. Pathogens, 2021, 10, 160.	2.8	4
6	Three new species and new records of eriophyoid mites (Acarci, Eriophyoidea) from Karadag Nature Reserve, Crimea. Systematic and Applied Acarology, 2018, 23, 628.	0.5	3
7	<p class="Body">Supplementary description of three species from the subfamily Cecidophyinae (Eriophyoidea: Eriophyidae) from the Nalepa collection. Systematic and Applied Acarology, 2018, 23, 838.	0.5	3
8	Framework for risk assessment of â€“Candidatus <i>Phytoplasma solaniâ€™</i> associated diseases outbreaks in agroecosystems in Serbia. Journal of Plant Pathology, 0, , 1.	1.2	3
9	New species and records of phytoptids (Eriophyoidea, Phytoptidae) from cinquefoils (Rosaceae,) Tj ETQq1 1 0.784314 rgBT /Overlock 10	0.5	2
10	The â€œcode redâ€ for Balkan vineyards: occurrence of <i>Orientus ishidae</i> (Matsumura, 1902) (Hemiptera:) Tj ETQq0 0 0 rgBT /Overlock 1.1	2	2
11	Comparative morphological analysis of apple blister mite, <i>Eriophyes malii</i> Nal., a new pest in Serbia. Pesticidi I Fitomedicina = Pesticides and Phytomedicine, 2014, 29, 123-130.	0.2	2
12	<p class="Body"><strong>Supplementary description of five species from the genus <em> <i>Cecidophyopsis</i> </em> (Eriophyoidea: Eriophyidae: Cecidophyinae)</strong></p>. Systematic and Applied Acarology, 2019, 24, 1555-1578.	0.5	1
13	Can polyphagous insect vectors contribute to phytoplasma inventory in diverse ecosystems?. Phytopathogenic Mollicutes, 2019, 9, 103.	0.1	0