

Anna-Maria Larsson

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4290434/publications.pdf>

Version: 2024-02-01

16
papers

541
citations

932766

10
h-index

940134

16
g-index

16
all docs

16
docs citations

16
times ranked

1082
citing authors

#	ARTICLE	IF	CITATIONS
1	Prognostic impact of circulating tumor cell apoptosis and clusters in serial blood samples from patients with metastatic breast cancer in a prospective observational cohort. <i>BMC Cancer</i> , 2016, 16, 433.	1.1	125
2	Systemic Monocytic-MDSCs Are Generated from Monocytes and Correlate with Disease Progression in Breast Cancer Patients. <i>PLoS ONE</i> , 2015, 10, e0127028.	1.1	116
3	Longitudinal enumeration and cluster evaluation of circulating tumor cells improve prognostication for patients with newly diagnosed metastatic breast cancer in a prospective observational trial. <i>Breast Cancer Research</i> , 2018, 20, 48.	2.2	80
4	Clinical relevance of systemic monocytic-MDSCs in patients with metastatic breast cancer. <i>Cancer Immunology, Immunotherapy</i> , 2020, 69, 435-448.	2.0	44
5	Molecular characterization of circulating tumor cells from patients with metastatic breast cancer reflects evolutionary changes in gene expression under the pressure of systemic therapy. <i>Oncotarget</i> , 2017, 8, 45544-45565.	0.8	38
6	A novel method for downstream characterization of breast cancer circulating tumor cells following CellSearch isolation. <i>Journal of Translational Medicine</i> , 2015, 13, 126.	1.8	23
7	Impact of systemic therapy on circulating leukocyte populations in patients with metastatic breast cancer. <i>Scientific Reports</i> , 2019, 9, 13451.	1.6	21
8	EPO-independent functional EPO receptor in breast cancer enhances estrogen receptor activity and promotes cell proliferation. <i>Biochemical and Biophysical Research Communications</i> , 2014, 445, 163-169.	1.0	14
9	Serial evaluation of serum thymidine kinase activity is prognostic in women with newly diagnosed metastatic breast cancer. <i>Scientific Reports</i> , 2020, 10, 4484.	1.6	14
10	Human G-MDSCs are neutrophils at distinct maturation stages promoting tumor growth in breast cancer. <i>Life Science Alliance</i> , 2020, 3, e202000893.	1.3	14
11	PAM50 Intrinsic Subtype Profiles in Primary and Metastatic Breast Cancer Show a Significant Shift toward More Aggressive Subtypes with Prognostic Implications. <i>Cancers</i> , 2021, 13, 1592.	1.7	11
12	The Distribution of Circulating Tumor Cells Is Different in Metastatic Lobular Compared to Ductal Carcinoma of the Breast—Long-Term Prognostic Significance. <i>Cells</i> , 2020, 9, 1718.	1.8	10
13	Evolution of Estrogen Receptor Status from Primary Tumors to Metastasis and Serially Collected Circulating Tumor Cells. <i>International Journal of Molecular Sciences</i> , 2020, 21, 2885.	1.8	9
14	Mammographic tumour appearance is related to clinicopathological factors and surrogate molecular breast cancer subtype. <i>Scientific Reports</i> , 2020, 10, 20814.	1.6	8
15	Peripheral Blood Mononuclear Cell Populations Correlate with Outcome in Patients with Metastatic Breast Cancer. <i>Cells</i> , 2022, 11, 1639.	1.8	8
16	High Levels of Expression of Cartilage Oligomeric Matrix Protein in Lymph Node Metastases in Breast Cancer Are Associated with Reduced Survival. <i>Cancers</i> , 2021, 13, 5876.	1.7	6