

# Jing-Jing Li

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4298259/publications.pdf>

Version: 2024-02-01

29  
papers

817  
citations

567281

15  
h-index

526287

27  
g-index

29  
all docs

29  
docs citations

29  
times ranked

1375  
citing authors

#	ARTICLE	IF	CITATIONS
1	Associations of Clinical and Molecular Characteristics with the Response to Immune Checkpoint Blockade in Advanced Gastric Cancers. <i>Journal of Oncology</i> , 2022, 2022, 1-10.	1.3	0
2	Targeting USP9X-AMPK Axis in ARID1A-Deficient Hepatocellular Carcinoma. <i>Cellular and Molecular Gastroenterology and Hepatology</i> , 2022, 14, 101-127.	4.5	17
3	Systemic Inflammatory Markers of Resectable Colorectal Cancer Patients with Different Mismatch Repair Gene Status. <i>Cancer Management and Research</i> , 2021, Volume 13, 2925-2935.	1.9	5
4	NET1 promotes HCC growth and metastasis in vitro and in vivo via activating the Akt signaling pathway. <i>Aging</i> , 2021, 13, 10672-10687.	3.1	5
5	PPDPF alleviates hepatic steatosis through inhibition of mTOR signaling. <i>Nature Communications</i> , 2021, 12, 3059.	12.8	18
6	Antifungal agent Terbinafine restrains tumor growth in preclinical models of hepatocellular carcinoma via AMPK-mTOR axis. <i>Oncogene</i> , 2021, 40, 5302-5313.	5.9	11
7	Clinicopathological Characteristics and Prognosis of Signet Ring Gastric Cancer: A Population-Based Study. <i>Frontiers in Oncology</i> , 2021, 11, 580545.	2.8	13
8	INTS6 promotes colorectal cancer progression by activating of AKT and ERK signaling. <i>Experimental Cell Research</i> , 2021, 407, 112826.	2.6	1
9	Ochratoxin A Induces Steatosis via PPAR $\gamma$ -CD36 Axis. <i>Toxins</i> , 2021, 13, 802.	3.4	12
10	Clinicopathologic characteristics of resectable colorectal cancer with mismatch repair protein defects in Chinese population. <i>Medicine (United States)</i> , 2020, 99, e20554.	1.0	4
11	Chromatin remodeling factor ARID2 suppresses hepatocellular carcinoma metastasis via DNMT1-Snail axis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020, 117, 4770-4780.	7.1	76
12	Epithelial V $\alpha$ 14e1 antigen 1 promotes hepatocellular carcinoma growth and metastasis via the ERBB $\beta$ -PI3K-AKT pathway. <i>Cancer Science</i> , 2020, 111, 1500-1513.	3.9	11
13	Expression levels of EPHB4, EFNB2 and caspase $\beta$ 8 are associated with clinicopathological features and progression of esophageal squamous cell cancer. <i>Oncology Letters</i> , 2020, 19, 917-929.	1.8	5
14	Scinderin suppresses cell proliferation and predicts the poor prognosis of hepatocellular carcinoma. <i>Oncology Letters</i> , 2020, 19, 2011-2020.	1.8	4
15	BMP10 suppresses hepatocellular carcinoma progression via PTPRS-STAT3 axis. <i>Oncogene</i> , 2019, 38, 7281-7293.	5.9	19
16	CHML promotes liver cancer metastasis by facilitating Rab14 recycle. <i>Nature Communications</i> , 2019, 10, 2510.	12.8	32
17	Liver cancer: WISP3 suppresses hepatocellular carcinoma progression by negative regulation of $\beta$ -catenin/TCF/LEF signalling. <i>Cell Proliferation</i> , 2019, 52, e12583.	5.3	18
18	Two Novel Long Noncoding RNAs - RP11-296E3.2 and LEF1-AS1 can - Separately Serve as Diagnostic and Prognostic Bio-Markers of Metastasis in Colorectal Cancer. <i>Medical Science Monitor</i> , 2019, 25, 7042-7051.	1.1	14

#	ARTICLE	IF	CITATIONS
19	hPCL3s Promotes Hepatocellular Carcinoma Metastasis by Activating $\beta$ -Catenin Signaling. <i>Cancer Research</i> , 2018, 78, 2536-2549.	0.9	34
20	Chemerin suppresses hepatocellular carcinoma metastasis through CMKLR1-PTEN-Akt axis. <i>British Journal of Cancer</i> , 2018, 118, 1337-1348.	6.4	62
21	PRMT1 Promoted HCC Growth and Metastasis In Vitro and In Vivo via Activating the STAT3 Signalling Pathway. <i>Cellular Physiology and Biochemistry</i> , 2018, 47, 1643-1654.	1.6	33
22	FABP4 suppresses proliferation and invasion of hepatocellular carcinoma cells and predicts a poor prognosis for hepatocellular carcinoma. <i>Cancer Medicine</i> , 2018, 7, 2629-2640.	2.8	55
23	Triosephosphate isomerase 1 suppresses growth, migration and invasion of hepatocellular carcinoma cells. <i>Biochemical and Biophysical Research Communications</i> , 2017, 482, 1048-1053.	2.1	44
24	Iron overload in hereditary tyrosinemia type 1 induces liver injury through the Sp1/Tfr2/hepcidin axis. <i>Journal of Hepatology</i> , 2016, 65, 137-145.	3.7	22
25	Sorafenib enriches epithelial cell adhesion molecule <sup>+</sup> positive tumor initiating cells and exacerbates a subtype of hepatocellular carcinoma through TSC2 $\rightarrow$ AKT cascade. <i>Hepatology</i> , 2015, 62, 1791-1803.	7.3	54
26	Recruitment of Phosphatase PP2A by RACK1 Adaptor Protein Deactivates Transcription Factor IRF3 and Limits Type I Interferon Signaling. <i>Immunity</i> , 2014, 40, 515-529.	14.3	94
27	RACK1 modulates NF- $\kappa$ B activation by interfering with the interaction between TRAF2 and the IKK complex. <i>Cell Research</i> , 2014, 24, 359-371.	12.0	42
28	RACK1 Promotes Non-small-cell Lung Cancer Tumorigenicity through Activating Sonic Hedgehog Signaling Pathway. <i>Journal of Biological Chemistry</i> , 2012, 287, 7845-7858.	3.4	79
29	Cleavage of focal adhesion kinase (FAK) is essential in adipocyte differentiation. <i>Biochemical and Biophysical Research Communications</i> , 2007, 357, 648-654.	2.1	33