

Cristiano De Stefanis

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/430270/publications.pdf>

Version: 2024-02-01

26
papers

1,451
citations

566801

15
h-index

610482

24
g-index

26
all docs

26
docs citations

26
times ranked

2877
citing authors

#	ARTICLE	IF	CITATIONS
1	Gut microbiota profiling of pediatric nonalcoholic fatty liver disease and obese patients unveiled by an integrated meta-omics-based approach. <i>Hepatology</i> , 2017, 65, 451-464.	3.6	572
2	LPS-induced TNF- β factor mediates pro-inflammatory and pro-fibrogenic pattern in non-alcoholic fatty liver disease. <i>Oncotarget</i> , 2015, 6, 41434-41452.	0.8	100
3	The Benefit of Sleeve Gastrectomy in Obese Adolescents on Nonalcoholic Steatohepatitis and Hepatic Fibrosis. <i>Journal of Pediatrics</i> , 2017, 180, 31-37.e2.	0.9	95
4	Association between Serum Atypical Fibroblast Growth Factors 21 and 19 and Pediatric Nonalcoholic Fatty Liver Disease. <i>PLoS ONE</i> , 2013, 8, e67160.	1.1	89
5	Docosahexanoic Acid Plus Vitamin D Treatment Improves Features of NAFLD in Children with Serum Vitamin D Deficiency: Results from a Single Centre Trial. <i>PLoS ONE</i> , 2016, 11, e0168216.	1.1	83
6	AMBRA1 regulates cyclin D to guard S-phase entry and genomic integrity. <i>Nature</i> , 2021, 592, 799-803.	13.7	78
7	Focal adhesion kinase depletion reduces human hepatocellular carcinoma growth by repressing enhancer of zeste homolog 2. <i>Cell Death and Differentiation</i> , 2017, 24, 889-902.	5.0	53
8	Macrophage Activation in Pediatric Nonalcoholic Fatty Liver Disease (NAFLD) Correlates with Hepatic Progenitor Cell Response via Wnt3a Pathway. <i>PLoS ONE</i> , 2016, 11, e0157246.	1.1	50
9	β -Klotho gene variation is associated with liver damage in children with NAFLD. <i>Journal of Hepatology</i> , 2020, 72, 411-419.	1.8	48
10	Altered gut-liver axis and hepatic adiponectin expression in OSAS: novel mediators of liver injury in paediatric non-alcoholic fatty liver. <i>Thorax</i> , 2015, 70, 769-781.	2.7	47
11	EZH2 Down-Regulation Exacerbates Lipid Accumulation and Inflammation in in Vitro and in Vivo NAFLD. <i>International Journal of Molecular Sciences</i> , 2013, 14, 24154-24168.	1.8	44
12	Hepatic farnesoid X receptor protein level and circulating fibroblast growth factor 19 concentration in children with NAFLD. <i>Liver International</i> , 2018, 38, 342-349.	1.9	37
13	Dual IGF1R/IR inhibitors in combination with GD2-CAR T-cells display a potent anti-tumor activity in diffuse midline glioma H3K27M-mutant. <i>Neuro-Oncology</i> , 2022, 24, 1150-1163.	0.6	31
14	Plasma high mobility group box 1 protein reflects fibrosis in pediatric nonalcoholic fatty liver disease. <i>Expert Review of Molecular Diagnostics</i> , 2014, 14, 763-771.	1.5	22
15	The exposure to uteroplacental insufficiency is associated with activation of unfolded protein response in postnatal life. <i>PLoS ONE</i> , 2018, 13, e0198490.	1.1	18
16	The Number of Liver Galectin-3 Positive Cells Is Dually Correlated with NAFLD Severity in Children. <i>International Journal of Molecular Sciences</i> , 2019, 20, 3460.	1.8	16
17	Focal adhesion kinase inhibitor TAE226 combined with Sorafenib slows down hepatocellular carcinoma by multiple epigenetic effects. <i>Journal of Experimental and Clinical Cancer Research</i> , 2021, 40, 364.	3.5	15
18	The Contribution of the Adipose Tissue-Liver Axis in Pediatric Patients with Nonalcoholic Fatty Liver Disease after Laparoscopic Sleeve Gastrectomy. <i>Journal of Pediatrics</i> , 2020, 216, 117-127.e2.	0.9	14

#	ARTICLE	IF	CITATIONS
19	Focal Adhesion Kinase (FAK) Over-Expression and Prognostic Implication in Pediatric Hepatocellular Carcinoma. <i>International Journal of Molecular Sciences</i> , 2020, 21, 5795.	1.8	12
20	HDL cholesterol protects from liver injury in mice with intestinal specific LXRI± activation. <i>Liver International</i> , 2020, 40, 3127-3139.	1.9	8
21	Arginase 1: A potential marker of a common pattern of liver steatosis in HCV and NAFLD children. <i>Journal of Hepatology</i> , 2015, 62, 1207-1208.	1.8	6
22	Expression of insulin-like growth factor I and its receptor in the liver of children with biopsy-proven NAFLD. <i>PLoS ONE</i> , 2018, 13, e0201566.	1.1	6
23	Changes in Total Homocysteine and Glutathione Levels After Laparoscopic Sleeve Gastrectomy in Children with Metabolic-Associated Fatty Liver Disease. <i>Obesity Surgery</i> , 2021, , 1.	1.1	6
24	Retinoids counteract insulin resistance and liver steatosis: What's the potential mechanism?. <i>Hepatology</i> , 2013, 58, 1185-1185.	3.6	1
25	TMOD-14. INNOVATIVE 3D MODEL FOR THE ESTABLISHMENT OF PRIMARY PAEDIATRIC LOW-GRADE GLIOMA (LGG) CULTURES: NEW PLATFORM FOR ADVANCED PRECLINICAL STUDIES OF INNOVATIVE AND IMMUNOTHERAPEUTIC APPROACHES. <i>Neuro-Oncology</i> , 2019, 21, ii123-ii124.	0.6	0
26	IMMU-13. DUAL IGF1R/IR INHIBITOR IN COMBINATION WITH GD2-CAR T-CELLS AS A POTENT THERAPEUTIC STRATEGY FOR H3K27M-MUTANT DIFFUSE MIDLINE GLIOMAS. <i>Neuro-Oncology</i> , 2020, 22, iii362-iii362.	0.6	0