

Seyed Vahid Razavi-Termeh

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/430289/publications.pdf>

Version: 2024-02-01

189
papers

17,472
citations

15001

68
h-index

17891

125
g-index

193
all docs

193
docs citations

193
times ranked

7411
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Spatial modelling of accidents risk caused by driver drowsiness with data mining algorithms. Geocarto International, 2022, 37, 2698-2716. | 1.7 | 14 |
| 2 | Coronavirus disease vulnerability map using a geographic information system (GIS) from 16 April to 16 May 2020. Physics and Chemistry of the Earth, 2022, 126, 103043. | 1.2 | 8 |
| 3 | Comparison of statistical and machine learning approaches in land subsidence modelling. Geocarto International, 2022, 37, 6165-6185. | 1.7 | 5 |
| 4 | Application of machine learning algorithms in hydrology. , 2022, , 585-591. | | 18 |
| 5 | Predicting areas affected by forest fire based on a machine learning algorithm. , 2022, , 351-362. | | 3 |
| 6 | A novel hybrid of support vector regression and metaheuristic algorithms for groundwater spring potential mapping. Science of the Total Environment, 2022, 807, 151055. | 3.9 | 16 |
| 7 | Investigation of water quality and its spatial distribution in the Kor River basin, Fars province, Iran. Environmental Research, 2022, 204, 112294. | 3.7 | 18 |
| 8 | Spatial modeling of land subsidence using machine learning models and statistical methods. Environmental Science and Pollution Research, 2022, 29, 28866-28883. | 2.7 | 17 |
| 9 | A spatially based machine learning algorithm for potential mapping of the hearing senses in an urban environment. Sustainable Cities and Society, 2022, 80, 103675. | 5.1 | 11 |
| 10 | Identification of morphometric features of alluvial fan and basins in predicting the erosion levels using ANN. Environmental Earth Sciences, 2022, 81, 1. | 1.3 | 4 |
| 11 | Multi-hazard spatial modeling via ensembles of machine learning and meta-heuristic techniques. Scientific Reports, 2022, 12, 1451. | 1.6 | 19 |
| 12 | A multi-criteria GIS-based model for wind farm site selection with the least impact on environmental pollution using the OWA-ANP method. Environmental Science and Pollution Research, 2022, 29, 43891-43912. | 2.7 | 14 |
| 13 | The topographic threshold of gully erosion and contributing factors. Natural Hazards, 2022, 112, 2013-2035. | 1.6 | 4 |
| 14 | Digital soil mapping and modeling in Loess-derived soils of Iranian Loess Plateau. Geocarto International, 2022, 37, 11633-11651. | 1.7 | 7 |
| 15 | Spatio-temporal modelling of asthma-prone areas using a machine learning optimized with metaheuristic algorithms. Geocarto International, 2022, 37, 9917-9942. | 1.7 | 6 |
| 16 | Investigating geometrical characteristics of collapsed pipes and the changing role of driving factors. Journal of Environmental Management, 2022, 312, 114910. | 3.8 | 2 |
| 17 | Aquifer vulnerability identification using DRASTIC-LU model modification by fuzzy analytic hierarchy process. Modeling Earth Systems and Environment, 2022, 8, 5365-5380. | 1.9 | 14 |
| 18 | Advanced machine learning algorithms for flood susceptibility modeling " performance comparison: Red Sea, Egypt. Environmental Science and Pollution Research, 2022, 29, 66768-66792. | 2.7 | 8 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Assessment of groundwater vulnerability in an urban area: a comparative study based on DRASTIC, EBF, and LR models. <i>Environmental Science and Pollution Research</i> , 2022, 29, 72908-72928. | 2.7 | 5 |
| 20 | Integration of machine learning algorithms and GIS-based approaches to cutaneous leishmaniasis prevalence risk mapping. <i>International Journal of Applied Earth Observation and Geoinformation</i> , 2022, 112, 102854. | 0.9 | 3 |
| 21 | Landslide susceptibility mapping using machine learning algorithms and comparison of their performance at Abha Basin, Asir Region, Saudi Arabia. <i>Geoscience Frontiers</i> , 2021, 12, 639-655. | 4.3 | 206 |
| 22 | Location-allocation modeling for emergency evacuation planning with GIS and remote sensing: A case study of Northeast Bangladesh. <i>Geoscience Frontiers</i> , 2021, 12, 101095. | 4.3 | 49 |
| 23 | Factors affecting methane emissions in OPEC member countries: does the agricultural production matter?. <i>Environment, Development and Sustainability</i> , 2021, 23, 6734-6748. | 2.7 | 17 |
| 24 | Assessment of land degradation using machine learning techniques: A case of declining rangelands. <i>Land Degradation and Development</i> , 2021, 32, 1452-1466. | 1.8 | 33 |
| 25 | Ecological risk potential assessment of heavy metal contaminated soils in Ophiolitic formations. <i>Environmental Research</i> , 2021, 192, 110305. | 3.7 | 23 |
| 26 | A linear/non-linear hybrid time-series model to investigate the depletion of inland water bodies. <i>Environment, Development and Sustainability</i> , 2021, 23, 10727-10742. | 2.7 | 2 |
| 27 | Spatial modeling of susceptibility to subsidence using machine learning techniques. <i>Stochastic Environmental Research and Risk Assessment</i> , 2021, 35, 1689. | 1.9 | 18 |
| 28 | Asthma-prone areas modeling using a machine learning model. <i>Scientific Reports</i> , 2021, 11, 1912. | 1.6 | 34 |
| 29 | RUSLE model coupled with RS-GIS for soil erosion evaluation compared with T value in Southwest Iran. <i>Arabian Journal of Geosciences</i> , 2021, 14, 1. | 0.6 | 20 |
| 30 | Groundwater recharge potential zonation using an ensemble of machine learning and bivariate statistical models. <i>Scientific Reports</i> , 2021, 11, 5587. | 1.6 | 47 |
| 31 | Spatial and temporal analysis of urban heat island using Landsat satellite images. <i>Environmental Science and Pollution Research</i> , 2021, 28, 41439-41450. | 2.7 | 21 |
| 32 | Field Monitoring-Based and Theoretical Analysis of Baota Mountain Landslide Stability. <i>Advances in Civil Engineering</i> , 2021, 2021, 1-16. | 0.4 | 1 |
| 33 | Land Subsidence Susceptibility Mapping Using Persistent Scatterer SAR Interferometry Technique and Optimized Hybrid Machine Learning Algorithms. <i>Remote Sensing</i> , 2021, 13, 1326. | 1.8 | 40 |
| 34 | Evaluation of multi-hazard map produced using MaxEnt machine learning technique. <i>Scientific Reports</i> , 2021, 11, 6496. | 1.6 | 63 |
| 35 | Morphometry of AFs in upstream and downstream of floods in Gribayegan, Iran. <i>Natural Hazards</i> , 2021, 108, 425-450. | 1.6 | 5 |
| 36 | Prioritization of water erosion-prone sub-watersheds using three ensemble methods in Qareaghaj catchment, southern Iran. <i>Environmental Science and Pollution Research</i> , 2021, 28, 37894-37917. | 2.7 | 14 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Investigation of plant contamination to Ni, Pb, Zn, and Cd and its relationship with spectral reflections. <i>Environmental Science and Pollution Research</i> , 2021, 28, 37830-37842. | 2.7 | 1 |
| 38 | Fire-susceptibility mapping in the natural areas of Iran using new and ensemble data-mining models. <i>Environmental Science and Pollution Research</i> , 2021, 28, 47395-47406. | 2.7 | 18 |
| 39 | Landslide susceptibility assessment and mapping using state-of-the art machine learning techniques. <i>Natural Hazards</i> , 2021, 108, 1291-1316. | 1.6 | 27 |
| 40 | Social networks` analysis of rural stakeholders in watershed management. <i>Environment, Development and Sustainability</i> , 2021, 23, 17535-17557. | 2.7 | 6 |
| 41 | Landslide susceptibility mapping using statistical bivariate models and their hybrid with normalized spatial-correlated scale index and weighted calibrated landslide potential model. <i>Environmental Earth Sciences</i> , 2021, 80, 1. | 1.3 | 27 |
| 42 | A comparative study between dynamic and soft computing models for sediment forecasting. <i>Soft Computing</i> , 2021, 25, 11005-11017. | 2.1 | 11 |
| 43 | Geohazards Susceptibility Assessment along the Upper Indus Basin Using Four Machine Learning and Statistical Models. <i>ISPRS International Journal of Geo-Information</i> , 2021, 10, 315. | 1.4 | 20 |
| 44 | Wildland Fire Susceptibility Mapping Using Support Vector Regression and Adaptive Neuro-Fuzzy Inference System-Based Whale Optimization Algorithm and Simulated Annealing. <i>ISPRS International Journal of Geo-Information</i> , 2021, 10, 382. | 1.4 | 24 |
| 45 | A multi-hazard map-based flooding, gully erosion, forest fires, and earthquakes in Iran. <i>Scientific Reports</i> , 2021, 11, 14889. | 1.6 | 30 |
| 46 | Determining the geographical model and environmental resilience patterns in Iranian provinces. <i>Environmental Monitoring and Assessment</i> , 2021, 193, 524. | 1.3 | 2 |
| 47 | Determining and forecasting drought susceptibility in southwestern Iran using multi-criteria decision-making (MCDM) coupled with CA-Markov model. <i>Science of the Total Environment</i> , 2021, 781, 146703. | 3.9 | 55 |
| 48 | Spatial Modeling of Asthma-Prone Areas Using Remote Sensing and Ensemble Machine Learning Algorithms. <i>Remote Sensing</i> , 2021, 13, 3222. | 1.8 | 14 |
| 49 | COVID-19 Risk Mapping with Considering Socio-Economic Criteria Using Machine Learning Algorithms. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 9657. | 1.2 | 17 |
| 50 | Evaluation of Tree-Based Machine Learning Algorithms for Accident Risk Mapping Caused by Driver Lack of Alertness at a National Scale. <i>Sustainability</i> , 2021, 13, 10239. | 1.6 | 18 |
| 51 | Effects of air pollution in Spatio-temporal modeling of asthma-prone areas using a machine learning model. <i>Environmental Research</i> , 2021, 200, 111344. | 3.7 | 27 |
| 52 | Integrating Landslide Typology with Weighted Frequency Ratio Model for Landslide Susceptibility Mapping: A Case Study from Lanzhou City of Northwestern China. <i>Remote Sensing</i> , 2021, 13, 3623. | 1.8 | 37 |
| 53 | Application of Granger-causality to study the climate change impacts on depletion patterns of inland water bodies. <i>Hydrological Sciences Journal</i> , 2021, 66, 1767-1776. | 1.2 | 5 |
| 54 | Application of stacking hybrid machine learning algorithms in delineating multi-type flooding in Bangladesh. <i>Journal of Environmental Management</i> , 2021, 295, 113086. | 3.8 | 51 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | Spatio-temporal modeling of PM2.5 risk mapping using three machine learning algorithms. <i>Environmental Pollution</i> , 2021, 289, 117859. | 3.7 | 45 |
| 56 | Assessment of Urban Infrastructures Exposed to Flood Using Susceptibility Map and Google Earth Engine. <i>IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing</i> , 2021, 14, 1923-1937. | 2.3 | 19 |
| 57 | Mapping of landslide susceptibility using the combination of neuro-fuzzy inference system (ANFIS), ant colony (ANFIS-ACOR), and differential evolution (ANFIS-DE) models. <i>Bulletin of Engineering Geology and the Environment</i> , 2021, 80, 2045-2067. | 1.6 | 31 |
| 58 | Modeling and Prediction of Habitat Suitability for <i>Ferula gummosa</i> Medicinal Plant in a Mountainous Area. <i>Natural Resources Research</i> , 2021, 30, 4861-4884. | 2.2 | 9 |
| 59 | Habitat potential modelling and mapping of <i>Teucrium polium</i> using machine learning techniques. <i>Environmental Monitoring and Assessment</i> , 2021, 193, 759. | 1.3 | 8 |
| 60 | Soil erosion assessment using RUSLE model and its validation by FR probability model. <i>Geocarto International</i> , 2020, 35, 1750-1768. | 1.7 | 51 |
| 61 | The potential of straw mulch as a nature-based solution for soil erosion in olive plantation treated with glyphosate: A biophysical and socioeconomic assessment. <i>Land Degradation and Development</i> , 2020, 31, 1877-1889. | 1.8 | 44 |
| 62 | Landslide susceptibility mapping using maximum entropy and support vector machine models along the highway corridor, Garhwal Himalaya. <i>Geocarto International</i> , 2020, 35, 168-187. | 1.7 | 70 |
| 63 | Is multi-hazard mapping effective in assessing natural hazards and integrated watershed management?. <i>Geoscience Frontiers</i> , 2020, 11, 1203-1217. | 4.3 | 67 |
| 64 | An assessment of metaheuristic approaches for flood assessment. <i>Journal of Hydrology</i> , 2020, 582, 124536. | 2.3 | 50 |
| 65 | Land-subsidence susceptibility zonation using remote sensing, GIS, and probability models in a Google Earth Engine platform. <i>Environmental Earth Sciences</i> , 2020, 79, 1. | 1.3 | 16 |
| 66 | Improving groundwater potential mapping using metaheuristic approaches. <i>Hydrological Sciences Journal</i> , 2020, 65, 2729-2749. | 1.2 | 31 |
| 67 | A machine learning framework for multi-hazards modeling and mapping in a mountainous area. <i>Scientific Reports</i> , 2020, 10, 12144. | 1.6 | 66 |
| 68 | Comparison of new individual and hybrid machine learning algorithms for modeling and mapping fire hazard: a supplementary analysis of fire hazard in different counties of Golestan Province in Iran. <i>Natural Hazards</i> , 2020, 104, 305-327. | 1.6 | 29 |
| 69 | Predicting non-carcinogenic hazard quotients of heavy metals in pepper (<i>Capsicum annum</i> L.) utilizing electromagnetic waves. <i>Frontiers of Environmental Science and Engineering</i> , 2020, 14, 1. | 3.3 | 8 |
| 70 | A new integrated data mining model to map spatial variation in the susceptibility of land to act as a source of aeolian dust. <i>Environmental Science and Pollution Research</i> , 2020, 27, 42022-42039. | 2.7 | 26 |
| 71 | Assessment of the outbreak risk, mapping and infection behavior of COVID-19: Application of the autoregressive integrated-moving average (ARIMA) and polynomial models. <i>PLoS ONE</i> , 2020, 15, e0236238. | 1.1 | 29 |
| 72 | Assessing the susceptibility of schools to flood events in Iran. <i>Scientific Reports</i> , 2020, 10, 18114. | 1.6 | 17 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 73 | Using Dempster-Shafer theory to model earthquake events. <i>Natural Hazards</i> , 2020, 103, 1943-1959. | 1.6 | 3 |
| 74 | A comparison of machine learning models for the mapping of groundwater spring potential. <i>Environmental Earth Sciences</i> , 2020, 79, 1. | 1.3 | 29 |
| 75 | Ubiquitous GIS-Based Forest Fire Susceptibility Mapping Using Artificial Intelligence Methods. <i>Remote Sensing</i> , 2020, 12, 1689. | 1.8 | 46 |
| 76 | Gully erosion susceptibility mapping using artificial intelligence and statistical models. <i>Geomatics, Natural Hazards and Risk</i> , 2020, 11, 821-844. | 2.0 | 40 |
| 77 | Assessing, mapping, and optimizing the locations of sediment control check dams construction. <i>Science of the Total Environment</i> , 2020, 739, 139954. | 3.9 | 20 |
| 78 | Spatial prediction of landslide susceptibility using hybrid support vector regression (SVR) and the adaptive neuro-fuzzy inference system (ANFIS) with various metaheuristic algorithms. <i>Science of the Total Environment</i> , 2020, 741, 139937. | 3.9 | 113 |
| 79 | Soil loss tolerance in calcareous soils of a semiarid region: evaluation, prediction, and influential parameters. <i>Land Degradation and Development</i> , 2020, 31, 2156-2167. | 1.8 | 29 |
| 80 | A comparative study on machine learning modeling for mass movement susceptibility mapping (a case study) in the Tj ETQq0 0,0 rgBT /Overlock 10 | 1.6 | 24 |
| 81 | Morphometric attributes-based soil erosion susceptibility mapping in Dnyanganga watershed of India using individual and ensemble models. <i>Environmental Earth Sciences</i> , 2020, 79, 1. | 1.3 | 17 |
| 82 | Spatial modeling, risk mapping, change detection, and outbreak trend analysis of coronavirus (COVID-19) in Iran (days between February 19 and June 14, 2020). <i>International Journal of Infectious Diseases</i> , 2020, 98, 90-108. | 1.5 | 94 |
| 83 | Assessing and mapping multi-hazard risk susceptibility using a machine learning technique. <i>Scientific Reports</i> , 2020, 10, 3203. | 1.6 | 126 |
| 84 | Application of learning vector quantization and different machine learning techniques to assessing forest fire influence factors and spatial modelling. <i>Environmental Research</i> , 2020, 184, 109321. | 3.7 | 72 |
| 85 | Evaluation of Recent Advanced Soft Computing Techniques for Gully Erosion Susceptibility Mapping: A Comparative Study. <i>Sensors</i> , 2020, 20, 335. | 2.1 | 33 |
| 86 | Prediction of drainage morphometry using a genetic landscape evolution algorithm. <i>Geocarto International</i> , 2020, , 1-14. | 1.7 | 2 |
| 87 | Using machine learning algorithms to map the groundwater recharge potential zones. <i>Journal of Environmental Management</i> , 2020, 265, 110525. | 3.8 | 52 |
| 88 | Title is missing!. , 2020, 15, e0236238. | | 0 |
| 89 | Title is missing!. , 2020, 15, e0236238. | | 0 |
| 90 | Title is missing!. , 2020, 15, e0236238. | | 0 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 91 | Title is missing!. , 2020, 15, e0236238. | | 0 |
| 92 | Land subsidence susceptibility assessment using random forest machine learning algorithm. Environmental Earth Sciences, 2019, 78, 1. | 1.3 | 80 |
| 93 | Groundwater Potential Mapping Using an Integrated Ensemble of Three Bivariate Statistical Models with Random Forest and Logistic Model Tree Models. Water (Switzerland), 2019, 11, 1596. | 1.2 | 55 |
| 94 | Flood Spatial Modeling in Northern Iran Using Remote Sensing and GIS: A Comparison between Evidential Belief Functions and Its Ensemble with a Multivariate Logistic Regression Model. Remote Sensing, 2019, 11, 1589. | 1.8 | 124 |
| 95 | Multi-hazard probability assessment and mapping in Iran. Science of the Total Environment, 2019, 692, 556-571. | 3.9 | 119 |
| 96 | GIS-based susceptibility assessment of the occurrence of gully headcuts and pipe collapses in a semi-arid environment: Golestan Province, NE Iran. Land Degradation and Development, 2019, 30, 2211-2225. | 1.8 | 26 |
| 97 | Maxent Data Mining Technique and Its Comparison with a Bivariate Statistical Model for Predicting the Potential Distribution of Astragalus Fasciculifolius Boiss. in Fars, Iran. Sustainability, 2019, 11, 3452. | 1.6 | 32 |
| 98 | Comparison analytic network and analytical hierarchical process approaches with feature selection algorithm to predict groundwater quality. Environmental Earth Sciences, 2019, 78, 1. | 1.3 | 5 |
| 99 | Identification of soil erosion-susceptible areas using fuzzy logic and analytical hierarchy process modeling in an agricultural watershed of Burdwan district, India. Environmental Earth Sciences, 2019, 78, 1. | 1.3 | 75 |
| 100 | A Comparative Assessment of Random Forest and k-Nearest Neighbor Classifiers for Gully Erosion Susceptibility Mapping. Water (Switzerland), 2019, 11, 2076. | 1.2 | 75 |
| 101 | SEVUCAS: A Novel GIS-Based Machine Learning Software for Seismic Vulnerability Assessment. Applied Sciences (Switzerland), 2019, 9, 3495. | 1.3 | 42 |
| 102 | Assessment of land subsidence susceptibility in Semnan plain (Iran): a comparison of support vector machine and weights of evidence data mining algorithms. Natural Hazards, 2019, 99, 951-971. | 1.6 | 49 |
| 103 | Predicting Habitat Suitability and Conserving Juniperus spp. Habitat Using SVM and Maximum Entropy Machine Learning Techniques. Water (Switzerland), 2019, 11, 2049. | 1.2 | 31 |
| 104 | Testing a New Ensemble Model Based on SVM and Random Forest in Forest Fire Susceptibility Assessment and Its Mapping in Serbia's Tara National Park. Forests, 2019, 10, 408. | 0.9 | 124 |
| 105 | Evaluation of factors affecting gully headcut location using summary statistics and the maximum entropy model: Golestan Province, NE Iran. Science of the Total Environment, 2019, 677, 281-298. | 3.9 | 36 |
| 106 | Gully erosion susceptibility assessment and management of hazard-prone areas in India using different machine learning algorithms. Science of the Total Environment, 2019, 668, 124-138. | 3.9 | 202 |
| 107 | Spatial prediction of groundwater potentiality using ANFIS ensembled with teaching-learning-based and biogeography-based optimization. Journal of Hydrology, 2019, 572, 435-448. | 2.3 | 150 |
| 108 | PMT: New analytical framework for automated evaluation of geo-environmental modelling approaches. Science of the Total Environment, 2019, 664, 296-311. | 3.9 | 84 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 109 | Assessing the performance of GIS- based machine learning models with different accuracy measures for determining susceptibility to gully erosion. <i>Science of the Total Environment</i> , 2019, 664, 1117-1132. | 3.9 | 137 |
| 110 | Habitat Suitability Mapping of <i>Artemisia aucheri</i> Boiss Based on the GLM Model in R. , 2019, , 213-227. | | 8 |
| 111 | Prioritization of Effective Factors on <i>Zataria multiflora</i> Habitat Suitability and its Spatial Modeling. , 2019, , 411-427. | | 6 |
| 112 | Spatial Modeling of Gully Erosion. , 2019, , 653-669. | | 19 |
| 113 | Producing a Spatially Focused Landslide Susceptibility Map Using an Ensemble of Shannon's Entropy and Fractal Dimension (Case Study: Ziarat Watershed, Iran). , 2019, , 689-732. | | 3 |
| 114 | Gully Erosion Susceptibility Mapping Using Multivariate Adaptive Regression Splines' Replications and Sample Size Scenarios. <i>Water (Switzerland)</i> , 2019, 11, 2319. | 1.2 | 25 |
| 115 | How do data-mining models consider arsenic contamination in sediments and variables importance?. <i>Environmental Monitoring and Assessment</i> , 2019, 191, 777. | 1.3 | 10 |
| 116 | Landslide Susceptibility Mapping Using GIS-Based Data Mining Algorithms. <i>Water (Switzerland)</i> , 2019, 11, 2292. | 1.2 | 40 |
| 117 | Assessment of the importance of gully erosion effective factors using Boruta algorithm and its spatial modeling and mapping using three machine learning algorithms. <i>Geoderma</i> , 2019, 340, 55-69. | 2.3 | 152 |
| 118 | Prioritization of effective factors in the occurrence of land subsidence and its susceptibility mapping using an SVM model and their different kernel functions. <i>Bulletin of Engineering Geology and the Environment</i> , 2019, 78, 4017-4034. | 1.6 | 99 |
| 119 | Effects of urbanization on river morphology of the Talar River, Mazandarn Province, Iran. <i>Geocarto International</i> , 2019, 34, 276-292. | 1.7 | 29 |
| 120 | Soil organic carbon mapping using remote sensing techniques and multivariate regression model. <i>Geocarto International</i> , 2019, 34, 215-226. | 1.7 | 58 |
| 121 | Flood susceptibility mapping using geospatial frequency ratio technique: a case study of Subarnarekha River Basin, India. <i>Modeling Earth Systems and Environment</i> , 2018, 4, 395-408. | 1.9 | 116 |
| 122 | Analysis and evaluation of landslide susceptibility: a review on articles published during 2005-2016 (periods of 2005-2012 and 2013-2016). <i>Arabian Journal of Geosciences</i> , 2018, 11, 1. | 0.6 | 166 |
| 123 | A comparative study of landslide susceptibility maps produced using support vector machine with different kernel functions and entropy data mining models in China. <i>Bulletin of Engineering Geology and the Environment</i> , 2018, 77, 647-664. | 1.6 | 161 |
| 124 | A comparison between ten advanced and soft computing models for groundwater qanat potential assessment in Iran using R and GIS. <i>Theoretical and Applied Climatology</i> , 2018, 131, 967-984. | 1.3 | 127 |
| 125 | Prioritization of landslide conditioning factors and its spatial modeling in Shangnan County, China using GIS-based data mining algorithms. <i>Bulletin of Engineering Geology and the Environment</i> , 2018, 77, 611-629. | 1.6 | 94 |
| 126 | Modeling and assessing the effects of land use changes on runoff generation with the CLUE-s and WetSpa models. <i>Theoretical and Applied Climatology</i> , 2018, 133, 459-471. | 1.3 | 28 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 127 | Spatial assessment of groundwater quality using water quality index and hydrochemical indices in the Kodavanan sub-basin, Tamil Nadu, India. <i>Sustainable Water Resources Management</i> , 2018, 4, 627-641. | 1.0 | 34 |
| 128 | Assessment of floodplain landuse and channel morphology within meandering reach of the Talar River in Iran using GIS and aerial photographs. <i>Geocarto International</i> , 2018, 33, 1367-1380. | 1.7 | 12 |
| 129 | Flood susceptibility mapping using novel ensembles of adaptive neuro fuzzy inference system and metaheuristic algorithms. <i>Science of the Total Environment</i> , 2018, 615, 438-451. | 3.9 | 330 |
| 130 | Erodibility prioritization of sub-watersheds using morphometric parameters analysis and its mapping: A comparison among TOPSIS, VIKOR, SAW, and CF multi-criteria decision making models. <i>Science of the Total Environment</i> , 2018, 613-614, 1385-1400. | 3.9 | 142 |
| 131 | Spatial modelling of gully erosion in Mazandaran Province, northern Iran. <i>Catena</i> , 2018, 161, 1-13. | 2.2 | 155 |
| 132 | Identification of erosion-prone areas using different multi-criteria decision-making techniques and GIS. <i>Geomatics, Natural Hazards and Risk</i> , 2018, 9, 1129-1155. | 2.0 | 74 |
| 133 | A novel hybrid bivariate statistical method entitled FROC for landslide susceptibility assessment. <i>Environmental Earth Sciences</i> , 2018, 77, 1. | 1.3 | 8 |
| 134 | Assessment of Landslide-Prone Areas and Their Zonation Using Logistic Regression, LogitBoost, and Naïve Bayes Machine-Learning Algorithms. <i>Sustainability</i> , 2018, 10, 3697. | 1.6 | 82 |
| 135 | Spatial modelling of gully erosion using evidential belief function, logistic regression, and a new ensemble of evidential belief function and logistic regression algorithm. <i>Land Degradation and Development</i> , 2018, 29, 4035-4049. | 1.8 | 98 |
| 136 | GIS-based gully erosion susceptibility mapping: a comparison among three data-driven models and AHP knowledge-based technique. <i>Environmental Earth Sciences</i> , 2018, 77, 1. | 1.3 | 125 |
| 137 | Spatial Modelling of Gully Erosion Using GIS and R Programming: A Comparison among Three Data Mining Algorithms. <i>Applied Sciences (Switzerland)</i> , 2018, 8, 1369. | 1.3 | 103 |
| 138 | Landslide susceptibility assessment in the Anfu County, China: comparing different statistical and probabilistic models considering the new topo-hydrological factor (HAND). <i>Earth Science Informatics</i> , 2018, 11, 605-622. | 1.6 | 21 |
| 139 | A GIS-based comparative study of Dempster-Shafer, logistic regression and artificial neural network models for landslide susceptibility mapping. <i>Geocarto International</i> , 2017, 32, 367-385. | 1.7 | 143 |
| 140 | Landslide susceptibility assessment in the Uttarakhand area (India) using GIS: a comparison study of prediction capability of naïve bayes, multilayer perceptron neural networks, and functional trees methods. <i>Theoretical and Applied Climatology</i> , 2017, 128, 255-273. | 1.3 | 264 |
| 141 | Identification of Critical Flood Prone Areas in Data-Scarce and Ungauged Regions: A Comparison of Three Data Mining Models. <i>Water Resources Management</i> , 2017, 31, 1473-1487. | 1.9 | 134 |
| 142 | Interplay between river dynamics and international borders: The Hirmand River between Iran and Afghanistan. <i>Science of the Total Environment</i> , 2017, 586, 492-501. | 3.9 | 17 |
| 143 | Spatial Mapping of Groundwater Potential Using Entropy Weighted Linear Aggregate Novel Approach and GIS. <i>Arabian Journal for Science and Engineering</i> , 2017, 42, 1185-1199. | 1.7 | 27 |
| 144 | Performance evaluation of GIS-based new ensemble data mining techniques of adaptive neuro-fuzzy inference system (ANFIS) with genetic algorithm (GA), differential evolution (DE), and particle swarm optimization (PSO) for landslide spatial modelling. <i>Catena</i> , 2017, 157, 310-324. | 2.2 | 267 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 145 | A comparative assessment between linear and quadratic discriminant analyses (LDA-QDA) with frequency ratio and weights-of-evidence models for forest fire susceptibility mapping in China. <i>Arabian Journal of Geosciences</i> , 2017, 10, 1. | 0.6 | 91 |
| 146 | Evaluation of different machine learning models for predicting and mapping the susceptibility of gully erosion. <i>Geomorphology</i> , 2017, 298, 118-137. | 1.1 | 195 |
| 147 | Spatial prediction of landslide susceptibility using an adaptive neuro-fuzzy inference system combined with frequency ratio, generalized additive model, and support vector machine techniques. <i>Geomorphology</i> , 2017, 297, 69-85. | 1.1 | 215 |
| 148 | Performance assessment of individual and ensemble data-mining techniques for gully erosion modeling. <i>Science of the Total Environment</i> , 2017, 609, 764-775. | 3.9 | 258 |
| 149 | GIS-based bivariate statistical techniques for groundwater potential analysis (an example of Iran). <i>Journal of Earth System Science</i> , 2017, 126, 1. | 0.6 | 25 |
| 150 | Landslide spatial modeling: Introducing new ensembles of ANN, MaxEnt, and SVM machine learning techniques. <i>Geoderma</i> , 2017, 305, 314-327. | 2.3 | 280 |
| 151 | Landslide susceptibility modeling in a landslide prone area in Mazandam Province, north of Iran: a comparison between GLM, GAM, MARS, and M-AHP methods. <i>Theoretical and Applied Climatology</i> , 2017, 130, 609-633. | 1.3 | 129 |
| 152 | Evaluating the influence of geo-environmental factors on gully erosion in a semi-arid region of Iran: An integrated framework. <i>Science of the Total Environment</i> , 2017, 579, 913-927. | 3.9 | 152 |
| 153 | Applying different scenarios for landslide spatial modeling using computational intelligence methods. <i>Environmental Earth Sciences</i> , 2017, 76, 1. | 1.3 | 49 |
| 154 | A GIS-based flood susceptibility assessment and its mapping in Iran: a comparison between frequency ratio and weights-of-evidence bivariate statistical models with multi-criteria decision-making technique. <i>Natural Hazards</i> , 2016, 83, 947-987. | 1.6 | 333 |
| 155 | GIS-based multivariate adaptive regression spline and random forest models for groundwater potential mapping in Iran. <i>Environmental Earth Sciences</i> , 2016, 75, 1. | 1.3 | 149 |
| 156 | Flash flood susceptibility analysis and its mapping using different bivariate models in Iran: a comparison between Shannon's entropy, statistical index, and weighting factor models. <i>Environmental Monitoring and Assessment</i> , 2016, 188, 656. | 1.3 | 202 |
| 157 | A comparative assessment of prediction capabilities of modified analytical hierarchy process (M-AHP) and Mamdani fuzzy logic models using Netcad-GIS for forest fire susceptibility mapping. <i>Geomatics, Natural Hazards and Risk</i> , 2016, 7, 861-885. | 2.0 | 127 |
| 158 | GIS-based forest fire susceptibility mapping in Iran: a comparison between evidential belief function and binary logistic regression models. <i>Scandinavian Journal of Forest Research</i> , 2016, 31, 80-98. | 0.5 | 99 |
| 159 | Random forests and evidential belief function-based landslide susceptibility assessment in Western Mazandaran Province, Iran. <i>Environmental Earth Sciences</i> , 2016, 75, 1. | 1.3 | 245 |
| 160 | Fractal analysis of rainfall-induced landslide and debris flow spread distribution in the Chenyulan Creek Basin, Taiwan. <i>Journal of Earth Science (Wuhan, China)</i> , 2016, 27, 151-159. | 1.1 | 13 |
| 161 | Gully erosion susceptibility mapping: the role of GIS-based bivariate statistical models and their comparison. <i>Natural Hazards</i> , 2016, 82, 1231-1258. | 1.6 | 189 |
| 162 | GIS-based landslide spatial modeling in Ganzhou City, China. <i>Arabian Journal of Geosciences</i> , 2016, 9, 1. | 0.6 | 123 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 163 | Assessment and comparison of combined bivariate and AHP models with logistic regression for landslide susceptibility mapping in the Chaharmahal-e-Bakhtiari Province, Iran. <i>Arabian Journal of Geosciences</i> , 2016, 9, 1. | 0.6 | 45 |
| 164 | GIS-based groundwater potential mapping using boosted regression tree, classification and regression tree, and random forest machine learning models in Iran. <i>Environmental Monitoring and Assessment</i> , 2016, 188, 44. | 1.3 | 489 |
| 165 | Investigation of general indicators influencing on forest fire and its susceptibility modeling using different data mining techniques. <i>Ecological Indicators</i> , 2016, 64, 72-84. | 2.6 | 178 |
| 166 | Application of GIS-based data driven random forest and maximum entropy models for groundwater potential mapping: A case study at Mehran Region, Iran. <i>Catena</i> , 2016, 137, 360-372. | 2.2 | 408 |
| 167 | Landslide susceptibility mapping using random forest, boosted regression tree, classification and regression tree, and general linear models and comparison of their performance at Wadi Tayyah Basin, Asir Region, Saudi Arabia. <i>Landslides</i> , 2016, 13, 839-856. | 2.7 | 530 |
| 168 | Assessment of a spatial multi-criteria evaluation to site selection underground dams in the Alborz Province, Iran. <i>Geocarto International</i> , 2016, 31, 628-646. | 1.7 | 43 |
| 169 | Flood susceptibility mapping using frequency ratio and weights-of-evidence models in the Golastan Province, Iran. <i>Geocarto International</i> , 2016, 31, 42-70. | 1.7 | 376 |
| 170 | Landslide susceptibility maps using different probabilistic and bivariate statistical models and comparison of their performance at Wadi Itwad Basin, Asir Region, Saudi Arabia. <i>Bulletin of Engineering Geology and the Environment</i> , 2016, 75, 63-87. | 1.6 | 68 |
| 171 | Landslide susceptibility assessment at Wadi Jawrah Basin, Jizan region, Saudi Arabia using two bivariate models in GIS. <i>Geosciences Journal</i> , 2015, 19, 449-469. | 0.6 | 58 |
| 172 | An integrated artificial neural network model for the landslide susceptibility assessment of Osado Island, Japan. <i>Natural Hazards</i> , 2015, 78, 1749-1776. | 1.6 | 182 |
| 173 | Automatic detection of sinkhole collapses at finer resolutions using a multi-component remote sensing approach. <i>Natural Hazards</i> , 2015, 78, 1021-1044. | 1.6 | 16 |
| 174 | Application of analytical hierarchy process, frequency ratio, and certainty factor models for groundwater potential mapping using GIS. <i>Earth Science Informatics</i> , 2015, 8, 867-883. | 1.6 | 389 |
| 175 | Accuracy assessment of land cover/land use classifiers in dry and humid areas of Iran. <i>Environmental Monitoring and Assessment</i> , 2015, 187, 641. | 1.3 | 30 |
| 176 | A Comparative Assessment Between Three Machine Learning Models and Their Performance Comparison by Bivariate and Multivariate Statistical Methods in Groundwater Potential Mapping. <i>Water Resources Management</i> , 2015, 29, 5217-5236. | 1.9 | 213 |
| 177 | Groundwater qanat potential mapping using frequency ratio and Shannon's entropy models in the Moghan watershed, Iran. <i>Earth Science Informatics</i> , 2015, 8, 171-186. | 1.6 | 259 |
| 178 | Groundwater potential mapping at Kurdistan region of Iran using analytic hierarchy process and GIS. <i>Arabian Journal of Geosciences</i> , 2015, 8, 7059-7071. | 0.6 | 417 |
| 179 | Assessment of a data-driven evidential belief function model and GIS for groundwater potential mapping in the Koohrang Watershed, Iran. <i>Geocarto International</i> , 2015, 30, 662-685. | 1.7 | 139 |
| 180 | Application of frequency ratio, statistical index, and weights-of-evidence models and their comparison in landslide susceptibility mapping in Central Nepal Himalaya. <i>Arabian Journal of Geosciences</i> , 2014, 7, 725-742. | 0.6 | 366 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 181 | GIS-based groundwater spring potential assessment and mapping in the Birjand Township, southern Khorasan Province, Iran. <i>Hydrogeology Journal</i> , 2014, 22, 643-662. | 0.9 | 240 |
| 182 | Landslide susceptibility mapping at Vaz Watershed (Iran) using an artificial neural network model: a comparison between multilayer perceptron (MLP) and radial basic function (RBF) algorithms. <i>Arabian Journal of Geosciences</i> , 2013, 6, 2873-2888. | 0.6 | 315 |
| 183 | Landslide susceptibility mapping using support vector machine and GIS at the Golestan Province, Iran. <i>Journal of Earth System Science</i> , 2013, 122, 349-369. | 0.6 | 278 |
| 184 | Application of weights-of-evidence and certainty factor models and their comparison in landslide susceptibility mapping at Haraz watershed, Iran. <i>Arabian Journal of Geosciences</i> , 2013, 6, 2351-2365. | 0.6 | 261 |
| 185 | Landslide susceptibility mapping using certainty factor, index of entropy and logistic regression models in GIS and their comparison at Muglingâ€™Narayanghat road section in Nepal Himalaya. <i>Natural Hazards</i> , 2013, 65, 135-165. | 1.6 | 559 |
| 186 | Application of fuzzy logic and analytical hierarchy process (AHP) to landslide susceptibility mapping at Haraz watershed, Iran. <i>Natural Hazards</i> , 2012, 63, 965-996. | 1.6 | 758 |
| 187 | Landslide susceptibility mapping at Golestan Province, Iran: A comparison between frequency ratio, Dempsterâ€™Shafer, and weights-of-evidence models. <i>Journal of Asian Earth Sciences</i> , 2012, 61, 221-236. | 1.0 | 378 |
| 188 | Landslide susceptibility mapping using index of entropy and conditional probability models in GIS: Safarood Basin, Iran. <i>Catena</i> , 2012, 97, 71-84. | 2.2 | 400 |
| 189 | Remote Sensing Data Derived Parameters and its Use in Landslide Susceptibility Assessment Using Shannonâ€™s Entropy and GIS. <i>Applied Mechanics and Materials</i> , 0, 225, 486-491. | 0.2 | 51 |