Kai Pun Wong

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4303391/publications.pdf

Version: 2024-02-01

		1163117	1474206
11	316	8	9
papers	citations	h-index	g-index
11	11	11	518
all docs	docs citations	times ranked	citing authors

#	Article	IF	CITATIONS
1	Non-invasive recording during intra-operative neuro-monitoring in thyroid surgery: omitting surface electrode and oversize endotracheal tube. Gland Surgery, 2019, 8, 312-314.	1.1	1
2	Role of surgery in pancreatic neuroendocrine tumor. Gland Surgery, 2018, 7, 36-41.	1.1	20
3	Effect of Thyroid Remnant Volume on the Risk of Hypothyroidism After Hemithyroidectomy: A Prospective Study. Annals of Surgical Oncology, 2017, 24, 1525-1532.	1.5	38
4	Systematic review and meta-analysis on intra-operative neuro-monitoring in high-risk thyroidectomy. International Journal of Surgery, 2017, 38, 21-30.	2.7	100
5	Indocyanine green fluorescence angiography for quantitative evaluation of in situ parathyroid gland perfusion and function after total thyroidectomy. Surgery, 2017, 161, 87-95.	1.9	77
6	The Association of Discolored Parathyroid Glands and Hypoparathyroidism Following Total Thyroidectomy. World Journal of Surgery, 2016, 40, 1611-1617.	1.6	26
7	Evaluating the Efficacy of Primary Treatment for Graves' Disease Complicated by Thyrotoxic Periodic Paralysis. International Journal of Endocrinology, 2014, 2014, 1-6.	1.5	10
8	A systematic review and meta-analysis comparing outcomes between robotic-assisted thyroidectomy and non-robotic endoscopic thyroidectomy. Journal of Surgical Research, 2014, 191, 389-398.	1.6	28
9	The pros and cons of routine central compartment neck dissection for clinically nodal negative (cN0) papillary thyroid cancer. Gland Surgery, 2013, 2, 186-95.	1.1	16
10	Impact of Routine Unilateral Central Neck Dissection on Preablative and Postablative Stimulated Thyroglobulin Levels after Total Thyroidectomy in Papillary Thyroid Carcinoma. Indian Journal of Surgical Oncology, 2012, 3, 191-198.	0.7	0
11	Significance of Metastatic Lymph Node Ratio on Stimulated Thyroglobulin Levels in Papillary Thyroid Carcinoma after Prophylactic Unilateral Central Neck Dissection. Indian Journal of Surgical Oncology, 2012, 3, 199-205.	0.7	О