

Sten Hammarström

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4315866/publications.pdf>

Version: 2024-02-01

16
papers

333
citations

933447

10
h-index

940533

16
g-index

17
all docs

17
docs citations

17
times ranked

510
citing authors

#	ARTICLE	IF	CITATIONS
1	<i>Lachnoanaerobaculum</i> gen. nov., a new genus in the Lachnospiraceae : characterization of <i>Lachnoanaerobaculum umeaense</i> gen. nov., sp. nov., isolated from the human small intestine, and <i>Lachnoanaerobaculum orale</i> sp. nov., isolated from saliva, and reclassification of <i>Eubacterium saburreum</i> (PrÄ©vot 1966) Holdeman and Moore 1970 as <i>Lachnoanaerobaculum saburreum</i> comb. nov.. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2012, 62, 2685-2690.	1.7	57
2	Ectopic expression of the chemokine CXCL17 in colon cancer cells. <i>British Journal of Cancer</i> , 2016, 114, 697-703.	6.4	43
3	Immunopathology of childhood celiac diseaseâ€”Key role of intestinal epithelial cells. <i>PLoS ONE</i> , 2017, 12, e0185025.	2.5	41
4	Noncontaminated Dietary Oats May Hamper Normalization of the Intestinal Immune Status in Childhood Celiac Disease. <i>Clinical and Translational Gastroenterology</i> , 2014, 5, e58.	2.5	31
5	<i>Prevotella jejuni</i> sp. nov., isolated from the small intestine of a child with coeliac disease. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2013, 63, 4218-4223.	1.7	29
6	Lymph node CEA and MUC2 mRNA as useful predictors of outcome in colorectal cancer. <i>International Journal of Cancer</i> , 2012, 130, 1833-1843.	5.1	23
7	Utility of G protein-coupled receptor 35 expression for predicting outcome in colon cancer. <i>Tumor Biology</i> , 2019, 41, 101042831985888.	1.8	23
8	The Chemokine CXCL16 Is a New Biomarker for Lymph Node Analysis of Colon Cancer Outcome. <i>International Journal of Molecular Sciences</i> , 2019, 20, 5793.	4.1	17
9	Lymph node CXCL17 messenger RNA: A new prognostic biomarker for colon cancer. <i>Tumor Biology</i> , 2018, 40, 101042831879925.	1.8	15
10	Lipids are a constitutive component of cytolytic granules. <i>Histochemistry and Cell Biology</i> , 2000, 114, 167-171.	1.7	12
11	Clinical Significance of Stem Cell Biomarkers EpCAM, LGR5 and LGR4 mRNA Levels in Lymph Nodes of Colon Cancer Patients. <i>International Journal of Molecular Sciences</i> , 2022, 23, 403.	4.1	12
12	Allocating colorectal cancer patients to different risk categories by using a five-biomarker mRNA combination in lymph node analysis. <i>PLoS ONE</i> , 2020, 15, e0229007.	2.5	11
13	Evaluating macrophage migration inhibitory factor 1 expression as a prognostic biomarker in colon cancer. <i>Tumor Biology</i> , 2020, 42, 101042832092452.	1.8	9
14	The myeloid cell biomarker EMR1 is ectopically expressed in colon cancer. <i>Tumor Biology</i> , 2021, 43, 209-223.	1.8	4
15	Prognostic Significance of GPR55 mRNA Expression in Colon Cancer. <i>International Journal of Molecular Sciences</i> , 2022, 23, 4556.	4.1	3
16	Biomarker mRNAs as prognostic tools for lymph node analysis in colorectal cancer. <i>Biomarkers in Medicine</i> , 2019, 13, 801-803.	1.4	2