

Jian Cui

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/4318454/jian-cui-publications-by-year.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

88
papers

1,710
citations

25
h-index

40
g-index

105
ext. papers

1,898
ext. citations

2.8
avg, IF

4.28
L-index

#	Paper	IF	Citations
88	Moderate whole body heating attenuates the exercise pressor reflex responses in older humans. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2021 , 320, R757-R769	3.2	0
87	Sympathetic activation due to limb venous distension is preserved during muscle metaboreceptor stimulation. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2021 , 321, R21-R28	3.2	1
86	Renal medullary oxygenation decreases with lower body negative pressure in healthy young adults. <i>Journal of Applied Physiology</i> , 2021 , 130, 48-56	3.7	
85	Acute effects of sublingual nitroglycerin on cardiovagal and sympathetic baroreflex sensitivity. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2021 , 321, R525-R536	3.2	1
84	Lower-limb venous distension reflex and orthostatic tolerance in young healthy humans. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2020 , 319, R142-R147	3.2	1
83	Systemic and regional hemodynamic response to activation of the exercise pressor reflex in patients with peripheral artery disease. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2020 , 318, H916-H924	5.2	3
82	Reliability of Skin Blood Flow Measurement with Multiple Laser Doppler Probes. <i>FASEB Journal</i> , 2020 , 34, 1-1	0.9	
81	Sympathetic Activation to Limb Venous Distension is Preserved during Muscle Metaboreceptor Stimulation. <i>FASEB Journal</i> , 2020 , 34, 1-1	0.9	
80	Habitual cigarette smoking raises pressor responses to spontaneous bursts of muscle sympathetic nerve activity. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2019 , 317, R280-R288	3.2	4
79	Hemodynamic Response to Activation of the Exercise Pressor Reflex during Dynamic Plantar Flexion in Peripheral Arterial Disease Patients. <i>FASEB Journal</i> , 2019 , 33, 540.1	0.9	
78	Diastolic Function at Rest and During Handgrip is Impaired in Patients with Peripheral Arterial Disease. <i>FASEB Journal</i> , 2019 , 33, 828.6	0.9	
77	The Baroreflex Control of Heart Rate is Impaired in Patients with Peripheral Arterial Disease. <i>FASEB Journal</i> , 2019 , 33, 746.3	0.9	2
76	Age and sex differences in sympathetic and hemodynamic responses to hypoxia and cold pressor test. <i>Physiological Reports</i> , 2019 , 7, e13988	2.6	11
75	Sympathetic responses induced by radiofrequency catheter ablation of atrial fibrillation. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2019 , 316, H476-H484	5.2	2
74	Renal Medullary Oxygenation Decreases in a Dose-Dependent Manner with Graded Lower Body Negative Pressure in Healthy Young Adults. <i>FASEB Journal</i> , 2018 , 32, 621.9	0.9	
73	Autonomic Responses during Atrial Fibrillation Ablation. <i>FASEB Journal</i> , 2018 , 32, 596.2	0.9	
72	Muscle sympathetic nerve activity response to heat stress is attenuated in chronic heart failure patients. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2017 , 312, R873-R882	3.2	7

71	Purinergic P2X Receptors and Heightened Exercise Pressor Reflex in Peripheral Artery Disease. <i>Internal Medicine Review (Washington, D C: Online)</i> , 2016 , 2,	2	1
70	Whole body heat stress attenuates the pressure response to muscle metaboreceptor stimulation in humans. <i>Journal of Applied Physiology</i> , 2016 , 121, 1178-1186	3.7	2
69	Seasonal variation in muscle sympathetic nerve activity. <i>Physiological Reports</i> , 2015 , 3, e12492	2.6	19
68	Limb suction evoked during arterial occlusion causes systemic sympathetic activity in humans. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2015 , 309, R482-8	3.2	5
67	Whole-Body Heat Stress Alters the Exercise Pressor Reflex in Humans. <i>FASEB Journal</i> , 2015 , 29, 827.2	0.9	
66	Role of Adenosine Receptors in Evoking Venous Distension Reflex in Humans. <i>FASEB Journal</i> , 2015 , 29, 649.4	0.9	
65	Blood Pressure Responses to Muscle Sympathetic Nerve Activity are Accentuated in Smokers. <i>FASEB Journal</i> , 2015 , 29, 830.2	0.9	
64	Cyclooxygenase Inhibition Attenuates the Muscle Sympathetic Nerve Activity Responses to Venous Distension in Humans. <i>FASEB Journal</i> , 2015 , 29, 649.5	0.9	
63	Cardiovascular responses to heat stress in chronic heart failure. <i>Current Heart Failure Reports</i> , 2014 , 11, 139-45	2.8	21
62	Increased pressure gradient across veins in human limbs induces sympathetic activation (1170.1). <i>FASEB Journal</i> , 2014 , 28, 1170.1	0.9	
61	Chronic heart failure does not attenuate the total activity of sympathetic outflow to skin during whole-body heating. <i>Circulation: Heart Failure</i> , 2013 , 6, 271-8	7.6	25
60	Distension of central great vein decreases sympathetic outflow in humans. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2013 , 305, H378-85	5.2	16
59	Effect of oxidative stress on sympathetic and renal vascular responses to ischemic exercise. <i>Physiological Reports</i> , 2013 , 1,	2.6	10
58	Renal vasoconstriction is augmented during exercise in patients with peripheral arterial disease. <i>Physiological Reports</i> , 2013 , 1, e00154	2.6	26
57	Tactile stimulation of the oropharynx elicits sympathoexcitation in conscious humans. <i>Journal of Applied Physiology</i> , 2013 , 115, 71-7	3.7	7
56	Distension of central great vein decreases sympathetic outflow in humans. <i>FASEB Journal</i> , 2013 , 27, 1118.6		
55	Oxidative stress contributes to the augmented exercise pressor reflex in peripheral arterial disease patients. <i>Journal of Physiology</i> , 2012 , 590, 6237-46	3.9	74
54	Human sympathetic outflows to skin and muscle target organs fluctuate concordantly over a wide range of time-varying frequencies. <i>Journal of Physiology</i> , 2012 , 590, 363-75	3.9	24

53	Limb venous distension evokes sympathetic activation via stimulation of the limb afferents in humans. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2012 , 303, H457-63	5.2	27
52	Negative pressure on an occluded limb induces sympathetic activation. <i>FASEB Journal</i> , 2012 , 26, 1091.1	0.9	1
51	Intravenous ascorbic acid does not block the pressor or sympathetic nerve response to ischemic fatiguing rhythmic handgrip exercise. <i>FASEB Journal</i> , 2012 , 26, 893.15	0.9	
50	Sympathetic response to fatiguing handgrip and muscle metaboreflex activation is attenuated in smokers compared to non-smokers. <i>FASEB Journal</i> , 2012 , 26, 1087.10	0.9	1
49	Effect of P2 receptor blockade with pyridoxine on sympathetic response to exercise pressor reflex in humans. <i>Journal of Physiology</i> , 2011 , 589, 685-95	3.9	21
48	Muscle sympathetic responses during orthostasis in heat-stressed individuals. <i>Clinical Autonomic Research</i> , 2011 , 21, 381-7	4.3	16
47	Sympathetic and cardiovascular responses to venous distension in an occluded limb. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2011 , 301, R1831-7	3.2	20
46	Heat stress attenuates the increase in arterial blood pressure during the cold pressor test. <i>Journal of Applied Physiology</i> , 2010 , 109, 1354-9	3.7	19
45	Local adenosine receptor blockade accentuates the sympathetic responses to fatiguing exercise. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2010 , 298, H2130-7	5.2	11
44	P2X receptor blockade attenuates the sympathetic response to exercise in humans. <i>FASEB Journal</i> , 2010 , 24, 807.5	0.9	
43	Whole body heat stress attenuates baroreflex control of muscle sympathetic nerve activity during postexercise muscle ischemia. <i>Journal of Applied Physiology</i> , 2009 , 106, 1125-31	3.7	12
42	Dynamic cerebral autoregulation during passive heat stress in humans. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2009 , 296, R1598-605	3.2	38
41	Sympathetic responses during saline infusion into the veins of an occluded limb. <i>Journal of Physiology</i> , 2009 , 587, 3619-28	3.9	22
40	Sustained impairments in cutaneous vasodilation and sweating in grafted skin following long-term recovery. <i>Journal of Burn Care and Research</i> , 2009 , 30, 675-85	0.8	33
39	Adenosine receptor blockade accentuates the responses of muscle sympathetic nerve activity to fatiguing exercise. <i>FASEB Journal</i> , 2009 , 23, 608.5	0.9	
38	Local prostaglandin blockade attenuates muscle mechanoreflex-mediated renal vasoconstriction during muscle stretch in humans. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2008 , 294, H2184-90	5.2	17
37	Effects of muscle metabolites on responses of muscle sympathetic nerve activity to mechanoreceptor(s) stimulation in healthy humans. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2008 , 294, R458-66	3.2	39
36	Cutaneous vasoconstriction during whole-body and local cooling in grafted skin five to nine months postsurgery. <i>Journal of Burn Care and Research</i> , 2008 , 29, 36-41	0.8	6

35	Commentary on viewpoint: the human cutaneous circulation as a model of generalized microvascular function. <i>Journal of Applied Physiology</i> , 2008 , 105, 386; author reply 389	3.7	1
34	Cyclooxygenase inhibition attenuates sympathetic responses to muscle stretch in humans. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2008 , 294, H2693-700	5.2	24
33	Cyclooxygenase inhibition attenuates muscle sympathetic nerve responses to passive muscle stretch. <i>FASEB Journal</i> , 2008 , 22, 957.5	0.9	
32	Baroreflex control of muscle sympathetic nerve activity during skin surface cooling. <i>Journal of Applied Physiology</i> , 2007 , 103, 1284-9	3.7	24
31	Endogenous nitric oxide attenuates neutrally mediated cutaneous vasoconstriction. <i>Journal of Physiology</i> , 2007 , 585, 627-34	3.9	41
30	The role of the cyclooxygenase products in evoking sympathetic activation in exercise. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2007 , 293, H1861-8	5.2	25
29	Skin grafting impairs postsynaptic cutaneous vasodilator and sweating responses. <i>Journal of Burn Care and Research</i> , 2007 , 28, 435-41	0.8	25
28	Impaired cutaneous vasodilation and sweating in grafted skin during whole-body heating. <i>Journal of Burn Care and Research</i> , 2007 , 28, 427-34	0.8	34
27	Endogenous NO decreases cutaneous vasoconstrictor responsiveness during lower-body negative pressure (LBNP) in the heat stressed individual. <i>FASEB Journal</i> , 2007 , 21, A1298	0.9	
26	Cutaneous Vasodilation and Sweating In Grafted Skin During Heat Stress 59 Months Post-Surgery: A 1-Year Follow-up. <i>FASEB Journal</i> , 2007 , 21, A1312	0.9	
25	Muscle metabolites accentuate muscle sympathetic nerve activity responses to passive muscle stretch. <i>FASEB Journal</i> , 2007 , 21, A569	0.9	
24	Effects of local prostaglandin blockade on renal vasoconstriction during muscle stretch. <i>FASEB Journal</i> , 2007 , 21, A568	0.9	
23	Heat stress attenuates increases in arterial blood pressure during a cold pressor test (CPT). <i>FASEB Journal</i> , 2007 , 21, A563	0.9	
22	Effects of systemic hypoxia and lower body negative pressure on heart rate variability and transfer function gain in humans. <i>FASEB Journal</i> , 2007 , 21, A564	0.9	
21	Heat stress decreases baroreflex sensitivity during muscle metaboreceptor stimulation. <i>FASEB Journal</i> , 2007 , 21, A571	0.9	
20	Skin blood flow influences near-infrared spectroscopy-derived measurements of tissue oxygenation during heat stress. <i>Journal of Applied Physiology</i> , 2006 , 100, 221-4	3.7	129
19	Spectral characteristics of skin sympathetic nerve activity in heat-stressed humans. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2006 , 290, H1601-9	5.2	45
18	Heat stress reduces cerebral blood velocity and markedly impairs orthostatic tolerance in humans. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2006 , 291, R1443-8	3.2	118

17	Neurally mediated vasoconstriction is capable of decreasing skin blood flow during orthostasis in the heat-stressed human. <i>Journal of Physiology</i> , 2006 , 575, 953-9	3.9	32
16	Muscle sympathetic nerve activity responses to dynamic passive muscle stretch in humans. <i>Journal of Physiology</i> , 2006 , 576, 625-34	3.9	83
15	Mechanisms of cutaneous vasoconstriction during orthostasis in heat stressed individuals. <i>Japanese Journal of Physical Fitness and Sports Medicine</i> , 2006 , 55, 20-20	0.1	
14	Muscle sympathetic nerve activity responses to dynamic passive muscle stretch.. <i>FASEB Journal</i> , 2006 , 20, A768	0.9	
13	Mean body temperature does not modulate eccrine sweat rate during upright tilt. <i>Journal of Applied Physiology</i> , 2005 , 98, 1207-12	3.7	31
12	Effect of skin surface cooling on central venous pressure during orthostatic challenge. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2005 , 289, H2429-33	5.2	44
11	Effects of heat stress on thermoregulatory responses in congestive heart failure patients. <i>Circulation</i> , 2005 , 112, 2286-92	16.7	83
10	Spectral analysis of muscle sympathetic nerve activity in heat-stressed humans. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2004 , 286, H1101-6	5.2	31
9	Active recovery attenuates the fall in sweat rate but not cutaneous vascular conductance after supine exercise. <i>Journal of Applied Physiology</i> , 2004 , 96, 668-73	3.7	28
8	Orthostatic challenge does not alter skin sympathetic nerve activity in heat-stressed humans. <i>Autonomic Neuroscience: Basic and Clinical</i> , 2004 , 116, 54-61	2.4	54
7	Muscle sympathetic nerve activity during lower body negative pressure is accentuated in heat-stressed humans. <i>Journal of Applied Physiology</i> , 2004 , 96, 2103-8	3.7	43
6	Exercise throughout 6 degrees head-down tilt bed rest preserves thermoregulatory responses. <i>Journal of Applied Physiology</i> , 2003 , 95, 1817-23	3.7	19
5	Nitric oxide synthase inhibition does not affect regulation of muscle sympathetic nerve activity during head-up tilt. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2003 , 285, H2105-10	5.2	8
4	Phenylephrine-induced elevations in arterial blood pressure are attenuated in heat-stressed humans. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2002 , 283, R1221-6	3.2	35
3	Baroreflex modulation of sympathetic nerve activity to muscle in heat-stressed humans. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2002 , 282, R252-8	3.2	48
2	Baroreflex modulation of muscle sympathetic nerve activity during cold pressor test in humans. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2002 , 282, H1717-23	5.2	77
1	Absence of arterial baroreflex modulation of skin sympathetic activity and sweat rate during whole-body heating in humans. <i>Journal of Physiology</i> , 2001 , 536, 615-23	3.9	81