

# Stefanie Endesfelder

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4325650/publications.pdf>

Version: 2024-02-01

54  
papers

1,765  
citations

279487

23  
h-index

276539

41  
g-index

56  
all docs

56  
docs citations

56  
times ranked

2309  
citing authors

#	ARTICLE	IF	CITATIONS
1	Dexmedetomidine Restores Autophagic Flux, Modulates Associated microRNAs and the Cholinergic Anti-inflammatory Pathway upon LPS-Treatment in Rats. <i>Journal of NeuroImmune Pharmacology</i> , 2022, 17, 261-276.	2.1	9
2	Adaptation of the Oxygen Sensing System during Lung Development. <i>Oxidative Medicine and Cellular Longevity</i> , 2022, 2022, 1-15.	1.9	3
3	Perinatal Hyperoxia and Developmental Consequences on the Lung-Brain Axis. <i>Oxidative Medicine and Cellular Longevity</i> , 2022, 2022, 1-17.	1.9	15
4	The Conflicting Role of Caffeine Supplementation on Hyperoxia-Induced Injury on the Cerebellar Granular Cell Neurogenesis of Newborn Rats. <i>Oxidative Medicine and Cellular Longevity</i> , 2022, 2022, 1-21.	1.9	5
5	Neonatal Oxidative Stress Impairs Cortical Synapse Formation and GABA Homeostasis in Parvalbumin-Expressing Interneurons. <i>Oxidative Medicine and Cellular Longevity</i> , 2022, 2022, 1-12.	1.9	9
6	In vitro P38MAPK inhibition in aged astrocytes decreases reactive astrocytes, inflammation and increases nutritive capacity after oxygen-glucose deprivation. <i>Aging</i> , 2021, 13, 6346-6358.	1.4	6
7	GABAB Receptor-Mediated Impairment of Intermediate Progenitor Maturation During Postnatal Hippocampal Neurogenesis of Newborn Rats. <i>Frontiers in Cellular Neuroscience</i> , 2021, 15, 651072.	1.8	16
8	Reduction of cortical parvalbumin-expressing GABAergic interneurons in a rodent hyperoxia model of preterm birth brain injury with deficits in social behavior and cognition. <i>Development (Cambridge)</i> , 2021, 148, .	1.2	7
9	Paracetamol (Acetaminophen) and the Developing Brain. <i>International Journal of Molecular Sciences</i> , 2021, 22, 11156.	1.8	15
10	Postnatal myelination of the immature rat cingulum is regulated by GABAB receptor activity. <i>Developmental Neurobiology</i> , 2021, , .	1.5	9
11	Repetitive Erythropoietin Treatment Improves Long-Term Neurocognitive Outcome by Attenuating Hyperoxia-Induced Hypomyelination in the Developing Brain. <i>Frontiers in Neurology</i> , 2020, 11, 804.	1.1	14
12	Prevention of Oxygen-Induced Inflammatory Lung Injury by Caffeine in Neonatal Rats. <i>Oxidative Medicine and Cellular Longevity</i> , 2020, 2020, 1-19.	1.9	29
13	Transient Improvement of Cerebellar Oligodendroglial Development in a Neonatal Hyperoxia Model by PDGFA Treatment. <i>Developmental Neurobiology</i> , 2019, 79, 222-235.	1.5	6
14	Vascular endothelial growth factor polymorphism rs2010963 status does not affect patent ductus arteriosus incidence or cyclooxygenase inhibitor treatment success in preterm infants. <i>Cardiology in the Young</i> , 2019, 29, 893-897.	0.4	2
15	Antioxidative effects of caffeine in a hyperoxia-based rat model of bronchopulmonary dysplasia. <i>Respiratory Research</i> , 2019, 20, 88.	1.4	39
16	Neonatal Hyperoxia Perturbs Neuronal Development in the Cerebellum. <i>Molecular Neurobiology</i> , 2018, 55, 3901-3915.	1.9	28
17	Caffeine Protects Against Anticonvulsant-Induced Impaired Neurogenesis in the Developing Rat Brain. <i>Neurotoxicity Research</i> , 2018, 34, 173-187.	1.3	19
18	Physostigmine Restores Impaired Autophagy in the Rat Hippocampus after Surgery Stress and LPS Treatment. <i>Journal of NeuroImmune Pharmacology</i> , 2018, 13, 383-395.	2.1	6

#	ARTICLE	IF	CITATIONS
19	Oxygen impairs oligodendroglial development via oxidative stress and reduced expression of HIF-1 $\alpha$ . Scientific Reports, 2017, 7, 43000.	1.6	13
20	Caffeine Protects Against Anticonvulsant-Induced Neurotoxicity in the Developing Rat Brain. Neurotoxicity Research, 2017, 32, 460-472.	1.3	15
21	Dexmedetomidine Prevents Lipopolysaccharide-Induced MicroRNA Expression in the Adult Rat Brain. International Journal of Molecular Sciences, 2017, 18, 1830.	1.8	28
22	Neuroprotection by Caffeine in Hyperoxia-Induced Neonatal Brain Injury. International Journal of Molecular Sciences, 2017, 18, 187.	1.8	75
23	Neuroprotective effects of dexmedetomidine against hyperoxia-induced injury in the developing rat brain. PLoS ONE, 2017, 12, e0171498.	1.1	64
24	Isoflurane but not Fentanyl Causes Apoptosis in Immature Primary Neuronal Cells. Open Anesthesiology Journal, 2017, 11, 39-47.	0.4	0
25	Erythropoietin Restores Long-Term Neurocognitive Function Involving Mechanisms of Neuronal Plasticity in a Model of Hyperoxia-Induced Preterm Brain Injury. Oxidative Medicine and Cellular Longevity, 2016, 2016, 1-13.	1.9	29
26	Effects of progesterone on hyperoxia-induced damage in mouse C8 $\alpha$ 1A astrocytes. Brain and Behavior, 2016, 6, e00435.	1.0	3
27	Oligodendroglial maldevelopment in the cerebellum after postnatal hyperoxia and its prevention by minocycline. Glia, 2015, 63, 1825-1839.	2.5	26
28	Neuroprotective Effect of Dexmedetomidine on Hyperoxia-Induced Toxicity in the Neonatal Rat Brain. Oxidative Medicine and Cellular Longevity, 2015, 2015, 1-10.	1.9	55
29	Neuroinflammation after traumatic injury to the developing brain. Scripta Scientifica Medica, 2015, 47, 47.	0.1	3
30	Inhibition of Acetylcholinesterase Modulates NMDA Receptor Antagonist Mediated Alterations in the Developing Brain. International Journal of Molecular Sciences, 2014, 15, 3784-3798.	1.8	4
31	Caffeine protects neuronal cells against injury caused by hyperoxia in the immature brain. Free Radical Biology and Medicine, 2014, 67, 221-234.	1.3	55
32	Minocycline protects the immature white matter against hyperoxia. Experimental Neurology, 2014, 254, 153-165.	2.0	45
33	Effects of moderate and deep hypothermia on RNA-binding proteins RBM3 and CIRP expressions in murine hippocampal brain slices. Brain Research, 2013, 1504, 74-84.	1.1	71
34	Oxygen Toxicity Is Reduced by Acetylcholinesterase Inhibition in the Developing Rat Brain. Developmental Neuroscience, 2013, 35, 255-264.	1.0	21
35	Prevention of hyperoxia-mediated pulmonary inflammation in neonatal rats by caffeine. European Respiratory Journal, 2013, 41, 966-973.	3.1	104
36	Erythropoietin Modulates Autophagy Signaling in the Developing Rat Brain in an In Vivo Model of Oxygen-Toxicity. International Journal of Molecular Sciences, 2012, 13, 12939-12951.	1.8	43

#	ARTICLE	IF	CITATIONS
37	Prevention of neonatal oxygen-induced brain damage by reduction of intrinsic apoptosis. <i>Cell Death and Disease</i> , 2012, 3, e250-e250.	2.7	38
38	Hyperoxia changes the balance of the thioredoxin/peroxiredoxin system in the neonatal rat brain. <i>Brain Research</i> , 2012, 1484, 68-75.	1.1	23
39	The Protective Role of Erythropoietin in the Developing Brain. , 2012, , .		0
40	Minocycline protects oligodendroglial precursor cells against injury caused by oxygen-glucose deprivation. <i>Journal of Neuroscience Research</i> , 2012, 90, 933-944.	1.3	29
41	Adolescent hyperactivity and impaired coordination after neonatal hyperoxia. <i>Experimental Neurology</i> , 2012, 235, 374-379.	2.0	31
42	Interaction of Inflammation and Hyperoxia in a Rat Model of Neonatal White Matter Damage. <i>PLoS ONE</i> , 2012, 7, e49023.	1.1	74
43	Hyperoxia-Induced Injury in Oligodendroglial Precursor Cells (OPC) is Attenuated by Minocycline. <i>Pediatric Research</i> , 2011, 70, 748-748.	1.1	0
44	Neurotoxic Effects of MDMA (Ecstasy) on the Developing Rodent Brain. <i>Developmental Neuroscience</i> , 2010, 32, 197-207.	1.0	8
45	Erythropoietin attenuates hyperoxia-induced oxidative stress in the developing rat brain. <i>Brain, Behavior, and Immunity</i> , 2010, 24, 792-799.	2.0	40
46	A critical role for Fas/CD95 dependent signaling pathways in the pathogenesis of hyperoxia-induced brain injury. <i>Annals of Neurology</i> , 2008, 64, 664-673.	2.8	39
47	IL-22 Induces Lipopolysaccharide-Binding Protein in Hepatocytes: A Potential Systemic Role of IL-22 in Crohn's Disease. <i>Journal of Immunology</i> , 2007, 178, 5973-5981.	0.4	254
48	IL-19 and IL-20: two novel cytokines with importance in inflammatory diseases. <i>Expert Opinion on Therapeutic Targets</i> , 2007, 11, 601-612.	1.5	89
49	Activation of caspase-1 dependent interleukins in developmental brain trauma. <i>Neurobiology of Disease</i> , 2007, 25, 614-622.	2.1	24
50	Acute and long-term proteome changes induced by oxidative stress in the developing brain. <i>Cell Death and Differentiation</i> , 2006, 13, 1097-1109.	5.0	53
51	Antisense oligonucleotides and short interfering RNAs silencing the cyclin-dependent kinase inhibitor p21 improve proliferation of Duchenne muscular dystrophy patients' primary skeletal myoblasts. <i>Journal of Molecular Medicine</i> , 2005, 83, 64-71.	1.7	8
52	NMDA antagonist inhibits the extracellular signal-regulated kinase pathway and suppresses cancer growth. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 15605-15610.	3.3	129
53	Transfection of normal primary human skeletal myoblasts with p21 and p57 antisense oligonucleotides to improve their proliferation: a first step towards an alternative molecular therapy approach of Duchenne muscular dystrophy. <i>Journal of Molecular Medicine</i> , 2003, 81, 355-362.	1.7	15
54	Elevated p21 mRNA level in skeletal muscle of DMD patients and mdx mice indicates either an exhausted satellite cell pool or a higher p21 expression in dystrophin-deficient cells per se. <i>Journal of Molecular Medicine</i> , 2000, 78, 569-574.	1.7	19