Sangram S Sisodia

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/4330024/sangram-s-sisodia-publications-by-year.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

67 13,794 41 70 g-index

70 14,819 11.8 5.82 ext. papers ext. citations avg, IF L-index

#	Paper	IF	Citations
67	Gut microbiota-driven brain Alamyloidosis in mice requires microglia. <i>Journal of Experimental Medicine</i> , 2022 , 219,	16.6	7
66	Infection and inflammation: New perspectives on Alzheimerß disease. <i>Brain, Behavior, & Immunity - Health</i> , 2022 , 22, 100462	5.1	2
65	An APP ectodomain mutation outside of the Aldomain promotes Alproduction in vitro and deposition in vivo. <i>Journal of Experimental Medicine</i> , 2021 , 218,	16.6	2
64	Alteration in synaptic nanoscale organization dictates amyloidogenic processing in Alzheimerß disease. <i>IScience</i> , 2021 , 24, 101924	6.1	6
63	Modulation of amyloid deposition and neuroinflammation by the microbiome. <i>Alzheimern</i> and <i>Dementia</i> , 2020 , 16, e044154	1.2	
62	Synergistic depletion of gut microbial consortia, but not individual antibiotics, reduces amyloidosis in APPPS1-21 Alzheimerß transgenic mice. <i>Scientific Reports</i> , 2020 , 10, 8183	4.9	25
61	Negative evidence for a role of APH1B T27I variant in Alzheimerß disease. <i>Human Molecular Genetics</i> , 2020 , 29, 955-966	5.6	3
60	Deficits in Enrichment-Dependent Neurogenesis and Enhanced Anxiety Behaviors Mediated by Expression of Alzheimer Disease-Linked Ps1 Variants Are Rescued by Microglial Depletion. <i>Journal of Neuroscience</i> , 2019 , 39, 6766-6780	6.6	15
59	Sex-specific effects of microbiome perturbations on cerebral Alamyloidosis and microglia phenotypes. <i>Journal of Experimental Medicine</i> , 2019 , 216, 1542-1560	16.6	93
58	EC-03-03: GUT MICROBIOME ALTERATIONS IN ALZHEIMER DISEASE: PRECLINICAL EVIDENCE 2018 , 14, P1007-P1007		
57	Microglia turnover with aging and in an Alzheimerß model via long-term in vivo single-cell imaging. <i>Nature Neuroscience</i> , 2017 , 20, 1371-1376	25.5	193
56	Antibiotic-induced perturbations in microbial diversity during post-natal development alters amyloid pathology in an aged APP/PS1 murine model of Alzheimerß disease. <i>Scientific Reports</i> , 2017 , 7, 10411	4.9	133
55	Antibiotic-induced perturbations in gut microbial diversity influences neuro-inflammation and amyloidosis in a murine model of Alzheimerß disease. <i>Scientific Reports</i> , 2016 , 6, 30028	4.9	314
54	Evidence That the "Lid" Domain of Nicastrin Is Not Essential for Regulating Esecretase Activity. Journal of Biological Chemistry, 2016 , 291, 6748-53	5.4	6
53	Acne inversa caused by missense mutations in NCSTN is not fully compatible with impairments in Notch signaling. <i>Journal of Investigative Dermatology</i> , 2015 , 135, 618-620	4.3	22
52	The topology of pen-2, a Becretase subunit, revisited: evidence for a reentrant loop and a single pass transmembrane domain. <i>Molecular Neurodegeneration</i> , 2015 , 10, 39	19	11
51	Differential release of Eamyloid from dendrite- versus axon-targeted APP. <i>Journal of Neuroscience</i> , 2014 , 34, 12313-27	6.6	29

(2002-2014)

50	Soluble Becretase modulators selectively inhibit the production of the 42-amino acid amyloid peptide variant and augment the production of multiple carboxy-truncated amyloid pecies. <i>Biochemistry</i> , 2014 , 53, 702-13	3.2	40
49	A synthetic antibody fragment targeting nicastrin affects assembly and trafficking of Elecretase. Journal of Biological Chemistry, 2014 , 289, 34851-61	5.4	5
48	Mutant presenilin 1 expression in excitatory neurons impairs enrichment-mediated phenotypes of adult hippocampal progenitor cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013 , 110, 9148-53	11.5	13
47	Endogenous expression of FAD-linked PS1 impairs proliferation, neuronal differentiation and survival of adult hippocampal progenitors. <i>Molecular Neurodegeneration</i> , 2013 , 8, 41	19	13
46	Trafficking and proteolytic processing of APP. Cold Spring Harbor Perspectives in Medicine, 2012, 2, a006	52 <u>57.</u> p	628
45	Identification of a tetratricopeptide repeat-like domain in the nicastrin subunit of Esecretase using synthetic antibodies. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012 , 109, 8534-9	11.5	29
44	Structure of gamma-secretase and its trimeric pre-activation intermediate by single-particle electron microscopy. <i>Journal of Biological Chemistry</i> , 2011 , 286, 21440-9	5.4	33
43	Amyloid beta from axons and dendrites reduces local spine number and plasticity. <i>Nature Neuroscience</i> , 2010 , 13, 190-6	25.5	241
42	Activation and intrinsic gamma-secretase activity of presenilin 1. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010 , 107, 21435-40	11.5	112
41	Modulation of gamma-secretase reduces beta-amyloid deposition in a transgenic mouse model of Alzheimer B disease. <i>Neuron</i> , 2010 , 67, 769-80	13.9	208
40	Non-cell-autonomous effects of presenilin 1 variants on enrichment-mediated hippocampal progenitor cell proliferation and differentiation. <i>Neuron</i> , 2008 , 59, 568-80	13.9	147
39	Expression of a familial Alzheimer disease-linked presentiin-1 variant enhances perforant pathway lesion-induced neuronal loss in the entorhinal cortex. <i>Journal of Neuroscience</i> , 2006 , 26, 429-34	6.6	24
38	A sequence within the first transmembrane domain of PEN-2 is critical for PEN-2-mediated endoproteolysis of presenilin 1. <i>Journal of Biological Chemistry</i> , 2005 , 280, 1992-2001	5.4	51
37	Evidence that the "NF" motif in transmembrane domain 4 of presenilin 1 is critical for binding with PEN-2. <i>Journal of Biological Chemistry</i> , 2005 , 280, 41953-66	5.4	67
36	Regulated hyperaccumulation of presenilin-1 and the "gamma-secretase" complex. Evidence for differential intramembranous processing of transmembrane subatrates. <i>Journal of Biological Chemistry</i> , 2003 , 278, 33992-4002	5.4	88
35	The Notch ligands, Delta1 and Jagged2, are substrates for presenilin-dependent "gamma-secretase" cleavage. <i>Journal of Biological Chemistry</i> , 2003 , 278, 7751-4	5.4	159
34	APP processing and synaptic function. <i>Neuron</i> , 2003 , 37, 925-37	13.9	1248
33	Evidence that synaptically released beta-amyloid accumulates as extracellular deposits in the hippocampus of transgenic mice. <i>Journal of Neuroscience</i> , 2002 , 22, 9785-93	6.6	255

32	gamma-Secretase, Notch, Abeta and Alzheimerß disease: where do the presenilins fit in?. <i>Nature Reviews Neuroscience</i> , 2002 , 3, 281-90	13.5	461
31	Characterization of a presenilin-mediated amyloid precursor protein carboxyl-terminal fragment gamma. Evidence for distinct mechanisms involved in gamma -secretase processing of the APP and Notch1 transmembrane domains. <i>Journal of Biological Chemistry</i> , 2001 , 276, 43756-60	5.4	171
30	Multiple effects of aspartate mutant presenilin 1 on the processing and trafficking of amyloid precursor protein. <i>Journal of Biological Chemistry</i> , 2001 , 276, 43343-50	5.4	78
29	Deficient neurogenesis in forebrain-specific presenilin-1 knockout mice is associated with reduced clearance of hippocampal memory traces. <i>Neuron</i> , 2001 , 32, 911-26	13.9	408
28	The value of transgenic models for the study of neurodegenerative diseases. <i>Annals of the New York Academy of Sciences</i> , 2000 , 920, 179-91	6.5	44
27	Requirement for presenilin 1 in facilitating lagged 2-mediated endoproteolysis and signaling of notch 1. <i>Journal of Molecular Neuroscience</i> , 2000 , 15, 189-204	3.3	44
26	Furin mediates enhanced production of fibrillogenic ABri peptides in familial British dementia. <i>Nature Neuroscience</i> , 1999 , 2, 984-8	25.5	128
25	Transgenic mouse models of Alzheimerß disease and amyotrophic lateral sclerosis. <i>Brain Pathology</i> , 1998 , 8, 735-57	6	20
24	Alzheimer R disease: genetic studies and transgenic models. <i>Annual Review of Genetics</i> , 1998 , 32, 461-93	314.5	349
23	Mutant genes in familial Alzheimerß disease and transgenic models. <i>Annual Review of Neuroscience</i> , 1998 , 21, 479-505	17	516
22	Post-translational processing and turnover kinetics of presynaptically targeted amyloid precursor superfamily proteins in the central nervous system. <i>Journal of Biological Chemistry</i> , 1998 , 273, 11100-6	5.4	67
21	Alzheimer amyloid protein precursor in the rat hippocampus: transport and processing through the perforant path. <i>Journal of Neuroscience</i> , 1998 , 18, 9629-37	6.6	234
20	Evidence that levels of presenilins (PS1 and PS2) are coordinately regulated by competition for limiting cellular factors. <i>Journal of Biological Chemistry</i> , 1997 , 272, 28415-22	5.4	275
19	Ectodomain phosphorylation of beta-amyloid precursor protein at two distinct cellular locations. Journal of Biological Chemistry, 1997 , 272, 1896-903	5.4	55
18	Altered metabolism of familial Alzheimer disease-linked amyloid precursor protein variants in yeast artificial chromosome transgenic mice. <i>Human Molecular Genetics</i> , 1997 , 6, 1535-41	5.6	90
17	Amyloid \$sZ-protein stimulates parallel increases in cellular levels of its precursor and amyloid precursor-like protein 2 (APLP2) in human cerebrovascular smooth muscle cells. <i>Amyloid: the International Journal of Experimental and Clinical Investigation: the Official Journal of the</i>	2.7	4
16	Accelerated amyloid deposition in the brains of transgenic mice coexpressing mutant presenilin 1 and amyloid precursor proteins. <i>Neuron</i> , 1997 , 19, 939-45	13.9	885
15	Hyperaccumulation of FAD-linked presenilin 1 variants in vivo. <i>Nature Medicine</i> , 1997 , 3, 756-60	50.5	131

LIST OF PUBLICATIONS

14	Presenilin 1 is required for Notch1 and DII1 expression in the paraxial mesoderm. <i>Nature</i> , 1997 , 387, 288-92	50.4	669
13	Familial Alzheimerß disease-linked presenilin 1 variants elevate Abeta1-42/1-40 ratio in vitro and in vivo. <i>Neuron</i> , 1996 , 17, 1005-13	13.9	1350
12	Endoproteolysis of presenilin 1 and accumulation of processed derivatives in vivo. <i>Neuron</i> , 1996 , 17, 181-90	13.9	999
11	Inherited neurodegenerative diseases and transgenic models. <i>Brain Pathology</i> , 1996 , 6, 467-80	6	8
10	A vector for expressing foreign genes in the brains and hearts of transgenic mice. <i>Genetic Analysis, Techniques and Applications</i> , 1996 , 13, 159-63		290
9	Metabolism of the "Swedish" amyloid precursor protein variant in neuro2a (N2a) cells. Evidence that cleavage at the "beta-secretase" site occurs in the golgi apparatus. <i>Journal of Biological Chemistry</i> , 1996 , 271, 9390-7	5.4	245
8	Motor neuron disease and model systems: aetiologies, mechanisms and therapies. <i>Novartis Foundation Symposium</i> , 1996 , 196, 3-13; discussion 13-7		1
7	A mouse model for Down syndrome exhibits learning and behaviour deficits. <i>Nature Genetics</i> , 1995 , 11, 177-84	36.3	739
6	Role of the beta-amyloid protein in Alzheimerß disease. FASEB Journal, 1995, 9, 366-70	0.9	237
5	Nucleotide sequence of the chromosome 14-encoded S182 cDNA and revised secondary structure prediction. <i>Amyloid: the International Journal of Experimental and Clinical Investigation: the Official Journal of the International Society of Amyloidosis</i> , 1995 , 2, 188-190	2.7	14
4	beta-Amyloid precursor protein-deficient mice show reactive gliosis and decreased locomotor activity. <i>Cell</i> , 1995 , 81, 525-31	56.2	573
3	Cellular and molecular biology of Alzheimerß disease and animal models. <i>Annual Review of Medicine</i> , 1994 , 45, 435-46	17.4	112
2	Isolation and characterization of APLP2 encoding a homologue of the Alzheimerß associated amyloid beta protein precursor. <i>Nature Genetics</i> , 1993 , 5, 95-100	36.3	340
1	Neuronal degeneration in human diseases and animal models. <i>Journal of Neurobiology</i> , 1992 , 23, 1277	-94	29