

Luc Teyton

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/4334521/luc-teyton-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

70
papers

9,395
citations

37
h-index

96
g-index

114
ext. papers

10,070
ext. citations

16.2
avg. IF

5.66
L-index

#	Paper	IF	Citations
70	Lipid presentation by the protein C receptor links coagulation with autoimmunity. <i>Science</i> , 2021 , 371,	33.3	23
69	Unravelling the structural complexity of glycolipids with cryogenic infrared spectroscopy. <i>Nature Communications</i> , 2021 , 12, 1201	17.4	16
68	Pathogenic lipid-binding antiphospholipid antibodies are associated with severity of COVID-19. <i>Journal of Thrombosis and Haemostasis</i> , 2021 , 19, 2335-2347	15.4	6
67	Chromogranin A Deficiency Confers Protection From Autoimmune Diabetes via Multiple Mechanisms. <i>Diabetes</i> , 2021 , 70, 2860-2870	0.9	1
66	Glycolipids as Antigens for Semi-Invariant Natural Killer T Cells 2021 , 470-484		1
65	Intestinal Delivery of Proinsulin and IL-10 via Combined With Low-Dose Anti-CD3 Restores Tolerance Outside the Window of Acute Type 1 Diabetes Diagnosis. <i>Frontiers in Immunology</i> , 2020 , 11, 1103	8.4	3
64	PLA2G1B is involved in CD4 anergy and CD4 lymphopenia in HIV-infected patients. <i>Journal of Clinical Investigation</i> , 2020 , 130, 2872-2887	15.9	12
63	Position 57 of I-A controls early anti-insulin responses in NOD mice, linking an MHC susceptibility allele to type 1 diabetes onset. <i>Science Immunology</i> , 2019 , 4,	28	18
62	Single-Cell Analysis of CD4 T Cells in Type 1 Diabetes: From Mouse to Man, How to Perform Mechanistic Studies. <i>Diabetes</i> , 2019 , 68, 1886-1891	0.9	3
61	High-affinity anti-glycan antibodies: challenges and strategies. <i>Current Opinion in Immunology</i> , 2019 , 59, 65-71	7.8	3
60	Role of a selecting ligand in shaping the murine TCR repertoire. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2018 , 115, 1889-1894	11.5	22
59	Role of lipid transfer proteins in loading CD1 antigen-presenting molecules. <i>Journal of Lipid Research</i> , 2018 , 59, 1367-1373	6.3	10
58	PLD3 and PLD4 are single-stranded acid exonucleases that regulate endosomal nucleic-acid sensing. <i>Nature Immunology</i> , 2018 , 19, 942-953	19.1	56
57	Gene Profiling and T Cell Receptor Sequencing from Antigen-Specific CD4 T Cells. <i>Methods in Molecular Biology</i> , 2018 , 1712, 217-238	1.4	4
56	Using single cell analysis for translational studies in immune mediated diseases: Opportunities and challenges. <i>Molecular Immunology</i> , 2018 , 103, 191-199	4.3	5
55	Rapid CLIP dissociation from MHC II promotes an unusual antigen presentation pathway in autoimmunity. <i>Journal of Experimental Medicine</i> , 2018 , 215, 2617-2635	16.6	13
54	Data Streaming for Metabolomics: Accelerating Data Processing and Analysis from Days to Minutes. <i>Analytical Chemistry</i> , 2017 , 89, 1254-1259	7.8	20

53	New Directions for Natural Killer T Cells in the Immunotherapy of Cancer. <i>Frontiers in Immunology</i> , 2017 , 8, 1480	8.4	7
52	T cells control the generation of nanomolar-affinity anti-glycan antibodies. <i>Journal of Clinical Investigation</i> , 2017 , 127, 1491-1504	15.9	47
51	The processing and presentation of lipids and glycolipids to the immune system. <i>Immunological Reviews</i> , 2016 , 272, 109-19	11.3	28
50	Restricting nonclassical MHC genes coevolve with TRAV genes used by innate-like T cells in mammals. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016 , 113, E2983-92	11.5	45
49	Endogenous ligands of natural killer T cells are alpha-linked glycosylceramides. <i>Molecular Immunology</i> , 2015 , 68, 94-7	4.3	34
48	The identification of the endogenous ligands of natural killer T cells reveals the presence of mammalian linked glycosylceramides. <i>Immunity</i> , 2014 , 41, 543-54	32.3	170
47	Efficacy of ABX196, a new NKT agonist, in prophylactic human vaccination. <i>Vaccine</i> , 2014 , 32, 6138-45	4.1	33
46	Transport and uptake of immunogenic lipids. <i>Molecular Immunology</i> , 2013 , 55, 179-81	4.3	9
45	Scavenger receptors target glycolipids for natural killer T cell activation. <i>Journal of Clinical Investigation</i> , 2012 , 122, 3943-54	15.9	40
44	The endothelial protein C receptor supports tissue factor ternary coagulation initiation complex signaling through protease-activated receptors. <i>Journal of Biological Chemistry</i> , 2011 , 286, 5756-67	5.4	70
43	Fatty acid amide hydrolase shapes NKT cell responses by influencing the serum transport of lipid antigen in mice. <i>Journal of Clinical Investigation</i> , 2010 , 120, 1873-84	15.9	22
42	The diabetogenic mouse MHC class II molecule I-Ag7 is endowed with a switch that modulates TCR affinity. <i>Journal of Clinical Investigation</i> , 2010 , 120, 1578-90	15.9	31
41	The Endothelial Protein C Receptor Supports TF-VIIa-Xa Ternary Complex Signaling through Protease-Activated Receptors.. <i>Blood</i> , 2010 , 116, 1144-1144	2.2	
40	New design of MHC class II tetramers to accommodate fundamental principles of antigen presentation. <i>Journal of Immunology</i> , 2009 , 183, 7949-57	5.3	46
39	Natural killer T cell recognition of lipid antigens. <i>F1000 Biology Reports</i> , 2009 , 1, 97		
38	The role of HLA-DQ8 beta57 polymorphism in the anti-gluten T-cell response in coeliac disease. <i>Nature</i> , 2008 , 456, 534-8	50.4	78
37	Crystal structures of mouse CD1d-iGb3 complex and its cognate Valpha14 T cell receptor suggest a model for dual recognition of foreign and self glycolipids. <i>Journal of Molecular Biology</i> , 2008 , 377, 1104-16	6.5	88
36	Corrugoside, a new immunostimulatory alpha-galactoglycosphingolipid from the marine sponge <i>Axinella corrugata</i> . <i>Bioorganic and Medicinal Chemistry</i> , 2008 , 16, 2077-85	3.4	24

35	Visualizing skin T cell antigen expression during wound repair. <i>FASEB Journal</i> , 2008 , 22, 1065-35	0.9	1
34	The X-ray Structure of IAd in Complex with a Self Peptide Offers New Insights Into the Basis of Autoimmune Gastritis (AIG). <i>FASEB Journal</i> , 2008 , 22, 667.4	0.9	
33	The Niemann-Pick type C2 protein loads isoglobotrihexosylceramide onto CD1d molecules and contributes to the thymic selection of NKT cells. <i>Journal of Experimental Medicine</i> , 2007 , 204, 841-52	16.6	81
32	A distal effect of microsomal triglyceride transfer protein deficiency on the lysosomal recycling of CD1d. <i>Journal of Experimental Medicine</i> , 2007 , 204, 921-8	16.6	43
31	The biology of NKT cells. <i>Annual Review of Immunology</i> , 2007 , 25, 297-336	34.7	1704
30	The saga of MHC-bound peptides: a renaissance for antigen presentation?. <i>Journal of Clinical Investigation</i> , 2007 , 117, 3164-6	15.9	4
29	A modified alpha-galactosyl ceramide for staining and stimulating natural killer T cells. <i>Journal of Immunological Methods</i> , 2006 , 312, 34-9	2.5	155
28	Mechanisms imposing the Vbeta bias of Valpha14 natural killer T cells and consequences for microbial glycolipid recognition. <i>Journal of Experimental Medicine</i> , 2006 , 203, 1197-207	16.6	85
27	Cutting edge: impaired glycosphingolipid trafficking and NKT cell development in mice lacking Niemann-Pick type C1 protein. <i>Journal of Immunology</i> , 2006 , 177, 26-30	5.3	66
26	Glycolipids for natural killer T cells. <i>Chemical Society Reviews</i> , 2006 , 35, 771-9	58.5	113
25	Structure and function of a potent agonist for the semi-invariant natural killer T cell receptor. <i>Nature Immunology</i> , 2005 , 6, 810-8	19.1	267
24	Exogenous and endogenous glycolipid antigens activate NKT cells during microbial infections. <i>Nature</i> , 2005 , 434, 525-9	50.4	930
23	Editing of CD1d-bound lipid antigens by endosomal lipid transfer proteins. <i>Science</i> , 2004 , 303, 523-7	33.3	282
22	Effects of lipid chain lengths in alpha-galactosylceramides on cytokine release by natural killer T cells. <i>Journal of the American Chemical Society</i> , 2004 , 126, 13602-3	16.4	177
21	Lysosomal glycosphingolipid recognition by NKT cells. <i>Science</i> , 2004 , 306, 1786-9	33.3	817
20	The paradox of immune molecular recognition of alpha-galactosylceramide: low affinity, low specificity for CD1d, high affinity for alpha beta TCRs. <i>Journal of Immunology</i> , 2003 , 170, 4673-82	5.3	84
19	Crystal structure of CD1a in complex with a sulfatide self antigen at a resolution of 2.15 A. <i>Nature Immunology</i> , 2003 , 4, 808-15	19.1	201
18	Detection of low-avidity CD4+ T cells using recombinant artificial APC: following the antiovalbumin immune response. <i>Journal of Immunology</i> , 2003 , 170, 123-31	5.3	50

17	Susceptible MHC alleles, not background genes, select an autoimmune T cell reactivity. <i>Journal of Clinical Investigation</i> , 2003 , 112, 902-14	15.9	79
16	Multiple defects in antigen presentation and T cell development by mice expressing cytoplasmic tail-truncated CD1d. <i>Nature Immunology</i> , 2002 , 3, 55-60	19.1	165
15	Thymocyte expression of cathepsin L is essential for NKT cell development. <i>Nature Immunology</i> , 2002 , 3, 1069-74	19.1	95
14	Synthesis and NKT cell stimulating properties of fluorophore- and biotin-appended 6"-amino-6"-deoxy-galactosylceramides. <i>Organic Letters</i> , 2002 , 4, 1267-70	6.2	95
13	Crystal structure of the murine NK cell-activating receptor NKG2D at 1.95 A. <i>Nature Immunology</i> , 2001 , 2, 248-54	19.1	75
12	Differential regulation of antiviral T-cell immunity results in stable CD8+ but declining CD4+ T-cell memory. <i>Nature Medicine</i> , 2001 , 7, 913-9	50.5	493
11	The mouse CD1d-restricted repertoire is dominated by a few autoreactive T cell receptor families. <i>Journal of Experimental Medicine</i> , 2001 , 193, 893-904	16.6	150
10	Function and dysfunction of T cell receptor: structural studies. <i>Immunologic Research</i> , 2000 , 21, 325-30	4.3	5
9	The I-Ag7 MHC class II molecule linked to murine diabetes is a promiscuous peptide binder. <i>Journal of Immunology</i> , 2000 , 165, 3214-25	5.3	111
8	A structural framework for deciphering the link between I-Ag7 and autoimmune diabetes. <i>Science</i> , 2000 , 288, 505-11	33.3	225
7	Effects of complementarity determining region mutations on the affinity of an alpha/beta T cell receptor: measuring the energy associated with CD4/CD8 repertoire skewing. <i>Journal of Experimental Medicine</i> , 1999 , 189, 461-70	16.6	28
6	Structural basis of T cell recognition. <i>Annual Review of Immunology</i> , 1999 , 17, 369-97	34.7	427
5	Probing the activation requirements for naive CD8+ T cells with Drosophila cell transfectants as antigen presenting cells. <i>Immunological Reviews</i> , 1998 , 165, 249-65	11.3	26
4	T-cell receptor peptide-MHC interactions: biological lessons from structural studies. <i>Current Opinion in Biotechnology</i> , 1998 , 9, 338-43	11.4	17
3	Structural basis of plasticity in T cell receptor recognition of a self peptide-MHC antigen. <i>Science</i> , 1998 , 279, 1166-72	33.3	595
2	Invariant chain distinguishes between the exogenous and endogenous antigen presentation pathways. <i>Nature</i> , 1990 , 348, 39-44	50.4	275
1	Intracellular transport of class II MHC molecules directed by invariant chain. <i>Nature</i> , 1990 , 348, 600-5	50.4	484