

# Pedro M Nieto

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4346872/publications.pdf>

Version: 2024-02-01

94  
papers

3,930  
citations

117571

34  
h-index

128225

60  
g-index

98  
all docs

98  
docs citations

98  
times ranked

3833  
citing authors

#	ARTICLE	IF	CITATIONS
1	GAG Multivalent Systems to Interact with Langerin. <i>Current Medicinal Chemistry</i> , 2022, 29, 1173-1192.	1.2	6
2	Fluorous-Tag-Assisted Synthesis of GAG-Like Oligosaccharides. <i>Methods in Molecular Biology</i> , 2022, 2303, 37-47.	0.4	3
3	The Interaction between Chondroitin Sulfate and Dermatan Sulfate Tetrasaccharides and Pleiotrophin. <i>International Journal of Molecular Sciences</i> , 2022, 23, 3026.	1.8	3
4	Pleiotrophin Interaction with Synthetic Glycosaminoglycan Mimetics. <i>Pharmaceuticals</i> , 2022, 15, 496.	1.7	0
5	Synthesis, structure and midkine binding of chondroitin sulfate oligosaccharide analogues. <i>Organic and Biomolecular Chemistry</i> , 2021, 19, 5312-5326.	1.5	3
6	Synthesis, self-assembly and Langerin recognition studies of a resorcinarene-based glycocluster exposing a hyaluronic acid thiodisaccharide mimetic. <i>Organic and Biomolecular Chemistry</i> , 2021, 19, 6455-6467.	1.5	0
7	Midkine Interaction with Chondroitin Sulfate Model Synthetic Tetrasaccharides and Their Mimetics: The Role of Aromatic Interactions. <i>Chemistry - A European Journal</i> , 2021, 27, 12395-12409.	1.7	7
8	Synthesis of (1 $\alpha$ '3) Thiodisaccharides of GlcNAc and the Serendipitous Formation of 2,3-Dideoxy-(1 $\alpha$ '2)-thiodisaccharides through a Vinyl Azide Intermediate. <i>Journal of Organic Chemistry</i> , 2020, 85, 306-317.	1.7	13
9	Second-Generation Dendrimers with Chondroitin Sulfate Type-E Disaccharides as Multivalent Ligands for Langerin. <i>Biomacromolecules</i> , 2020, 21, 2726-2734.	2.6	6
10	Langerin-Heparin Interaction: Analysis of the Binding to the Non-Lectin Site. <i>Natural Product Communications</i> , 2019, 14, 1934578X1985159.	0.2	3
11	Synthesis of a Fluorous-Tagged Hexasaccharide and Interaction with Growth Factors Using Sugar-Coated Microplates. <i>Molecules</i> , 2019, 24, 1591.	1.7	4
12	Unexpected loss of stereoselectivity in glycosylation reactions during the synthesis of chondroitin sulfate oligosaccharides. <i>Beilstein Journal of Organic Chemistry</i> , 2019, 15, 137-144.	1.3	3
13	Fluorous-tag assisted synthesis of a glycosaminoglycan mimetic tetrasaccharide as a high-affinity FGF-2 and midkine ligand. <i>Bioorganic and Medicinal Chemistry</i> , 2018, 26, 1076-1085.	1.4	12
14	Rational-Differential Design of Highly Specific Glycomimetic Ligands: Targeting DC-SIGN and Excluding Langerin Recognition. <i>ACS Chemical Biology</i> , 2018, 13, 600-608.	1.6	56
15	The Use of NMR to Study Transient Carbohydrate-Protein Interactions. <i>Frontiers in Molecular Biosciences</i> , 2018, 5, 33.	1.6	12
16	Glycodendrimers as Chondroitin Sulfate Mimetics: Synthesis and Binding to Growth Factor Midkine. <i>Chemistry - A European Journal</i> , 2017, 23, 11338-11345.	1.7	26
17	Interactions between a Heparin Trisaccharide Library and FGF-1 Analyzed by NMR Methods. <i>International Journal of Molecular Sciences</i> , 2017, 18, 1293.	1.8	13
18	Flavonoid glycosides from <i>Persea caerulea</i> . Unraveling their interactions with SDS-micelles through matrix-assisted DOSY, PGSE, mass spectrometry, and NOESY. <i>Magnetic Resonance in Chemistry</i> , 2016, 54, 718-728.	1.1	4

#	ARTICLE	IF	CITATIONS
19	Chondroitin Sulfate Tetrasaccharides: Synthesis, Three-Dimensional Structure and Interaction with Midkine. <i>Chemistry - A European Journal</i> , 2016, 22, 2356-2369.	1.7	45
20	Improvement on binding of chondroitin sulfate derivatives to midkine by increasing hydrophobicity. <i>Organic and Biomolecular Chemistry</i> , 2016, 14, 3506-3509.	1.5	12
21	Detection and quantitative analysis of two independent binding modes of a small ligand responsible for DC-SIGN clustering. <i>Organic and Biomolecular Chemistry</i> , 2016, 14, 335-344.	1.5	18
22	Mimicking Tyrosine Phosphorylation in Human Cytochrome <i>c</i> by the Evolved tRNA Synthetase Technique. <i>Chemistry - A European Journal</i> , 2015, 21, 15004-15012.	1.7	32
23	Langerin-Heparin Interaction: Two Binding Sites for Small and Large Ligands As Revealed by a Combination of NMR Spectroscopy and Cross-Linking Mapping Experiments. <i>Journal of the American Chemical Society</i> , 2015, 137, 4100-4110.	6.6	61
24	Structures of Glycans Bound to Receptors from Saturation Transfer Difference (STD) NMR Spectroscopy: Quantitative Analysis by Using CORCEMA-ST. <i>Methods in Molecular Biology</i> , 2015, 1273, 475-487.	0.4	5
25	A STD-NMR Study of the Interaction of the Anabaena Ferredoxin-NADP+ Reductase with the Coenzyme. <i>Molecules</i> , 2014, 19, 672-685.	1.7	1
26	Importance of the polarity of the glycosaminoglycan chain on the interaction with FGF-1. <i>Glycobiology</i> , 2014, 24, 1004-1009.	1.3	24
27	Synthesis of Chondroitin Sulfate Oligosaccharides Using <i>N</i> -(Tetrachlorophthaloyl)- and <i>N</i> -(Trifluoroacetyl)galactosamine Building Blocks. <i>European Journal of Organic Chemistry</i> , 2014, 2014, 3868-3884.	1.2	27
28	Synthesis of hyaluronic acid oligosaccharides and exploration of a fluorine-assisted approach. <i>Carbohydrate Research</i> , 2014, 394, 17-25.	1.1	18
29	Heparin Modulates the Mitogenic Activity of Fibroblast Growth Factor by Inducing Dimerization of its Receptor. A 3D View by Using NMR. <i>ChemBioChem</i> , 2013, 14, 1732-1744.	1.3	40
30	Structure of a Glycomimetic Ligand in the Carbohydrate Recognition Domain of C-type Lectin DC-SIGN. Structural Requirements for Selectivity and Ligand Design. <i>Journal of the American Chemical Society</i> , 2013, 135, 2518-2529.	6.6	75
31	Synthesis of chondroitin/dermatan sulfate-like oligosaccharides and evaluation of their protein affinity by fluorescence polarization. <i>Organic and Biomolecular Chemistry</i> , 2013, 11, 3510.	1.5	36
32	Insights into the Glycosaminoglycan-Mediated Cytotoxic Mechanism of Eosinophil Cationic Protein Revealed by NMR. <i>ACS Chemical Biology</i> , 2013, 8, 144-151.	1.6	27
33	NMR studies on carbohydrate interactions with DC-SIGN towards a quantitative STD analysis. <i>Pure and Applied Chemistry</i> , 2013, 85, 1771-1787.	0.9	4
34	Conformations of the iduronate ring in short heparin fragments described by time-averaged distance restrained molecular dynamics. <i>Glycobiology</i> , 2013, 23, 1220-1229.	1.3	27
35	3D structure of a heparin mimetic analogue of a FGF-1 activator. A NMR and molecular modelling study. <i>Organic and Biomolecular Chemistry</i> , 2013, 11, 8269.	1.5	22
36	Synthesis and Characterization of Linker-Armed Fucose-Based Glycomimetics. <i>European Journal of Organic Chemistry</i> , 2013, 2013, 5303-5314.	1.2	18

#	ARTICLE	IF	CITATIONS
37	Synthesis of amine-functionalized heparin oligosaccharides for the investigation of carbohydrate-protein interactions in microtiter plates. <i>Organic and Biomolecular Chemistry</i> , 2012, 10, 2146.	1.5	28
38	Effect of the Substituents of the Neighboring Ring in the Conformational Equilibrium of Iduronate in Heparin-like Trisaccharides. <i>Chemistry - A European Journal</i> , 2012, 18, 16319-16331.	1.7	32
39	A Non-damaging Method to Analyze the Configuration and Dynamics of Nitrotyrosines in Proteins. <i>Chemistry - A European Journal</i> , 2012, 18, 3872-3878.	1.7	9
40	sp <sup>2</sup> -Aminosugar O-, S-, and N-Glycosides as Conformational Mimics of $\beta$ -Linked Disaccharides; Implications for Glycosidase Inhibition. <i>Chemistry - A European Journal</i> , 2012, 18, 8527-8539.	1.7	51
41	Insights into molecular recognition of LewisX mimics by DC-SIGN using NMR and molecular modelling. <i>Organic and Biomolecular Chemistry</i> , 2011, 9, 7705.	1.5	21
42	STD-NMR: application to transient interactions between biomolecules—a quantitative approach. <i>European Biophysics Journal</i> , 2011, 40, 1357-1369.	1.2	140
43	Conformational Selection of the AGA <sup>1A</sup> Heparin Pentasaccharide when Bound to the Fibroblast Growth Factor Receptor. <i>Chemistry - A European Journal</i> , 2011, 17, 11204-11209.	1.7	32
44	Microwave-assisted sulfonation of heparin oligosaccharides. <i>Tetrahedron Letters</i> , 2011, 52, 441-443.	0.7	21
45	Nitration of tyrosine 74 prevents human cytochrome c to play a key role in apoptosis signaling by blocking caspase-9 activation. <i>Biochimica Et Biophysica Acta - Bioenergetics</i> , 2010, 1797, 981-993.	0.5	72
46	Structural and functional changes induced by tyrosine nitration in cytochrome c, a bi-functional protein. <i>Biochimica Et Biophysica Acta - Bioenergetics</i> , 2010, 1797, 70.	0.5	0
47	Polymer-Supported Synthesis of Oligosaccharides Using a Diisopropylsiloxane Linker and Trichloroacetimidate Donors. <i>European Journal of Organic Chemistry</i> , 2010, 2010, 2138-2147.	1.2	12
48	Ligand-Receptor Binding Affinities from Saturation Transfer Difference (STD) NMR Spectroscopy: The Binding Isotherm of STD Initial Growth Rates. <i>Chemistry - A European Journal</i> , 2010, 16, 7803-7812.	1.7	161
49	Exploration of the use of an acylsulfonamide safety-catch linker for the polymer-supported synthesis of hyaluronic acid oligosaccharides. <i>Carbohydrate Research</i> , 2010, 345, 565-571.	1.1	12
50	Mutants of the Arabidopsis thaliana Cation/H <sup>+</sup> Antiporter AtNHX1 Conferring Increased Salt Tolerance in Yeast. <i>Journal of Biological Chemistry</i> , 2009, 284, 14276-14285.	1.6	71
51	Synthesis of Novel DC-SIGN Ligands with an $\beta$ -Fucosylamide Anchor. <i>ChemBioChem</i> , 2008, 9, 1921-1930.	1.3	58
52	Saturation Transfer Difference (STD) NMR Spectroscopy Characterization of Dual Binding Mode of a Mannose Disaccharide to DC-SIGN. <i>ChemBioChem</i> , 2008, 9, 2225-2227.	1.3	63
53	Experimental Measurement of Carbohydrate-Aromatic Stacking in Water by Using a Dangling-Ended DNA Model System. <i>Chemistry - A European Journal</i> , 2008, 14, 7828-7835.	1.7	33
54	Docking, synthesis, and NMR studies of mannosyl trisaccharide ligands for DC-SIGN lectin. <i>Organic and Biomolecular Chemistry</i> , 2008, 6, 2743.	1.5	37

#	ARTICLE	IF	CITATIONS
55	NMR Structural Studies of Oligosaccharides Related to Cancer Processes. <i>Anti-Cancer Agents in Medicinal Chemistry</i> , 2008, 8, 52-63.	0.9	13
56	1,2-Mannobioside Mimic: Synthesis, DC-SIGN Interaction by NMR and Docking, and Antiviral Activity. <i>ChemMedChem</i> , 2007, 2, 1030-1036.	1.6	73
57	Solution NMR structure of a human FGF-1 monomer, activated by a hexasaccharide heparin-analogue. <i>FEBS Journal</i> , 2006, 273, 4716-4727.	2.2	57
58	Backbone dynamics of a biologically active human FGF-1 monomer, complexed to a hexasaccharide heparin-analogue, by <sup>15</sup> N NMR relaxation methods. <i>Journal of Biomolecular NMR</i> , 2006, 35, 225-239.	1.6	20
59	Structure and dynamics of the conserved protein GPI anchor core inserted into detergent micelles. <i>Glycobiology</i> , 2006, 16, 969-980.	1.3	21
60	Conformational Study of GPI Anchors: the Common Oligosaccharide GPI Anchor Backbone. <i>European Journal of Organic Chemistry</i> , 2005, 2005, 3489-3498.	1.2	10
61	GENDER DIMORPHISM AND ALTITUDINAL VARIATION OF SECONDARY COMPOUNDS IN LEAVES OF THE GYNODIOECIOUS SHRUB <i>Daphne laureola</i> . <i>Journal of Chemical Ecology</i> , 2005, 31, 139-150.	0.9	19
62	NMR Analysis of the Transient Complex between Membrane Photosystem I and Soluble Cytochrome c6. <i>Journal of Biological Chemistry</i> , 2005, 280, 7925-7931.	1.6	37
63	Dynamic properties of biologically active synthetic heparin-like hexasaccharides. <i>Glycobiology</i> , 2005, 15, 1008-1015.	1.3	33
64	Conformational Flexibility of a Synthetic Glycosylaminoglycan Bound to a Fibroblast Growth Factor. FGF-1 Recognizes Both the 1C4 and 2SO Conformations of a Bioactive Heparin-like Hexasaccharide. <i>Journal of the American Chemical Society</i> , 2005, 127, 5778-5779.	6.6	69
65	Tachykinins and Tachykinin Receptors: Structure and Activity Relationships. <i>Current Medicinal Chemistry</i> , 2004, 11, 2045-2081.	1.2	274
66	The Activation of Fibroblast Growth Factors (FGFs) by Glycosaminoglycans: Influence of the Sulfation Pattern on the Biological Activity of FGF-1. <i>ChemBioChem</i> , 2004, 5, 55-61.	1.3	59
67	The heparin-Ca <sup>2+</sup> interaction: the influence of the O-sulfation pattern on binding. <i>Carbohydrate Research</i> , 2004, 339, 975-983.	1.1	36
68	Synthesis and structural study of two new heparin-like hexasaccharides. <i>Organic and Biomolecular Chemistry</i> , 2003, 1, 2253-2266.	1.5	40
69	Glycodendritic Structures Based on Boltorn Hyperbranched Polymers and Their Interactions with <i>Lens culinaris</i> Lectin. <i>Bioconjugate Chemistry</i> , 2003, 14, 817-823.	1.8	82
70	A molecular dynamics description of the conformational flexibility of the iduronate ring in glycosaminoglycans. <i>Chemical Communications</i> , 2003, , 1512-1513.	2.2	26
71	The Interactions of Cyanobacterial Cytochrome c6 and Cytochrome f, Characterized by NMR. <i>Journal of Biological Chemistry</i> , 2002, 277, 48685-48689.	1.6	33
72	The activation of fibroblast growth factors by heparin: Synthesis and structural study of rationally modified heparin-like oligosaccharides. <i>Canadian Journal of Chemistry</i> , 2002, 80, 917-936.	0.6	37

#	ARTICLE	IF	CITATIONS
73	Synthesis and Structure of 1-D-6-O-(2-Amino-2-deoxy- $\beta$ - and $\beta$ -D-gluco- and Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 747 Td (-gal 889-898.	1.2	17
74	The Heparin $\beta$ Ca <sup>2+</sup> Interaction: Structure of the Ca <sup>2+</sup> Binding Site. European Journal of Organic Chemistry, 2002, 2002, 2367.	1.2	26
75	The Activation of Fibroblast Growth Factors by Heparin: Synthesis, Structure, and Biological Activity of Heparin-Like Oligosaccharides. ChemBioChem, 2001, 2, 673-685.	1.3	89
76	Changes in the Expression of Tachykinin Receptors in the Rat Uterus During the Course of Pregnancy <sup>1</sup> . Biology of Reproduction, 2001, 65, 538-543.	1.2	36
77	Inositolphosphoglycan Mediators Structurally Related to Glycosyl Phosphatidylinositol Anchors: Synthesis, Structure and Biological Activity. Chemistry - A European Journal, 2000, 6, 3608-3621.	1.7	72
78	The solution conformation of glycosyl inositols related to inositolphosphoglycan (IPG) mediators. Tetrahedron: Asymmetry, 2000, 11, 37-51.	1.8	12
79	Structural basis for chitin recognition by defense proteins: GlcNAc residues are bound in a multivalent fashion by extended binding sites in hevein domains. Chemistry and Biology, 2000, 7, 529-543.	6.2	131
80	Interaction of heparin with Ca <sup>2+</sup> : A model study with a synthetic heparin-like hexasaccharide. Israel Journal of Chemistry, 2000, 40, 289-299.	1.0	17
81	$\beta$ -Alkyl- $\beta$ -Amino- $\beta$ -Lactam Peptides: Design, Synthesis, and Conformational Features. Angewandte Chemie - International Edition, 1999, 38, 3056-3058.	7.2	29
82	<sup>1</sup> H/ <sup>15</sup> N HSQC NMR Studies of Ligand Carboxylate Group Interactions with Arginine Residues in Complexes of Brodimoprim Analogues and Lactobacillus casei Dihydrofolate Reductase $\beta$ , $\beta$ . Biochemistry, 1999, 38, 2127-2134.	1.2	26
83	Interplay of Steric Hindrance and Hydrogen Bonding To Restrict Mono-O-substituted p-tert-Butylcalix[6]arenes in Cone Conformation. Journal of Organic Chemistry, 1998, 63, 1079-1085.	1.7	15
84	Correlated bond rotations in interactions of arginine residues with ligand carboxylate groups in protein ligand complexes. FEBS Letters, 1997, 405, 16-20.	1.3	41
85	NMR Detection of Arginine-Ligand Interactions in Complexes of Lactobacillus casei Dihydrofolate Reductase. FEBS Journal, 1996, 238, 435-439.	0.2	21
86	Synthesis of calix[6]arenes partially functionalized at the upper rim. Tetrahedron, 1995, 51, 12699-12720.	1.0	38
87	Calix[4]arene Sulfonates: Palladium-Catalyzed Intermolecular Migration of Sulfonyl Groups and Isolation of a Calix[4]arene in a Chiral 1,2-Alternate Conformation. Journal of Organic Chemistry, 1995, 60, 7419-7423.	1.7	29
88	Control of Calix[6]arene Conformations by Self-Inclusion of 1,3,5-Tri-O-alkyl Substituents: Synthesis and NMR Studies. Journal of the American Chemical Society, 1994, 116, 5814-5822.	6.6	110
89	Dipole moments can be used to determine the conformation of calix[4]arenes. Recueil Des Travaux Chimiques Des Pays-Bas, 1993, 112, 367-369.	0.0	16
90	Procedures for the Selective Alkylation of Calix[6]arenes at the Lower Rim. Synthesis, 1993, 1993, 380-386.	1.2	79

#	ARTICLE	IF	CITATIONS
91	Solution structure and conformational equilibria of a symmetrical calix[6]arene. Complete sequential and cyclostereospecific assignment of the low-temperature NMR spectra of a cycloasymmetric molecule. <i>Journal of Organic Chemistry</i> , 1992, 57, 6924-6931.	1.7	55
92	Carbon-13 NMR chemical shifts. A single rule to determine the conformation of calix[4]arenes. <i>Journal of Organic Chemistry</i> , 1991, 56, 3372-3376.	1.7	563
93	A stepwise synthesis of functionalized calix [4]arenes and a calix[6]arene with alternate electron-withdrawing substituents. <i>Tetrahedron</i> , 1990, 46, 671-682.	1.0	42
94	NMR of Sulfated Oligo- and Polysaccharides. , 0, , 189-229.		7