

Graham R Leggatt

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/4348609/graham-r-leggatt-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

86

papers

2,993

citations

30

h-index

52

g-index

92

ext. papers

3,414

ext. citations

5.6

avg, IF

5.1

L-index

#	Paper	IF	Citations
86	Evolution of Cancer Vaccines-Challenges, Achievements, and Future Directions. <i>Vaccines</i> , 2021 , 9,	5.3	9
85	PD-1 and beyond to Activate T Cells in Cutaneous Squamous Cell Cancers: The Case for 4-1BB and VISTA Antibodies in Combination Therapy. <i>Cancers</i> , 2021 , 13,	6.6	1
84	Regulatory T Cells but Not IL-10 Impair Cell-Mediated Immunity in Human Papillomavirus E7+ Hyperplastic Epithelium. <i>Journal of Investigative Dermatology</i> , 2021 , 141, 1264-1273.e3	4.3	2
83	Increased lipid metabolism impairs NK cell function and mediates adaptation to the lymphoma environment. <i>Blood</i> , 2020 , 136, 3004-3017	2.2	21
82	Serum antibodies against <i>Toxoplasma gondii</i> and <i>Neospora caninum</i> in southeast Queensland dugongs. <i>Marine Mammal Science</i> , 2020 , 36, 180-194	1.9	4
81	NKT Cell-Driven Enhancement of Antitumor Immunity Induced by Clec9a-Targeted Tailorable Nanoemulsion. <i>Cancer Immunology Research</i> , 2019 , 7, 952-962	12.5	6
80	HPV16 E7-Driven Epithelial Hyperplasia Promotes Impaired Antigen Presentation and Regulatory T-Cell Development. <i>Journal of Investigative Dermatology</i> , 2019 , 139, 2467-2476.e3	4.3	8
79	Cytokine/chemokine profiles in squamous cell carcinoma correlate with precancerous and cancerous disease stage. <i>Scientific Reports</i> , 2019 , 9, 17754	4.9	7
78	An Ex Vivo Human Tumor Assay Shows Distinct Patterns of EGFR Trafficking in Squamous Cell Carcinoma Correlating to Therapeutic Outcomes. <i>Journal of Investigative Dermatology</i> , 2019 , 139, 213-223	4.3	14
77	HPV16E7-Induced Hyperplasia Promotes CXCL9/10 Expression and Induces CXCR3 T-Cell Migration to Skin. <i>Journal of Investigative Dermatology</i> , 2018 , 138, 1348-1359	4.3	14
76	B cell lymphoma progression promotes the accumulation of circulating Ly6Clo monocytes with immunosuppressive activity. <i>Oncolmmunology</i> , 2018 , 7, e1393599	7.2	10
75	Development of a polyclonal anti-dugong immunoglobulin G (IgG) antibody with evaluation of total plasma IgG in a living dugong (<i>Dugong dugon</i>) population. <i>Veterinary Immunology and Immunopathology</i> , 2018 , 200, 16-25	2	1
74	Therapeutic vaccination with 4-1BB co-stimulation eradicates mouse acute myeloid leukemia. <i>Oncolmmunology</i> , 2018 , 7, e1486952	7.2	4
73	Recruitment of Antigen Presenting Cells to Skin Draining Lymph Node From HPV16E7-Expressing Skin Requires E7-Rb Interaction. <i>Frontiers in Immunology</i> , 2018 , 9, 2896	8.4	6
72	The Role of CXCR3 and Its Chemokine Ligands in Skin Disease and Cancer. <i>Frontiers in Medicine</i> , 2018 , 5, 271	4.9	54
71	Modulation of antigen presenting cell functions during chronic HPV infection. <i>Papillomavirus Research (Amsterdam, Netherlands)</i> , 2017 , 4, 58-65	4.6	28
70	Elevated frequencies of CD8 T cells expressing PD-1, CTLA-4 and Tim-3 within tumour from perineural squamous cell carcinoma patients. <i>PLoS ONE</i> , 2017 , 12, e0175755	3.7	24

69	DNA Vaccine Encoding HPV16 Oncogenes E6 and E7 Induces Potent Cell-mediated and Humoral Immunity Which Protects in Tumor Challenge and Drives E7-expressing Skin Graft Rejection. <i>Journal of Immunotherapy</i> , 2017 , 40, 62-70	5	33
68	HPV16-E7-Specific Activated CD8 T Cells in E7 Transgenic Skin and Skin Grafts. <i>Frontiers in Immunology</i> , 2017 , 8, 524	8.4	7
67	Interferon- γ -Derived from cytotoxic lymphocytes directly enhances their motility and cytotoxicity. <i>Cell Death and Disease</i> , 2017 , 8, e2836	9.8	159
66	Control of B-cell lymphoma by therapeutic vaccination and acquisition of immune resistance is independent of direct tumour IFN-gamma signalling. <i>Immunology and Cell Biology</i> , 2016 , 94, 554-62	5	5
65	Galectin-1 is associated with poor prognosis in patients with cutaneous head and neck cancer with perineural spread. <i>Cancer Immunology, Immunotherapy</i> , 2016 , 65, 213-22	7.4	7
64	Recent progress in vaccination against human papillomavirus-mediated cervical cancer. <i>Reviews in Medical Virology</i> , 2015 , 25 Suppl 1, 54-71	11.7	29
63	NKT cell-targeted vaccination plus anti-4-1BB antibody generates persistent CD8 T cell immunity against B cell lymphoma. <i>Oncot Immunology</i> , 2015 , 4, e990793	7.2	27
62	Aurora A Is Critical for Survival in HPV-Transformed Cervical Cancer. <i>Molecular Cancer Therapeutics</i> , 2015 , 14, 2753-61	6.1	23
61	Langerhans cell homeostasis and activation is altered in hyperplastic human papillomavirus type 16 E7 expressing epidermis. <i>PLoS ONE</i> , 2015 , 10, e0127155	3.7	19
60	The kinematics of cytotoxic lymphocytes influence their ability to kill target cells. <i>PLoS ONE</i> , 2014 , 9, e95248	3.7	10
59	Peptide Dose and/or Structure in Vaccines as a Determinant of T Cell Responses. <i>Vaccines</i> , 2014 , 2, 537-48	4.8	9
58	HPV16-E7 expression in squamous epithelium creates a local immune suppressive environment via CCL2- and CCL5- mediated recruitment of mast cells. <i>PLoS Pathogens</i> , 2014 , 10, e1004466	7.6	45
57	Immunosuppressive roles of natural killer T (NKT) cells in the skin. <i>Journal of Leukocyte Biology</i> , 2014 , 96, 49-54	6.5	11
56	Indoleamine 2,3-dioxygenase activity contributes to local immune suppression in the skin expressing human papillomavirus oncoprotein e7. <i>Journal of Investigative Dermatology</i> , 2013 , 133, 2686-2694	4.2	44
55	Expression of a single, viral oncoprotein in skin epithelium is sufficient to recruit lymphocytes. <i>PLoS ONE</i> , 2013 , 8, e57798	3.7	24
54	Γ cells augment rejection of skin grafts by enhancing cross-priming of CD8 T cells to skin-derived antigen. <i>Journal of Investigative Dermatology</i> , 2012 , 132, 1656-64	4.3	17
53	Histone deacetylase inhibitors in the generation of the anti-tumour immune response. <i>Immunology and Cell Biology</i> , 2012 , 90, 33-8	5	22
52	Impact of sex steroid ablation on viral, tumour and vaccine responses in aged mice. <i>PLoS ONE</i> , 2012 , 7, e42677	3.7	19

51	Role of intratumoural heterogeneity in cancer drug resistance: molecular and clinical perspectives. <i>EMBO Molecular Medicine</i> , 2012 , 4, 675-84	12	164
50	A combination of local inflammation and central memory T cells potentiates immunotherapy in the skin. <i>Journal of Immunology</i> , 2012 , 189, 5622-31	5.3	14
49	Human papillomavirus 16 E7 protein inhibits interferon- γ -mediated enhancement of keratinocyte antigen processing and T-cell lysis. <i>FEBS Journal</i> , 2011 , 278, 955-63	5.7	18
48	Regulation of immune responses to HPV infection and during HPV-directed immunotherapy. <i>Immunological Reviews</i> , 2011 , 239, 85-98	11.3	53
47	Prevention and treatment of papillomavirus-related cancers through immunization. <i>Annual Review of Immunology</i> , 2011 , 29, 111-38	34.7	82
46	NKT cells inhibit antigen-specific effector CD8 T cell induction to skin viral proteins. <i>Journal of Immunology</i> , 2011 , 187, 1601-8	5.3	30
45	Enhanced tumor growth in the NaS1 sulfate transporter null mouse. <i>Cancer Science</i> , 2010 , 101, 369-73	6.9	11
44	Secretion of IFN- γ but not IL-17 by CD1d-restricted NKT cells enhances rejection of skin grafts expressing epithelial cell-derived antigen. <i>Journal of Immunology</i> , 2010 , 184, 5663-9	5.3	29
43	Invariant NKT cells in hyperplastic skin induce a local immune suppressive environment by IFN- γ production. <i>Journal of Immunology</i> , 2010 , 184, 1242-50	5.3	52
42	Antigen-specific CD4 cells assist CD8 T-effector cells in eliminating keratinocytes. <i>Journal of Investigative Dermatology</i> , 2010 , 130, 1581-9	4.3	13
41	IL-1 signalling determines the fate of skin grafts expressing non-self protein in keratinocytes. <i>Experimental Dermatology</i> , 2010 , 19, 723-9	4	6
40	Both treated and untreated tumors are eliminated by short hairpin RNA-based induction of target-specific immune responses. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009 , 106, 8314-9	11.5	26
39	IFN- γ promotes generation of IL-10 secreting CD4+ T cells that suppress generation of CD8 responses in an antigen-experienced host. <i>Journal of Immunology</i> , 2009 , 183, 51-8	5.3	38
38	Epithelial expression of human papillomavirus type 16 E7 protein results in peripheral CD8 T-cell suppression mediated by CD4+CD25+ T cells. <i>European Journal of Immunology</i> , 2009 , 39, 481-90	6.1	32
37	Keratinocytes efficiently process endogenous antigens for cytotoxic T-cell mediated lysis. <i>Experimental Dermatology</i> , 2009 , 18, 1053-9	4	6
36	TLR7 stimulation augments T effector-mediated rejection of skin expressing neo-self antigen in keratinocytes. <i>European Journal of Immunology</i> , 2008 , 38, 73-81	6.1	16
35	Secondary immunisation with high-dose heterologous peptide leads to CD8 T cell populations with reduced functional avidity. <i>European Journal of Immunology</i> , 2007 , 37, 406-15	6.1	23
34	HPV vaccines: the beginning of the end for cervical cancer. <i>Current Opinion in Immunology</i> , 2007 , 19, 232-88	34	

33	Overcoming original antigenic sin to generate new CD8 T cell IFN-gamma responses in an antigen-experienced host. <i>Journal of Immunology</i> , 2006 , 177, 2873-9	5.3	27
32	Inhibition of cervical cancer cell growth in vitro and in vivo with lentiviral-vector delivered short hairpin RNA targeting human papillomavirus E6 and E7 oncogenes. <i>Cancer Gene Therapy</i> , 2006 , 13, 1023-32	5.4	105
31	RNA interference against human papillomavirus oncogenes in cervical cancer cells results in increased sensitivity to cisplatin. <i>Molecular Pharmacology</i> , 2005 , 68, 1311-9	4.3	94
30	Impaired antigen presentation and effectiveness of combined active/passive immunotherapy for epithelial tumors. <i>Journal of the National Cancer Institute</i> , 2004 , 96, 1611-9	9.7	51
29	Changes to peptide structure, not concentration, contribute to expansion of the lowest avidity cytotoxic T lymphocytes. <i>Journal of Leukocyte Biology</i> , 2004 , 76, 787-95	6.5	7
28	Human growth hormone presented by K14hGH-transgenic skin grafts induces a strong immune response but no graft rejection. <i>Immunology and Cell Biology</i> , 2004 , 82, 577-86	5	6
27	Tumour susceptibility to innate and adaptive immunotherapy changes during tumour maturation. <i>Immunology and Cell Biology</i> , 2004 , 82, 455-61	5	3
26	Functional memory CD8+ T cells can be generated in vivo without evident T help. <i>Vaccine</i> , 2004 , 23, 739-42	4.1	0
25	IL-10 mediates suppression of the CD8 T cell IFN-gamma response to a novel viral epitope in a primed host. <i>Journal of Immunology</i> , 2003 , 171, 4765-72	5.3	46
24	Expression of the HPV16E7 oncoprotein by thymic epithelium is accompanied by disrupted T cell maturation and a failure of the thymus to involute with age. <i>Clinical and Developmental Immunology</i> , 2003 , 10, 91-103		6
23	Paucity of functional CTL epitopes in the E7 oncoprotein of cervical cancer associated human papillomavirus type 16. <i>Immunology and Cell Biology</i> , 2003 , 81, 1-7	5	20
22	Inhibition of early tumor growth requires J alpha 18-positive (natural killer T) cells. <i>Cancer Research</i> , 2003 , 63, 3058-60	10.1	39
21	The number of long-lasting functional memory CD8+ T cells generated depends on the nature of the initial nonspecific stimulation. <i>European Journal of Immunology</i> , 2002 , 32, 1541-9	6.1	41
20	Route of administration of chimeric BPV1 VLP determines the character of the induced immune responses. <i>Immunology and Cell Biology</i> , 2002 , 80, 21-9	5	24
19	Interferon-gamma enhances cytotoxic T lymphocyte recognition of endogenous peptide in keratinocytes without lowering the requirement for surface peptide. <i>Immunology and Cell Biology</i> , 2002 , 80, 415-24	5	24
18	Chimeric human papilloma virus-simian/human immunodeficiency virus virus-like-particle vaccines: immunogenicity and protective efficacy in macaques. <i>Virology</i> , 2002 , 301, 176-87	3.6	56
17	Antigen-specific CD4+ T-cell help is required to activate a memory CD8+ T cell to a fully functional tumor killer cell. <i>Cancer Research</i> , 2002 , 62, 6438-41	10.1	105
16	Nonspecific down-regulation of CD8+ T-cell responses in mice expressing human papillomavirus type 16 E7 oncoprotein from the keratin-14 promoter. <i>Journal of Virology</i> , 2001 , 75, 5985-97	6.6	17

15	Tolerance or immunity to a tumor antigen expressed in somatic cells can be determined by systemic proinflammatory signals at the time of first antigen exposure. <i>Journal of Immunology</i> , 2001 , 167, 6180-7	5.3	43
14	Polynucleotide viral vaccines: codon optimisation and ubiquitin conjugation enhances prophylactic and therapeutic efficacy. <i>Vaccine</i> , 2001 , 20, 862-9	4.1	60
13	Papillomavirus virus-like particles for the delivery of multiple cytotoxic T cell epitopes. <i>Virology</i> , 2000 , 273, 374-82	3.6	48
12	Cytotoxic T-cell adherence assay (CAA). <i>Methods in Molecular Biology</i> , 2000 , 134, 277-81	1.4	
11	Potential strategies utilised by papillomavirus to evade host immunity. <i>Immunological Reviews</i> , 1999 , 168, 131-42	11.3	81
10	Split tolerance to a viral antigen expressed in thymic epithelium and keratinocytes. <i>European Journal of Immunology</i> , 1998 , 28, 2791-800	6.1	33
9	Cytotoxic T lymphocyte (CTL) adherence assay (CAA): a non-radioactive assay for murine CTL recognition of peptide-MHC class I complexes. <i>Journal of Immunological Methods</i> , 1997 , 201, 1-10	2.5	1
8	Cloning and expression of a cDNA encoding a nonintegrin laminin-binding protein from <i>Echinococcus granulosus</i> with localization of the laminin-binding domain. <i>Molecular and Biochemical Parasitology</i> , 1997 , 87, 183-92	1.9	11
7	Inactivation of human immunodeficiency virus (HIV)-1 envelope-specific CD8+ cytotoxic T lymphocytes by free antigenic peptide: a self-veto mechanism?. <i>Journal of Experimental Medicine</i> , 1996 , 183, 879-89	16.6	51
6	Selective expansion of high- or low-avidity cytotoxic T lymphocytes and efficacy for adoptive immunotherapy. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1996 , 93, 4102-7	11.5	452
5	Further characterization of the 38 kDa antigen from <i>Echinococcus granulosus</i> (hydatid disease) cyst fluid: evidence for antigenic heterogeneity and reactivity with anti-P1 antibodies. <i>Parasite Immunology</i> , 1995 , 17, 287-96	2.2	8
4	Identification and diagnostic value of a major antibody epitope on the 12 kDa antigen from <i>Echinococcus granulosus</i> (hydatid disease) cyst fluid. <i>Parasite Immunology</i> , 1994 , 16, 87-96	2.2	31
3	Hydatid immunoblot test and cross-reactivity with sera from patients with cysticercosis. <i>Transactions of the Royal Society of Tropical Medicine and Hygiene</i> , 1993 , 87, 350	2	2
2	Sequence homology between two immunodiagnostic fusion proteins from <i>Echinococcus multilocularis</i> . <i>International Journal for Parasitology</i> , 1992 , 22, 831-3	4.3	3
1	Serological evaluation of the 12 kDa subunit of antigen B in <i>Echinococcus granulosus</i> cyst fluid by immunoblot analysis. <i>Transactions of the Royal Society of Tropical Medicine and Hygiene</i> , 1992 , 86, 189-92 ²		49