

Cárdenas Mondragón María Guadalupe

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4353937/publications.pdf>

Version: 2024-02-01

18
papers

463
citations

759233

12
h-index

794594

19
g-index

19
all docs

19
docs citations

19
times ranked

578
citing authors

#	ARTICLE	IF	CITATIONS
1	Detection of Epstein-Barr Virus DNA in Gastric Biopsies of Pediatric Patients with Dyspepsia. <i>Pathogens</i> , 2020, 9, 623.	2.8	7
2	Differential expression of coxsackievirus and adenovirus receptor in endomyocardial tissue of patients with myocarditis. <i>Molecular Medicine Reports</i> , 2019, 20, 2189-2198.	2.4	5
3	Elevated Levels of Interferon- γ Are Associated with High Levels of Epstein-Barr Virus Reactivation in Patients with the Intestinal Type of Gastric Cancer. <i>Journal of Immunology Research</i> , 2017, 2017, 1-10.	2.2	15
4	Transcriptional Profiling of Type II Toxin-Antitoxin Genes of <i>Helicobacter pylori</i> under Different Environmental Conditions: Identification of HP0967-HP0968 System. <i>Frontiers in Microbiology</i> , 2016, 7, 1872.	3.5	33
5	Epstein-Barr Virus Association with Peptic Ulcer Disease. <i>Analytical Cellular Pathology</i> , 2015, 2015, 1-7.	1.4	16
6	Case-control study of Epstein-Barr virus and <i>Helicobacter pylori</i> serology in Latin American patients with gastric disease. <i>British Journal of Cancer</i> , 2015, 112, 1866-1873.	6.4	44
7	Preeclampsia Is Associated with Lower Production of Vascular Endothelial Growth Factor by Peripheral Blood Mononuclear Cells. <i>Archives of Medical Research</i> , 2014, 45, 561-569.	3.3	13
8	Epstein Barr Virus and <i>Helicobacter pylori</i> Co-Infection Are Positively Associated with Severe Gastritis in Pediatric Patients. <i>PLoS ONE</i> , 2013, 8, e62850.	2.5	70
9	Urinary Prolactin as a Reliable Marker for Preeclampsia, Its Severity, and the Occurrence of Adverse Pregnancy Outcomes. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2008, 93, 2492-2499.	3.6	68
10	Anti-prolactin autoantibodies in pregnant women with systemic lupus erythematosus: maternal and fetal outcome. <i>Lupus</i> , 2007, 16, 342-349.	1.6	21
11	Protein:Creatinine Ratio in Random Urine Samples Is a Reliable Marker of Increased 24-Hour Protein Excretion in Hospitalized Women with Hypertensive Disorders of Pregnancy. <i>Clinical Chemistry</i> , 2007, 53, 1623-1628.	3.2	64
12	Application of new homologous in vitro bioassays for human lactogens to assess the actual bioactivity of human prolactin isoforms in hyperprolactinaemic patients. <i>Clinical Endocrinology</i> , 2006, 65, 146-153.	2.4	24
13	Relation of Antimyocardium Antibodies to Mortality in Patients with Acute Myocardial Infarction. <i>Archives of Medical Research</i> , 2006, 37, 517-521.	3.3	1
14	Serum free prolactin concentrations in patients with systemic lupus erythematosus are associated with lupus activity. <i>Rheumatology</i> , 2006, 45, 97-101.	1.9	57
15	Immunization with a tetramer derivative of an anti-inflammatory pentapeptide produced by <i>Entamoeba histolytica</i> protects gerbils (<i>Meriones unguiculatus</i>) against experimental amoebic abscess of the liver. <i>Parasite Immunology</i> , 2004, 26, 343-349.	1.5	16
16	Eosinopenia Induced by Anti-IL-5 Antibody, but Not <i>Eimeria nieschulzi</i> Extract, Increases Susceptibility to Experimental Amebic Abscess of the Liver (EAAL) in Gerbils. <i>Archives of Medical Research</i> , 2002, 33, 316.	3.3	2
17	Increased Susceptibility to Experimental Amebic Abscess of the Liver (EAAL) in Gerbils (<i>Meriones</i>) Tj ETQq1 1 0.784314 rgBT /Overlock 1	3.3	2
18	Induction of Protective Immunity Against Experimental Amebic Abscess of the Liver (EAAL) Using the Monocyte Locomotion Inhibitory Factor (MLIF) Produced by <i>Entamoeba histolytica</i> . <i>Archives of Medical Research</i> , 2000, 31, S110-S111.	3.3	1