

Veronique Baud

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4354441/publications.pdf>

Version: 2024-02-01

56
papers

7,283
citations

156536

32
h-index

162838

57
g-index

59
all docs

59
docs citations

59
times ranked

14264
citing authors

#	ARTICLE	IF	CITATIONS
1	The alternative RelB NF- κ B subunit is a novel critical player in diffuse large B-cell lymphoma. <i>Blood</i> , 2022, 139, 384-398.	0.6	29
2	Differential Expression of Genes Involved in Metabolism and Immune Response in Diffuse and Intestinal Gastric Cancers, a Pilot Ptudy. <i>Biomedicines</i> , 2022, 10, 240.	1.4	1
3	The Alternative RelB NF- κ B Subunit Exerts a Critical Survival Function upon Metabolic Stress in Diffuse Large B-Cell Lymphoma-Derived Cells. <i>Biomedicines</i> , 2022, 10, 348.	1.4	5
4	Ultrahigh-Resolution NMR with Water Signal Suppression for a Deeper Understanding of the Action of Antimetabolic Drugs on Diffuse Large B-Cell Lymphoma. <i>Journal of Proteome Research</i> , 2022, 21, 1041-1051.	1.8	9
5	c-Rel Is the Pivotal NF- κ B Subunit in Germinal Center Diffuse Large B-Cell Lymphoma: A LYSA Study. <i>Frontiers in Oncology</i> , 2021, 11, 638897.	1.3	7
6	Tumor necrosis factor receptor family costimulation increases regulatory T cell activation and function via NF- κ B. <i>European Journal of Immunology</i> , 2020, 50, 972-985.	1.6	55
7	NF- κ B in the New Era of Cancer Therapy. <i>Trends in Cancer</i> , 2020, 6, 677-687.	3.8	49
8	The NF- κ B RelA Transcription Factor Is Critical for Regulatory T Cell Activation and Stability. <i>Frontiers in Immunology</i> , 2019, 10, 2487.	2.2	35
9	AXL Targeting Overcomes Human Lung Cancer Cell Resistance to NK- and CTL-Mediated Cytotoxicity. <i>Cancer Immunology Research</i> , 2019, 7, 1789-1802.	1.6	52
10	Cutting Edge: NANOG Activates Autophagy under Hypoxic Stress by Binding to BNIP3L Promoter. <i>Journal of Immunology</i> , 2017, 198, 1423-1428.	0.4	36
11	Molecular analysis of vascular smooth muscle cells from patients with giant cell arteritis: Targeting endothelin-1 receptor to control proliferation. <i>Autoimmunity Reviews</i> , 2017, 16, 398-406.	2.5	28
12	Model of the Interaction between the NF- κ B Inhibitory Protein p100 and the E3 Ubiquitin Ligase κ 2-TrCP based on NMR and Docking Experiments. <i>Journal of Chemical Information and Modeling</i> , 2017, 57, 223-233.	2.5	7
13	Post-Translational Modifications of RelB NF- κ B Subunit and Associated Functions. <i>Cells</i> , 2016, 5, 22.	1.8	38
14	Renal Cell Carcinoma Programmed Death-ligand 1, a New Direct Target of Hypoxia-inducible Factor-2 Alpha, is Regulated by von Hippel-Lindau Gene Mutation Status. <i>European Urology</i> , 2016, 70, 623-632.	0.9	115
15	Astrocyte Transcriptome from the Mecp2308-Truncated Mouse Model of Rett Syndrome. <i>NeuroMolecular Medicine</i> , 2015, 17, 353-363.	1.8	31
16	Rapid childhood T-ALL growth in xenograft models correlates with mature phenotype and NF- κ B pathway activation but not with poor prognosis. <i>Leukemia</i> , 2015, 29, 977-980.	3.3	10
17	ITPR1 Protects Renal Cancer Cells against Natural Killer Cells by Inducing Autophagy. <i>Cancer Research</i> , 2014, 74, 6820-6832.	0.4	97
18	Thrombopoietin promotes NHEJ DNA repair in hematopoietic stem cells through specific activation of Erk and NF- κ B pathways and their target, IEX-1. <i>Blood</i> , 2014, 123, 509-519.	0.6	51

#	ARTICLE	IF	CITATIONS
19	Interleukin-18 produced by bone marrow-derived stromal cells supports T-cell acute leukaemia progression. <i>EMBO Molecular Medicine</i> , 2014, 6, 821-834.	3.3	29
20	GH Administration Rescues Fatty Liver Regeneration Impairment by Restoring GH/EGFR Pathway Deficiency. <i>Endocrinology</i> , 2014, 155, 2545-2554.	1.4	40
21	IKK phosphorylates RelB to modulate its promoter specificity and promote fibroblast migration downstream of TNF receptors. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014, 111, 14794-14799.	3.3	22
22	RelB inhibits cell proliferation and tumor growth through p53 transcriptional activation. <i>Oncogene</i> , 2013, 32, 2661-2669.	2.6	22
23	A Mendelian predisposition to B-cell lymphoma caused by IL-10R deficiency. <i>Blood</i> , 2013, 122, 3713-3722.	0.6	116
24	Phagocytosis of apoptotic cells expressing PR3 impaired macrophage anti-inflammatory reprogramming. <i>Presse Medicale</i> , 2013, 42, 656.	0.8	0
25	Cutting Edge: Hypoxia-Induced Nanog Favors the Intratumoral Infiltration of Regulatory T Cells and Macrophages via Direct Regulation of TGF- β 1. <i>Journal of Immunology</i> , 2013, 191, 5802-5806.	0.4	97
26	Frequent Engagement of RelB Activation Is Critical for Cell Survival in Multiple Myeloma. <i>PLoS ONE</i> , 2013, 8, e59127.	1.1	29
27	A Novel Cellular Model to Study Angiotensin II AT2 Receptor Function in Breast Cancer Cells. <i>International Journal of Peptides</i> , 2012, 12, 1-6.	0.7	6
28	Autophagy is required for the activation of NF- κ B. <i>Cell Cycle</i> , 2012, 11, 194-199.	1.3	107
29	Antineoplastic activity of ouabain and pyriithione zinc in acute myeloid leukemia. <i>Oncogene</i> , 2012, 31, 3536-3546.	2.6	57
30	Oncogenic β -catenin triggers an inflammatory response that determines the aggressiveness of hepatocellular carcinoma in mice. <i>Journal of Clinical Investigation</i> , 2012, 122, 586-599.	3.9	155
31	Attenuation of Soft-Tissue Sarcomas Resistance to the Cytotoxic Action of TNF- α by Restoring p53 Function. <i>PLoS ONE</i> , 2012, 7, e38808.	1.1	8
32	Similar NF- κ B Gene Signatures in TNF- α Treated Human Endothelial Cells and Breast Tumor Biopsies. <i>PLoS ONE</i> , 2011, 6, e21589.	1.1	37
33	Deregulation of Aiolos expression in chronic lymphocytic leukemia is associated with epigenetic modifications. <i>Blood</i> , 2011, 117, 1917-1927.	0.6	38
34	Alternatively spliced NKp30 isoforms affect the prognosis of gastrointestinal stromal tumors. <i>Nature Medicine</i> , 2011, 17, 700-707.	15.2	282
35	GH Receptor Plays a Major Role in Liver Regeneration through the Control of EGFR and ERK1/2 Activation. <i>Endocrinology</i> , 2011, 152, 2731-2741.	1.4	45
36	A20-binding Inhibitor of Nuclear Factor- κ B (NF- κ B)-2 (ABIN-2) Is an Activator of Inhibitor of NF- κ B (I κ B) Kinase β (IKK β)-mediated NF- κ B Transcriptional Activity. <i>Journal of Biological Chemistry</i> , 2011, 286, 32277-32288.	1.6	28

#	ARTICLE	IF	CITATIONS
37	IKK connects autophagy to major stress pathways. <i>Autophagy</i> , 2010, 6, 189-191.	4.3	46
38	Î²B kinase overcomes PI3K/Akt and ERK/MAPK to control FOXO3a activity in acute myeloid leukemia. <i>Blood</i> , 2010, 116, 4240-4250.	0.6	69
39	Dichotomy between factors inducing the immunosuppressive enzyme IL-4-induced gene 1 (IL41) in B lymphocytes and mononuclear phagocytes. <i>European Journal of Immunology</i> , 2010, 40, 2557-2568.	1.6	61
40	The IKK complex contributes to the induction of autophagy. <i>EMBO Journal</i> , 2010, 29, 619-631.	3.5	274
41	Control of NF-Î²B Activity by Proteolysis. <i>Current Topics in Microbiology and Immunology</i> , 2010, 349, 97-114.	0.7	19
42	Is NF-Î²B a good target for cancer therapy? Hopes and pitfalls. <i>Nature Reviews Drug Discovery</i> , 2009, 8, 33-40.	21.5	913
43	RelA repression of RelB activity induces selective gene activation downstream of TNF receptors. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 14635-14640.	3.3	97
44	Identification and characterization of p100HB, a new mutant form of p100/NF-Î²B2. <i>Biochemical and Biophysical Research Communications</i> , 2003, 308, 744-749.	1.0	18
45	RelB/p50 Dimers Are Differentially Regulated by Tumor Necrosis Factor-Î± and Lymphotoxin-Î² Receptor Activation. <i>Journal of Biological Chemistry</i> , 2003, 278, 23278-23284.	1.6	139
46	IKKÎ² Is Essential for Protecting T Cells from TNFÎ±-Induced Apoptosis. <i>Immunity</i> , 2001, 14, 217-230.	6.6	198
47	Oxidative stress and gene expression: The AP-1 and NF-Î²B connections. <i>BioFactors</i> , 2001, 15, 87-89.	2.6	145
48	IKKÎ± controls formation of the epidermis independently of NF-Î²B. <i>Nature</i> , 2001, 410, 710-714.	13.7	335
49	Signal transduction by tumor necrosis factor and its relatives. <i>Trends in Cell Biology</i> , 2001, 11, 372-377.	3.6	1,475
50	Abnormal Morphogenesis But Intact IKK Activation in Mice Lacking the IKK Subunit of IB Kinase. <i>Science</i> , 1999, 284, 316-320.	6.0	799
51	Signaling by proinflammatory cytokines: oligomerization of TRAF2 and TRAF6 is sufficient for JNK and IKK activation and target gene induction via an amino-terminal effector domain. <i>Genes and Development</i> , 1999, 13, 1297-1308.	2.7	422
52	PAK4, a novel effector for Cdc42Hs, is implicated in the reorganization of the actin cytoskeleton and in the formation of filopodia. <i>EMBO Journal</i> , 1998, 17, 6527-6540.	3.5	317
53	EMR1, an unusual member in the family of hormone receptors with seven transmembrane segments. <i>Genomics</i> , 1995, 26, 334-344.	1.3	109
54	The E subunit of vacuolar H ⁺ -ATPase localizes close to the centromere on human chromosome 22. <i>Human Molecular Genetics</i> , 1994, 3, 335-339.	1.4	17

#	ARTICLE	IF	CITATIONS
55	Isolation of cosmids and fetal brain cDNAs from the proximal long arm of human chromosome 22. Human Molecular Genetics, 1993, 2, 535-540.	1.4	8
56	The human homolog of the mouse common viral integration region, FLI1, maps to 11q23-q24. Genomics, 1991, 11, 223-224.	1.3	17