

WÅ,adysÅ,aw Cabaj

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4357295/publications.pdf>

Version: 2024-02-01

29
papers

485
citations

623734

14
h-index

713466

21
g-index

29
all docs

29
docs citations

29
times ranked

538
citing authors

#	ARTICLE	IF	CITATIONS
1	The parasitic fauna of the European bison (<i>Bison bonasus</i>) (Linnaeus, 1758) and their impact on the conservation. Part 1 The summarising list of parasites noted. <i>Acta Parasitologica</i> , 2014, 59, 363-71.	1.1	48
2	Comparative analysis of excretory-secretory antigens of <i>Trichinella spiralis</i> and <i>Trichinella britovi</i> muscle larvae by two-dimensional difference gel electrophoresis and immunoblotting. <i>Proteome Science</i> , 2012, 10, 10.	1.7	34
3	Proteomic analysis of potential immunoreactive proteins from muscle larvae and adult worms of <i>Trichinella spiralis</i> in experimentally infected pigs. <i>Folia Parasitologica</i> , 2015, 62, .	1.3	34
4	Seroprevalence of <i>Toxoplasma gondii</i> and <i>Neospora caninum</i> infection in sheep, goats, and fallow deer farmed on the same area. <i>Journal of Animal Science</i> , 2018, 96, 2468-2473.	0.5	33
5	Antibodies to <i>Neospora caninum</i> in the blood of European bison (<i>Bison bonasus bonasus</i> L.) living in Poland. <i>Veterinary Parasitology</i> , 2005, 128, 163-168.	1.8	28
6	The parasitic fauna of the European bison (<i>Bison bonasus</i>) (Linnaeus, 1758) and their impact on the conservation. Part 2 The structure and changes over time. <i>Acta Parasitologica</i> , 2014, 59, 372-9.	1.1	24
7	Prevalence of antibodies against <i>Neospora caninum</i> in dogs from urban areas in Central Poland. <i>Parasitology Research</i> , 2011, 108, 991-996.	1.6	23
8	The first detection of <i>Neospora caninum</i> DNA in the colostrum of infected cows. <i>Parasitology Research</i> , 2006, 100, 633-636.	1.6	22
9	First report of <i>Trichinella pseudospiralis</i> in Poland, in red foxes (<i>Vulpes vulpes</i>). <i>Acta Parasitologica</i> , 2013, 58, 149-54.	1.1	22
10	Multilocus genotyping of <i>Giardia duodenalis</i> isolates from red deer (<i>Cervus elaphus</i>) and roe deer (<i>Capreolus capreolus</i>) from Poland. <i>Folia Parasitologica</i> , 2012, 59, 237-240.	1.3	22
11	In vitro isolation and identification of the first <i>Neospora caninum</i> isolate from European bison (<i>Bison</i>) Tj ETQq1 1 0.784314 rgBT /Overlock	1.8	20
12	Molecular identification of <i>Trichinella britovi</i> in martens (<i>Martes martes</i>) and badgers (<i>Meles meles</i>); new host records in Poland. <i>Acta Parasitologica</i> , 2012, 57, 402-5.	1.1	20
13	The first identification of a blood-sucking abomasal nematode <i>Ashworthius sidemi</i> in cattle (<i>Bos</i>) Tj ETQq1 1 0.784314 rgBT /Overlock	1.8	19
14	Susceptibility of adult <i>Heligmosomoides polygyrus</i> to intestinal inflammatory responses induced by heterologous infection. <i>International Journal for Parasitology</i> , 1992, 22, 75-86.	3.1	17
15	The Occurrence of <i>Trichinella</i> spp. in Red Foxes (<i>Vulpes vulpes</i>) in Different Regions of Poland: Current Data. <i>Vector-Borne and Zoonotic Diseases</i> , 2016, 16, 717-721.	1.5	15
16	Wild boars meat as a potential source of human trichinellosis in Poland: current data. <i>Acta Parasitologica</i> , 2015, 60, 530-5.	1.1	14
17	The occurrence of nematodes of the genus <i>Trichinella</i> in wolves (<i>Canis lupus</i>) from the Bieszczady Mountains and Augustowska Forest in Poland. <i>Veterinary Parasitology</i> , 2016, 231, 115-117.	1.8	14
18	Molecular identification of <i>Trichinella spiralis</i> and <i>Trichinella britovi</i> by diagnostic multiprimer large mitochondrial rRNA amplification. <i>Parasitology Research</i> , 2003, 91, 374-377.	1.6	10

#	ARTICLE	IF	CITATIONS
19	Recognition of antigens of three different stages of the <i>Trichinella spiralis</i> by antibodies from pigs infected with <i>T. spiralis</i> . <i>Experimental Parasitology</i> , 2013, 134, 129-137.	1.2	10
20	The first report of <i>Toxoplasma gondii</i> antibodies in free-living European bison (<i>Bison bonasus bonasus</i>) Tj ETQq0 0 Q ggBT /Overlock 10 T	1.3	10
21	Characterization of the first Polish isolate of <i>Neospora caninum</i> from cattle. <i>Acta Parasitologica</i> , 2007, 52, 295.	1.1	7
22	Detection of antibodies to <i>Neospora caninum</i> in moose (<i>Alces alces</i>): the first report in Europe. <i>Folia Parasitologica</i> , 2014, 61, 34-36.	1.3	7
23	Studies on <i>Neospora caninum</i> DNA detection in the oocytes and embryos collected from infected cows. <i>Veterinary Parasitology</i> , 2008, 158, 370-375.	1.8	6
24	The usefulness of DNA derived from third stage larvae in the detection of <i>Ashworthius sidemi</i> infection in European bison, by a simple polymerase chain reaction. <i>Parasites and Vectors</i> , 2014, 7, 215.	2.5	6
25	First <i>Toxoplasma gondii</i> isolate from an aborted foetus of European bison (<i>Bison bonasus bonasus</i> L.). <i>Parasitology Research</i> , 2017, 116, 2457-2461.	1.6	6
26	The first report of <i>Toxoplasma gondii</i> antibodies in free-living European bison (<i>Bison bonasus bonasus</i>) Tj ETQq0 0 Q ggBT /Overlock 10 T	1.3	6
27	<i>Sarcocystis cruzi</i> infection in free-living European bison (<i>Bison bonasus bonasus</i> L.) from the Bia&owie&14a Forest, Poland â&€“ A molecular analysis based on the <i>cox1</i> gene. <i>International Journal for Parasitology: Parasites and Wildlife</i> , 2021, 16, 59-63.	1.5	3
28	Detection of antibodies to <i>Neospora caninum</i> in moose (<i>Alces alces</i>): the first report in Europe. <i>Folia Parasitologica</i> , 2014, 61, 34-6.	1.3	3
29	Molecular identification of sarcocysts from tissue of fallow deer (<i>Dama dama</i>) farmed in the open pasture system based on <i>ssu rRNA</i> gene. <i>Acta Parasitologica</i> , 2020, 65, 354-360.	1.1	2