Anna WaÅ>kiewicz

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/43666/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Prevalence of general and abdominal obesity and overweight among adults in Poland. Results of the WOBASZ II study (2013–2014) and comparison with the WOBASZ study (2003–2005). Polish Archives of Internal Medicine, 2016, 126, 662-671.	0.3	53
2	Are dietary habits of the Polish population consistent with the recommendations for prevention of cardiovascular disease? — WOBASZ II project. Kardiologia Polska, 2016, 74, 969-977.	0.3	45
3	Comparison of Various Databases for Estimation of Dietary Polyphenol Intake in the Population of Polish Adults. Nutrients, 2015, 7, 9299-9308.	1.7	41
4	Dietary Total Antioxidant Capacity and Dietary Polyphenol Intake and Prevalence of Metabolic Syndrome in Polish Adults: A Nationwide Study. Oxidative Medicine and Cellular Longevity, 2018, 2018, 1-10.	1.9	32
5	Dietary Antioxidant and Flavonoid Intakes Are Reduced in the Elderly. Oxidative Medicine and Cellular Longevity, 2015, 2015, 1-8.	1.9	27
6	Are Total and Individual Dietary Lignans Related to Cardiovascular Disease and Its Risk Factors in Postmenopausal Women? A Nationwide Study. Nutrients, 2018, 10, 865.	1.7	25
7	Dietary Polyphenol Intake, but Not the Dietary Total Antioxidant Capacity, Is Inversely Related to Cardiovascular Disease in Postmenopausal Polish Women: Results of WOBASZ and WOBASZ II Studies. Oxidative Medicine and Cellular Longevity, 2017, 2017, 1-11.	1.9	22
8	Socioeconomic factors and the risk of metabolic syndrome in the adult Polish population: the WOBASZ study. Kardiologia Polska, 2012, 70, 718-27.	0.3	22
9	Dietary acid load and cardiometabolic risk in the Polish adult population. Advances in Clinical and Experimental Medicine, 2018, 27, 1347-1345.	0.6	21
10	Dietary Habits and Dietary Antioxidant Intake Are Related to Socioeconomic Status in Polish Adults: A Nationwide Study. Nutrients, 2020, 12, 518.	1.7	20
11	Metabolic syndrome and its components in Polish women of childbearing age: a nationwide study. BMC Public Health, 2018, 18, 15.	1.2	17
12	Dietary antioxidant capacity of the patients with cardiovascular disease in a cross-sectional study. Nutrition Journal, 2015, 14, 26.	1.5	16
13	Polyphenols and dietary antioxidant potential, and their relationship with arterial hypertension: A cross-sectional study of theadult population in Poland (WOBASZ II). Advances in Clinical and Experimental Medicine, 2019, 28, 797-806.	0.6	13
14	Assessment of Plant Sterols in the Diet of Adult Polish Population with the Use of a Newly Developed Database. Nutrients, 2021, 13, 2722.	1.7	12
15	Dietary Plant Sterols and Phytosterol-Enriched Margarines and Their Relationship with Cardiovascular Disease among Polish Men and Women: The WOBASZ II Cross-Sectional Study. Nutrients, 2022, 14, 2665.	1.7	11
16	The Consumption of Nuts is Associated with Better Dietary and Lifestyle Patterns in Polish Adults: Results of WOBASZ and WOBASZ II Surveys. Nutrients, 2019, 11, 1410.	1.7	10
17	Diet quality in the population of Norway and Poland: differences in the availability and consumption of food considering national nutrition guidelines and food market. BMC Public Health, 2021, 21, 319.	1.2	9
18	Relationship between the dietary glycemic load of the adult Polish population and socio-demographic and lifestyle factors – results of the WOBASZ II study. Advances in Clinical and Experimental Medicine, 2019, 28, 891-897.	0.6	5

#	Article	IF	CITATIONS
19	Metabolic disorders in women at procreative age living in Warsaw. Roczniki Panstwowego Zakladu Higieny, 2015, 66, 245-51.	0.5	3
20	Differences in food consumption in regions of Poland that had been previously under Austrian, Prussian and Russian annexations. Roczniki Panstwowego Zakladu Higieny, 2019, 70, 287-294.	0.5	0