

# Liubing Li

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4367164/publications.pdf>

Version: 2024-02-01

39  
papers

675  
citations

759233

12  
h-index

610901

24  
g-index

39  
all docs

39  
docs citations

39  
times ranked

1188  
citing authors

#	ARTICLE	IF	CITATIONS
1	Ferritin in the coronavirus disease 2019 (COVID-19): A systematic review and meta-analysis. <i>Journal of Clinical Laboratory Analysis</i> , 2020, 34, e23618.	2.1	244
2	Anti-MDA5 antibody as a potential diagnostic and prognostic biomarker in patients with dermatomyositis. <i>Oncotarget</i> , 2017, 8, 26552-26564.	1.8	39
3	Syndecan-1, an indicator of endothelial glycocalyx degradation, predicts outcome of patients admitted to an ICU with COVID-19. <i>Molecular Medicine</i> , 2021, 27, 151.	4.4	36
4	Assessment of anti-MDA5 antibody as a diagnostic biomarker in patients with dermatomyositis-associated interstitial lung disease or rapidly progressive interstitial lung disease. <i>Oncotarget</i> , 2017, 8, 76129-76140.	1.8	31
5	Meta-Analysis: Diagnostic Accuracy of Anti-Carbamylated Protein Antibody for Rheumatoid Arthritis. <i>PLoS ONE</i> , 2016, 11, e0159000.	2.5	28
6	Myositis-specific autoantibodies in dermatomyositis/polymyositis with interstitial lung disease. <i>Journal of the Neurological Sciences</i> , 2019, 397, 123-128.	0.6	28
7	Evaluation of the diagnostic potential of antibodies to beta2-glycoprotein 1 domain 1 in Chinese patients with antiphospholipid syndrome. <i>Scientific Reports</i> , 2016, 6, 23839.	3.3	23
8	Single-nucleotide rs738409 polymorphisms in the PNPLA3 gene are strongly associated with alcoholic liver disease in Han Chinese males. <i>Hepatology International</i> , 2018, 12, 429-437.	4.2	19
9	Assessment of diagnostic utility, clinical phenotypic associations, and prognostic significance of anti-NXP2 autoantibody in patients with idiopathic inflammatory myopathies: a systematic review and meta-analysis. <i>Clinical Rheumatology</i> , 2021, 40, 819-832.	2.2	16
10	Evaluation of the clinical relevance of anti-annexin-A5 antibodies in Chinese patients with antiphospholipid syndrome. <i>Clinical Rheumatology</i> , 2017, 36, 407-412.	2.2	14
11	Relationship of HLA-B*51 and HLA-B*52 alleles and TNF- $\alpha$ -308A/G polymorphism with susceptibility to Takayasu arteritis: a meta-analysis. <i>Clinical Rheumatology</i> , 2017, 36, 173-181.	2.2	14
12	Positive Association between ANKRD55 Polymorphism 7731626 and Dermatomyositis/Polymyositis with Interstitial Lung Disease in Chinese Han Population. <i>BioMed Research International</i> , 2017, 2017, 1-9.	1.9	14
13	High levels of antibodies to citrullinated $\alpha$ -enolase peptide-1 (CEP-1) identify erosions and interstitial lung disease (ILD) in a Chinese rheumatoid arthritis cohort. <i>Clinical Immunology</i> , 2019, 200, 10-15.	3.2	14
14	Evaluation of a commercial immunoassay for autoantibodies in Chinese Han systemic sclerosis population. <i>Clinica Chimica Acta</i> , 2019, 491, 121-125.	1.1	13
15	Genetic variant rs72613567 of <i>HSD17B13</i> gene reduces alcohol-related liver disease risk in Chinese Han population. <i>Liver International</i> , 2020, 40, 2194-2202.	3.9	13
16	Identification of Novel Serological Autoantibodies in Takayasu Arteritis Patients Using HuProt Arrays. <i>Molecular and Cellular Proteomics</i> , 2021, 20, 100036.	3.8	13
17	Association of FCGR2A/FCGR3A variant rs2099684 with Takayasu arteritis in the Han Chinese population. <i>Oncotarget</i> , 2017, 8, 17239-17245.	1.8	11
18	Association of GTF2I, NFKB1, and TYK2 Regional Polymorphisms With Systemic Sclerosis in a Chinese Han Population. <i>Frontiers in Immunology</i> , 2021, 12, 640083.	4.8	10

#	ARTICLE	IF	CITATIONS
19	Single nucleotide polymorphisms of IL12B are associated with Takayasu arteritis in Chinese Han population. <i>Rheumatology International</i> , 2017, 37, 547-555.	3.0	9
20	Meta-analysis of anti-Saccharomyces cerevisiae antibodies as diagnostic markers of Behçet's disease with gastrointestinal involvement. <i>BMJ Open</i> , 2020, 10, e033880.	1.9	9
21	Assessing serum IgG4 glycosylation profiles of IgG4-related disease using lectin microarray. <i>Clinical and Experimental Rheumatology</i> , 2021, 39, 393-402.	0.8	9
22	Clinical performance of antibodies to prothrombin and thrombin in Chinese patients with antiphospholipid syndrome: potential interest in discriminating patients with thrombotic events and non-thrombotic events. <i>Rheumatology International</i> , 2017, 37, 579-584.	3.0	7
23	Analysis of myositis autoantibodies in Chinese patients with cancer-associated myositis. <i>Journal of Clinical Laboratory Analysis</i> , 2020, 34, e23307.	2.1	7
24	Variation of red blood cell parameters in Behçet's disease: association with disease severity and vascular involvement. <i>Clinical Rheumatology</i> , 2021, 40, 1457-1464.	2.2	7
25	Association between genetic variants in the human leukocyte antigen-B/*MICA and Takayasu arteritis in Chinese Han population. <i>International Journal of Rheumatic Diseases</i> , 2018, 21, 271-277.	1.9	6
26	Analysis of anti-RNA polymerase III antibodies in Chinese Han systemic sclerosis patients. <i>Clinical Rheumatology</i> , 2020, 39, 1191-1197.	2.2	6
27	Single nucleotide polymorphisms in the ETS1 gene are associated with idiopathic inflammatory myopathies in a northern Chinese Han population. <i>Scientific Reports</i> , 2017, 7, 13128.	3.3	5
28	NAF1 rs4691896 Is Significantly Associated with Coal Workers' Pneumoconiosis in a Chinese Han Population: A Case-Control Study. <i>Medical Science Monitor</i> , 2020, 26, e918709.	1.1	5
29	Association between the BANK1 rs3733197 polymorphism and polymyositis/dermatomyositis in a Chinese Han population. <i>Clinical Rheumatology</i> , 2019, 38, 431-436.	2.2	4
30	Adult reference intervals for IgG subclasses with Siemens immunonephelometric assays in Chinese population. <i>Allergy, Asthma and Clinical Immunology</i> , 2017, 13, 44.	2.0	3
31	Peripheral differentials by Cytodiff flow cytometric system predict disease activity in Chinese patients with inflammatory bowel disease. <i>Clinica Chimica Acta</i> , 2017, 471, 17-22.	1.1	3
32	Comprehensive analysis of immunoglobulin and clinical variables identifies functional linkages and diagnostic indicators associated with Behçet's disease patients receiving immunomodulatory treatment. <i>BMC Immunology</i> , 2021, 22, 16.	2.2	3
33	Diagnostic value of anti-citrullinated Î±-enolase peptide 1 antibody in patients with rheumatoid arthritis: A systematic review and meta-analysis. <i>International Journal of Rheumatic Diseases</i> , 2021, 24, 633-646.	1.9	3
34	Serum C1q Concentration Positively Correlates with Erythrocyte Sedimentation Rate in Polymyositis/Dermatomyositis. <i>Annals of Clinical and Laboratory Science</i> , 2019, 49, 237-241.	0.2	3
35	Risk alleles for IgA nephropathy-associated SNPs conferred completely opposite effects to idiopathic membranous nephropathy in Chinese Han. <i>Immunologic Research</i> , 2017, 65, 1059-1064.	2.9	2
36	Assessing serum IgG4 glycosylation profiles of IgG4-related disease using lectin microarray. <i>Clinical and Experimental Rheumatology</i> , 2021, 39, 393-402.	0.8	2

#	ARTICLE	IF	CITATIONS
37	<i>GLIS3</i> and <i>TYK2</i> Single Nucleotide Polymorphisms Are Not Associated with Dermatomyositis/Polymyositis in Chinese Han Population. <i>Genetic Testing and Molecular Biomarkers</i> , 2017, 21, 565-570.	0.7	1
38	Establishing a reference interval for serum anti-dsDNA antibody: A large Chinese Han population-based multi-center study. <i>PLoS ONE</i> , 2017, 12, e0168871.	2.5	1
39	Prealbumin Correlates with C-reactive Protein in Rheumatic Musculoskeletal Diseases. <i>Annals of Clinical and Laboratory Science</i> , 2021, 51, 82-89.	0.2	0