

Peter Gmeiner

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/4371919/peter-gmeiner-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

289
papers

9,068
citations

45
h-index

81
g-index

319
ext. papers

10,213
ext. citations

5.7
avg, IF

5.95
L-index

#	Paper	IF	Citations
289	Structural insights into ligand recognition, activation, and signaling of the β adrenergic receptor.. <i>Science Advances</i> , 2022 , 8, eabj5347	14.3	1
288	Development of disulfide-functionalized peptides covalently binding G protein-coupled receptors.. <i>Bioorganic and Medicinal Chemistry</i> , 2022 , 61, 116720	3.4	
287	Activation and allosteric regulation of the orphan GPR88-Gi1 signaling complex.. <i>Nature Communications</i> , 2022 , 13, 2375	17.4	2
286	N-Terminus to Arginine Side-Chain Cyclization of Linear Peptidic Neuropeptide Y Y Receptor Ligands Results in Picomolar Binding Constants. <i>Journal of Medicinal Chemistry</i> , 2021 , 64, 16746-16769	8.3	0
285	Functional Reconstitution of Dopamine D2 Receptor into a Supported Model Membrane in a Nanometric Confinement. <i>Advanced Biology</i> , 2021 , 5, e2100636		
284	Dibenzodiazepinone-type muscarinic receptor antagonists conjugated to basic peptides: Impact of the linker moiety and unnatural amino acids on MR selectivity. <i>European Journal of Medicinal Chemistry</i> , 2021 , 213, 113159	6.8	2
283	Abolishing Dopamine D/D Receptor Affinity of Subtype-Selective Carbamoylguanidine-Type Histamine H Receptor Agonists. <i>Journal of Medicinal Chemistry</i> , 2021 , 64, 8684-8709	8.3	3
282	Visualization of β adrenergic receptor dynamics and differential localization in cardiomyocytes. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021 , 118,	11.5	7
281	Radiosynthesis and evaluation of F-labeled dopamine D-receptor ligands. <i>Nuclear Medicine and Biology</i> , 2021 , 92, 43-52	2.1	1
280	Binding pathway determines norepinephrine selectivity for the human β AR over β AR. <i>Cell Research</i> , 2021 , 31, 569-579	24.7	23
279	Presynaptic vesicular accumulation is required for antipsychotic efficacy in psychotic-like rats. <i>Journal of Psychopharmacology</i> , 2021 , 35, 65-77	4.6	2
278	Sensitization to amphetamine psychostimulant effect: A key role for ventral tegmental area neurotensin type 2 receptors and MAP kinase pathway. <i>Addiction Biology</i> , 2021 , 26, e13008	4.6	
277	Homobivalent Dopamine D Receptor Ligands Modulate the Dynamic Equilibrium of D Monomers and Homo- and Heterodimers. <i>ACS Chemical Biology</i> , 2021 , 16, 371-379	4.9	3
276	Functionally selective activation of the dopamine receptor D is mirrored by the protein expression profiles. <i>Scientific Reports</i> , 2021 , 11, 3501	4.9	1
275	Optimizing the Expression of Human Dopamine Receptors in. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	1
274	Bivalent ligands promote endosomal trafficking of the dopamine D3 receptor-neurotensin receptor 1 heterodimer. <i>Communications Biology</i> , 2021 , 4, 1062	6.7	2
273	Synthesis, Radiosynthesis and Biological Evaluation of Buprenorphine-Derived Phenylazocarboxamides as Novel μ Opioid Receptor Ligands. <i>ChemMedChem</i> , 2020 , 15, 1175-1186	3.7	3

272	An allosteric modulator binds to a conformational hub in the β -adrenergic receptor. <i>Nature Chemical Biology</i> , 2020 , 16, 749-755	11.7	16
271	Discovery of Novel Nonpeptidic PAR2 Ligands. <i>ACS Medicinal Chemistry Letters</i> , 2020 , 11, 1316-1323	4.3	3
270	Activation of the β -adrenoceptor by the sedative sympatholytic dexmedetomidine. <i>Nature Chemical Biology</i> , 2020 , 16, 507-512	11.7	20
269	Investigation of Inactive-State μ Opioid Receptor Homodimerization via Single-Molecule Microscopy Using New Antagonistic Fluorescent Probes. <i>Journal of Medicinal Chemistry</i> , 2020 , 63, 3596-3609	8.3	6
268	Regiospecific Introduction of Halogens on the 2-Aminobiphenyl Subunit Leading to Highly Potent and Selective M3 Muscarinic Acetylcholine Receptor Antagonists and Weak Inverse Agonists. <i>Journal of Medicinal Chemistry</i> , 2020 , 63, 4349-4369	8.3	3
267	Red-Emitting Dibenzodiazepinone Derivatives as Fluorescent Dualsteric Probes for the Muscarinic Acetylcholine M Receptor. <i>Journal of Medicinal Chemistry</i> , 2020 , 63, 4133-4154	8.3	7
266	Innenrücktitelbild: Selective and Wash-Resistant Fluorescent Dihydrocodeinone Derivatives Allow Single-Molecule Imaging of μ Opioid Receptor Dimerization (Angew. Chem. 15/2020). <i>Angewandte Chemie</i> , 2020 , 132, 6348-6348	3.6	1
265	Structure-based development of caged dopamine D/D receptor antagonists. <i>Scientific Reports</i> , 2020 , 10, 829	4.9	6
264	Fluorescent ligands for dopamine D/D receptors. <i>Scientific Reports</i> , 2020 , 10, 21842	4.9	4
263	Fluorescence Labeling of Neurotensin(8-13) via Arginine Residues Gives Molecular Tools with High Receptor Affinity. <i>ACS Medicinal Chemistry Letters</i> , 2020 , 11, 16-22	4.3	8
262	Selective and Wash-Resistant Fluorescent Dihydrocodeinone Derivatives Allow Single-Molecule Imaging of μ Opioid Receptor Dimerization. <i>Angewandte Chemie - International Edition</i> , 2020 , 59, 5958-5964	16.4	9
261	Selective and Wash-Resistant Fluorescent Dihydrocodeinone Derivatives Allow Single-Molecule Imaging of μ Opioid Receptor Dimerization. <i>Angewandte Chemie</i> , 2020 , 132, 6014-6020	3.6	3
260	Structure-based development of a subtype-selective orexin 1 receptor antagonist. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020 , 117, 18059-18067	11.5	18
259	Pharmacological Characterization of Low-to-Moderate Affinity Opioid Receptor Agonists and Brain Imaging with F-Labeled Derivatives in Rats. <i>Journal of Medicinal Chemistry</i> , 2020 , 63, 9484-9499	8.3	3
258	Rational design of agonists for bitter taste receptor TAS2R14: from modeling to bench and back. <i>Cellular and Molecular Life Sciences</i> , 2020 , 77, 531-542	10.3	18
257	Structure-based exploration of an allosteric binding pocket in the NTS1 receptor using bitopic NT(8-13) derivatives and molecular dynamics simulations. <i>Journal of Molecular Modeling</i> , 2019 , 25, 193	2	2
256	Development of covalent antagonists for α - and β -adrenergic receptors. <i>Bioorganic and Medicinal Chemistry</i> , 2019 , 27, 2959-2971	3.4	4
255	Conjugation of Short Peptides to Dibenzodiazepinone-Type Muscarinic Acetylcholine Receptor Ligands Determines MR Selectivity. <i>Journal of Medicinal Chemistry</i> , 2019 , 62, 5358-5369	8.3	8

254	Conformational Complexity and Dynamics in a Muscarinic Receptor Revealed by NMR Spectroscopy. <i>Molecular Cell</i> , 2019 , 75, 53-65.e7	17.6	31
253	Synthesis of pyrazolylvinyl ketones from furan derivatives. <i>Organic and Biomolecular Chemistry</i> , 2019 , 17, 4850-4855	3.9	9
252	Hybridization of β Adrenergic Agonists and Antagonists Confers G Protein Bias. <i>Journal of Medicinal Chemistry</i> , 2019 , 62, 5111-5131	8.3	5
251	Perquinoline AII: neuartige bakterielle Tetrahydroisochinoline mit einer bemerkenswerten Biosynthese. <i>Angewandte Chemie</i> , 2019 , 131, 13063-13068	3.6	
250	Perquinolines A-C: Unprecedented Bacterial Tetrahydroisoquinolines Involving an Intriguing Biosynthesis. <i>Angewandte Chemie - International Edition</i> , 2019 , 58, 12930-12934	16.4	5
249	Benzyl Phenylsemicarbazides: A Chemistry-Driven Approach Leading to G Protein-Biased Dopamine D Receptor Agonists with High Subtype Selectivity. <i>Journal of Medicinal Chemistry</i> , 2019 , 62, 9658-9679	8.3	4
248	Dual-Acting Cholinesterase-Human Cannabinoid Receptor 2 Ligands Show Pronounced Neuroprotection in Vitro and Overadditive and Disease-Modifying Neuroprotective Effects in Vivo. <i>Journal of Medicinal Chemistry</i> , 2019 , 62, 9078-9102	8.3	18
247	Differential allosteric modulation within dopamine DR - neurotensin NTS1R and DR - serotonin 5-HT ₂ receptor complexes gives bias to intracellular calcium signalling. <i>Scientific Reports</i> , 2019 , 9, 16312	4.9	11
246	Mast Cell Degranulation and Fibroblast Activation in the Morphine-induced Spinal Mass: Role of Mas-related G Protein-coupled Receptor Signaling. <i>Anesthesiology</i> , 2019 , 131, 132-147	4.3	11
245	Monitoring of the dopamine D ₂ receptor agonists hordenine and N-methyltyramine during the brewing process and in commercial beer samples. <i>Food Chemistry</i> , 2019 , 276, 745-753	8.5	11
244	Structure-guided development of selective M ₃ muscarinic acetylcholine receptor antagonists. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2018 , 115, 12046-12050	11.5	39
243	Derivatization of common antidepressant drugs increases inhibition of acid sphingomyelinase and reduces induction of phospholipidosis. <i>Journal of Neural Transmission</i> , 2018 , 125, 1837-1845	4.3	9
242	Identification of the Beer Component Hordenine as Food-Derived Dopamine D ₂ Receptor Agonist by Virtual Screening a 3D Compound Database. <i>Scientific Reports</i> , 2017 , 7, 44201	4.9	20
241	Discovery of G Protein-Biased Dopaminergics with a Pyrazolo[1,5-a]pyridine Substructure. <i>Journal of Medicinal Chemistry</i> , 2017 , 60, 2908-2929	8.3	38
240	Hydroxy-Substituted Heteroaryl piperazines: Novel Scaffolds for β Arrestin-Biased DR Agonists. <i>Journal of Medicinal Chemistry</i> , 2017 , 60, 4693-4713	8.3	18
239	Development of molecular tools based on the dopamine D receptor ligand FAUC 329 showing inhibiting effects on drug and food maintained behavior. <i>Bioorganic and Medicinal Chemistry</i> , 2017 , 25, 3491-3499	3.4	5
238	Potent haloperidol derivatives covalently binding to the dopamine D ₂ receptor. <i>Bioorganic and Medicinal Chemistry</i> , 2017 , 25, 5084-5094	3.4	6
237	Radiolabeled Dibenzodiazepinone-Type Antagonists Give Evidence of Dualsteric Binding at the M Muscarinic Acetylcholine Receptor. <i>Journal of Medicinal Chemistry</i> , 2017 , 60, 3314-3334	8.3	15

236	Structure-Based Design and Discovery of New M Receptor Agonists. <i>Journal of Medicinal Chemistry</i> , 2017 , 60, 9239-9250	8.3	16
235	Heterodimerization of Dibenzodiazepinone-Type Muscarinic Acetylcholine Receptor Ligands Leads to Increased MR Affinity and Selectivity. <i>ACS Omega</i> , 2017 , 2, 6741-6754	3.9	8
234	Arrestin biased dopamine D2 receptor partial agonists: Synthesis and pharmacological evaluation. <i>Bioorganic and Medicinal Chemistry</i> , 2017 , 25, 5613-5628	3.4	8
233	Structure-Guided Screening for Functionally Selective D Dopamine Receptor Ligands from a Virtual Chemical Library. <i>ACS Chemical Biology</i> , 2017 , 12, 2652-2661	4.9	27
232	Visualization of ligand-induced dopamine D and D receptor internalization by TIRF microscopy. <i>Scientific Reports</i> , 2017 , 7, 10894	4.9	10
231	[F]Fluorophenylazocarboxylates: Design and Synthesis of Potential Radioligands for Dopamine D3 and Opioid Receptor. <i>ACS Omega</i> , 2017 , 2, 8649-8659	3.9	12
230	Photochromic Dopamine Receptor Ligands Based on Dithienylethenes and Fulgides. <i>Chemistry - A European Journal</i> , 2017 , 23, 13423-13434	4.8	23
229	NTS2-selective neurotensin mimetics with tetrahydrofuran amino acids. <i>Bioorganic and Medicinal Chemistry</i> , 2017 , 25, 350-359	3.4	6
228	Theranostic Value of Multimers: Lessons Learned from Trimerization of Neurotensin Receptor Ligands and Other Targeting Vectors. <i>Pharmaceuticals</i> , 2017 , 10,	5.2	13
227	Structure-based discovery of opioid analgesics with reduced side effects. <i>Nature</i> , 2016 , 537, 185-190	50.4	547
226	Dibenzo[b,f][1,4]oxazepines and dibenzo[b,e]oxepines: Influence of the chlorine substitution pattern on the pharmacology at the HR, HR, 5-HTR and other selected GPCRs. <i>Pharmacological Research</i> , 2016 , 113, 610-625	10.2	7
225	Visualization and ligand-induced modulation of dopamine receptor dimerization at the single molecule level. <i>Scientific Reports</i> , 2016 , 6, 33233	4.9	66
224	Structure-guided development of heterodimer-selective GPCR ligands. <i>Nature Communications</i> , 2016 , 7, 12298	17.4	65
223	(18)F- and (68)Ga-Labeled Neurotensin Peptides for PET Imaging of Neurotensin Receptor 1. <i>Journal of Medicinal Chemistry</i> , 2016 , 59, 6480-92	8.3	25
222	Spontaneous butenolide ring formation of pregnane-21-O-malonyl hemiesters under mild reaction conditions is facilitated by the 14hydroxy group present in all natural cardenolides. <i>Tetrahedron</i> , 2016 , 72, 4556-4563	2.4	4
221	Development of Covalent Ligand-Receptor Pairs to Study the Binding Properties of Nonpeptidic Neurotensin Receptor 1 Antagonists. <i>ACS Chemical Biology</i> , 2016 , 11, 869-75	4.9	13
220	Mimicking of Arginine by Functionalized N-Carbamoylated Arginine As a New Broadly Applicable Approach to Labeled Bioactive Peptides: High Affinity Angiotensin, Neuropeptide Y, Neuropeptide FF, and Neurotensin Receptor Ligands As Examples. <i>Journal of Medicinal Chemistry</i> , 2016 , 59, 1925-45	8.3	19
219	Comparative MD Simulations Indicate a Dual Role for Arg1323.50 in Dopamine-Dependent D2R Activation. <i>PLoS ONE</i> , 2016 , 11, e0146612	3.7	8

218	Optimization and synthesis of an (18) F-labeled dopamine D3 receptor ligand using [(18) F]fluorophenylazocarboxylic tert-butylester. <i>Journal of Labelled Compounds and Radiopharmaceuticals</i> , 2016 , 59, 48-53	1.9	6
217	Structure-guided development of dual α adrenergic/dopamine D2 receptor agonists. <i>Bioorganic and Medicinal Chemistry</i> , 2016 , 24, 2641-53	3.4	10
216	Structural insights into μ -opioid receptor activation. <i>Nature</i> , 2015 , 524, 315-21	50.4	558
215	Multicomponent Synthesis and Biological Evaluation of a Piperazine-Based Dopamine Receptor Ligand Library. <i>ACS Medicinal Chemistry Letters</i> , 2015 , 6, 882-7	4.3	8
214	Improved radiosynthesis and preliminary in vivo evaluation of a (18)F-labeled glycopeptide-peptoid hybrid for PET imaging of neurotensin receptor 2. <i>Bioorganic and Medicinal Chemistry</i> , 2015 , 23, 4026-33	3.4	18
213	Molecular determinants of biased agonism at the dopamine D α receptor. <i>Journal of Medicinal Chemistry</i> , 2015 , 58, 2703-17	8.3	35
212	1,4-Disubstituted aromatic piperazines with high 5-HT $2A/D2$ selectivity: Quantitative structure-selectivity investigations, docking, synthesis and biological evaluation. <i>Bioorganic and Medicinal Chemistry</i> , 2015 , 23, 6195-209	3.4	13
211	The broad-spectrum anti-infective drug artesunate interferes with the canonical nuclear factor kappa B (NF- κ B) pathway by targeting RelA/p65. <i>Antiviral Research</i> , 2015 , 124, 101-9	10.8	37
210	Die Rhodopsin-Arrestin-Kristallstruktur und ihre Bedeutung für die Entwicklung funktionell selektiver GPCR-Wirkstoffe. <i>Angewandte Chemie</i> , 2015 , 127, 13362-13364	3.6	
209	Arrestin-Bound Rhodopsin: A Molecular Structure and its Impact on the Development of Biased GPCR Ligands. <i>Angewandte Chemie - International Edition</i> , 2015 , 54, 13166-8	16.4	2
208	Fluoro-substituted phenylazocarboxamides: Dopaminergic behavior and N-arylation properties for irreversible binding. <i>Bioorganic and Medicinal Chemistry</i> , 2015 , 23, 3938-47	3.4	3
207	Selective GPCR ligands. <i>Bioorganic and Medicinal Chemistry</i> , 2015 , 23, 3879	3.4	
206	Covalent molecular probes for class A G protein-coupled receptors: advances and applications. <i>ACS Chemical Biology</i> , 2015 , 10, 1376-86	4.9	45
205	GPCR crystal structures: Medicinal chemistry in the pocket. <i>Bioorganic and Medicinal Chemistry</i> , 2015 , 23, 3880-906	3.4	88
204	GPCRs als hochrelevante Targets für die Wirkstoffentwicklung. <i>BioSpektrum</i> , 2014 , 20, 15-18	0.1	1
203	Functionally selective dopamine D α D α receptor partial agonists. <i>Journal of Medicinal Chemistry</i> , 2014 , 57, 4861-75	8.3	60
202	Synthesis and evaluation of fluoro substituted pyridinylcarboxamides and their phenylazo analogues for potential dopamine D3 receptor PET imaging. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2014 , 24, 5399-403	2.9	10
201	Synthesis and binding profile of haloperidol-based bivalent ligands targeting dopamine D(2)-like receptors. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2014 , 24, 3753-6	2.9	10

200	Fast and efficient (18) F-labeling by [(18) F]fluorophenylazocarboxylic esters. <i>Chemistry - A European Journal</i> , 2014 , 20, 370-5	4.8	29
199	Active-state model of a dopamine D2 receptor-Gβγ complex stabilized by aripiprazole-type partial agonists. <i>PLoS ONE</i> , 2014 , 9, e100069	3.7	26
198	Covalent agonists for studying G protein-coupled receptor activation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014 , 111, 10744-8	11.5	69
197	H-CRRETAWAC-OH, a lead structure for the development of radiotracer targeting integrin α _v β ₃ ?. <i>BioMed Research International</i> , 2014 , 2014, 243185	3	11
196	Structure-based evolution of subtype-selective neurotensin receptor ligands. <i>ChemistryOpen</i> , 2014 , 3, 206-18	2.3	8
195	In vivo monitoring of the antiangiogenic effect of neurotensin receptor-mediated radiotherapy by small-animal positron emission tomography: a pilot study. <i>Pharmaceuticals</i> , 2014 , 7, 464-81	5.2	13
194	Biodistribution studies of two 18F-labeled pyridinylphenyl amides as subtype selective radioligands for the dopamine D3 receptor. <i>Nuclear Medicine and Biology</i> , 2014 , 41, 223-8	2.1	13
193	The pH probe CypHer488 is effectively quenched by FM dyes. <i>Journal of Fluorescence</i> , 2013 , 23, 487-94	2.4	5
192	Click chemistry based synthesis of dopamine D4 selective receptor ligands for the selection of potential PET tracers. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2013 , 23, 6079-82	2.9	11
191	Activation and allosteric modulation of a muscarinic acetylcholine receptor. <i>Nature</i> , 2013 , 504, 101-6	50.4	639
190	Synthesis and evaluation of a (18)F-labeled diarylpyrazole glycoconjugate for the imaging of NTS1-positive tumors. <i>Journal of Medicinal Chemistry</i> , 2013 , 56, 9361-5	8.3	31
189	Development of a metabolically stable neurotensin receptor 2 (NTS2) ligand. <i>ChemMedChem</i> , 2013 , 8, 75-81	3.7	28
188	Dopamine agonist-induced penile erection and yawning: a comparative study in outbred Roman high- and low-avoidance rats. <i>Pharmacology Biochemistry and Behavior</i> , 2013 , 109, 59-66	3.9	17
187	Discovery of dopamine D ₁ receptor antagonists with planar chirality. <i>Bioorganic and Medicinal Chemistry</i> , 2013 , 21, 1680-4	3.4	7
186	Class A G-protein-coupled receptor (GPCR) dimers and bivalent ligands. <i>Journal of Medicinal Chemistry</i> , 2013 , 56, 6542-59	8.3	87
185	Functionally selective dopamine D ₂ /D ₃ receptor agonists comprising an enyne moiety. <i>Journal of Medicinal Chemistry</i> , 2013 , 56, 5130-41	8.3	49
184	<i>Argyrea nervosa</i> (Burm. f.): receptor profiling of lysergic acid amide and other potential psychedelic LSD-like compounds by computational and binding assay approaches. <i>Journal of Ethnopharmacology</i> , 2013 , 148, 492-7	5	18
183	Efficient Synthesis of Heterocyclic Neurotensin Receptor Ligands by Microwave-Assisted Aminocarbonylation. <i>Synthesis</i> , 2013 , 45, 2474-2480	2.9	5

182	Pharmacological profile of 2-bromoterguride at human dopamine D2, porcine serotonin 5-hydroxytryptamine 2A, and α C-adrenergic receptors, and its antipsychotic-like effects in rats. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2013 , 347, 57-68	4.7	9
181	Muscarinic receptors as model targets and antitargets for structure-based ligand discovery. <i>Molecular Pharmacology</i> , 2013 , 84, 528-40	4.3	49
180	Syntheses, receptor bindings, in vitro and in vivo stabilities and biodistributions of DOTA-neurotensin(8-13) derivatives containing amino acid residues - a lesson about the importance of animal experiments. <i>Chemistry and Biodiversity</i> , 2013 , 10, 2101-21	2.5	19
179	Impact of the proline residue on ligand binding of neurotensin receptor 2 (NTS2)-selective peptide-peptoid hybrids. <i>ChemMedChem</i> , 2013 , 8, 772-8	3.7	14
178	Active-state models of ternary GPCR complexes: determinants of selective receptor-G-protein coupling. <i>PLoS ONE</i> , 2013 , 8, e67244	3.7	33
177	Bivalent molecular probes for dopamine D2-like receptors. <i>Bioorganic and Medicinal Chemistry</i> , 2012 , 20, 455-66	3.4	24
176	Selective agonists for dopamine/neurotensin receptor heterodimers. <i>ChemMedChem</i> , 2012 , 7, 509-14	3.7	11
175	Novel azulene derivatives for the treatment of erectile dysfunction. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2012 , 22, 7151-4	2.9	24
174	Biotransformation of 21-O-acetyl-deoxycorticosterone by cell suspension cultures of <i>Digitalis lanata</i> (strain W.1.4). <i>Steroids</i> , 2012 , 77, 1373-80	2.8	15
173	A new D μ dopamine receptor agonist allosterically modulates A(2A) adenosine receptor signalling by interacting with the A(2A)/D μ receptor heteromer. <i>Cellular Signalling</i> , 2012 , 24, 951-60	4.9	14
172	Conformationally Restricted Peptide Mimetics by Ring-Closing Olefin Metathesis. <i>Synthesis</i> , 2012 , 44, 2682-2694	2.9	10
171	A highly efficient type I turn mimetic simulating an Asx-Pro-turn-like structure. <i>Organic Letters</i> , 2011 , 13, 3502-5	6.2	10
170	Evaluation of 18F-labeled benzodioxine piperazine-based dopamine D4 receptor ligands: lipophilicity as a determinant of nonspecific binding. <i>Journal of Medicinal Chemistry</i> , 2011 , 54, 8343-52	8.3	26
169	Development of a bivalent dopamine D μ receptor agonist. <i>Journal of Medicinal Chemistry</i> , 2011 , 54, 7911-9.3	9.3	29
168	Discovery of highly potent and neurotensin receptor 2 selective neurotensin mimetics. <i>Journal of Medicinal Chemistry</i> , 2011 , 54, 2915-23	8.3	49
167	Radical arylation of tyrosine and its application in the synthesis of a highly selective neurotensin receptor 2 ligand. <i>Organic and Biomolecular Chemistry</i> , 2011 , 9, 3746-52	3.9	25
166	Dopamine D2-like receptor agonists induce penile erection in male rats: differential role of D2, D3 and D4 receptors in the paraventricular nucleus of the hypothalamus. <i>Behavioural Brain Research</i> , 2011 , 225, 169-76	3.4	29
165	Molecular dynamics simulations of the effect of the G-protein and diffusible ligands on the α -adrenergic receptor. <i>Journal of Molecular Biology</i> , 2011 , 414, 611-23	6.5	29

164	Recent advances in the search for D3- and D4-selective drugs: probes, models and candidates. <i>Trends in Pharmacological Sciences</i> , 2011 , 32, 148-57	13.2	90
163	Bivalent dopamine D2 receptor ligands: synthesis and binding properties. <i>Journal of Medicinal Chemistry</i> , 2011 , 54, 4896-903	8.3	61
162	Quinpramine ameliorates rat experimental autoimmune neuritis and redistributes MHC class II molecules. <i>PLoS ONE</i> , 2011 , 6, e21223	3.7	8
161	Structure and function of an irreversible agonist- α_2 adrenoceptor complex. <i>Nature</i> , 2011 , 469, 236-40	50.4	664
160	Highly potent 5-aminotetrahydropyrazolopyridines: enantioselective dopamine D3 receptor binding, functional selectivity, and analysis of receptor-ligand interactions. <i>Journal of Medicinal Chemistry</i> , 2011 , 54, 2477-91	8.3	50
159	On the terminal homologation of physiologically active peptides as a means of increasing stability in human serum--neurotensin, opiorphin, B27-KK10 epitope, NPY. <i>Chemistry and Biodiversity</i> , 2011 , 8, 711-39	2.5	23
158	Cross-receptor interactions between dopamine D2L and neurotensin NTS1 receptors modulate binding affinities of dopaminergics. <i>ACS Chemical Neuroscience</i> , 2011 , 2, 308-16	5.7	36
157	Aromatic ring functionalization of benzolactam derivatives: new potent dopamine D3 receptor ligands. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2011 , 21, 2670-4	2.9	17
156	The bulky N6 substituent of cabergoline is responsible for agonism of this drug at 5-hydroxytryptamine 5-HT2A and 5-HT2B receptors and thus is a determinant of valvular heart disease. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2011 , 338, 381-91	4.7	7
155	Histidine 6.55 is a major determinant of ligand-biased signaling in dopamine D2L receptor. <i>Molecular Pharmacology</i> , 2011 , 79, 575-85	4.3	67
154	Tetracycline sensing using novel doxycycline derivatives immobilized on different surface plasmon resonance biosensor surfaces. <i>Bioanalysis</i> , 2010 , 2, 217-27	2.1	4
153	Functional characterization of a partial loss-of-function mutation of the epithelial sodium channel (ENaC) associated with atypical cystic fibrosis. <i>Cellular Physiology and Biochemistry</i> , 2010 , 25, 145-58	3.9	25
152	The therapeutically anti-prion active antibody-fragment scFv-W226: paramagnetic relaxation-enhanced NMR spectroscopy aided structure elucidation of the paratope-epitope interface. <i>Journal of Biomolecular Structure and Dynamics</i> , 2010 , 28, 13-22	3.6	12
151	Novel pyridylmethylamines as highly selective 5-HT(1A) superagonists. <i>Journal of Medicinal Chemistry</i> , 2010 , 53, 7167-79	8.3	20
150	Bioisosteric replacement leading to biologically active [2.2]paracyclophanes with altered binding profiles for aminergic G-protein-coupled receptors. <i>Journal of Medicinal Chemistry</i> , 2010 , 53, 7219-28	8.3	12
149	Synthesis of a (68)ga-labeled peptoid-Peptide hybrid for imaging of neurotensin receptor expression in vivo. <i>ACS Medicinal Chemistry Letters</i> , 2010 , 1, 224-8	4.3	20
148	Engineering a GPCR-ligand pair that simulates the activation of D(2L) by Dopamine. <i>ACS Chemical Neuroscience</i> , 2010 , 1, 25-35	5.7	19
147	Anhydrotetracycline-peptide conjugates as representatives for ligand-based transactivating systems. <i>Bioorganic and Medicinal Chemistry</i> , 2010 , 18, 6127-33	3.4	1

- 146 A series of ¹⁸F-labelled pyridinylphenyl amides as subtype-selective radioligands for the dopamine D3 receptor. *ChemMedChem*, **2010**, 5, 941-8 3.7 13
- 145 Click-chemistry-derived tetracycline-amino acid conjugates exhibiting exceptional potency and exclusive recognition of the reverse tet repressor. *ChemBioChem*, **2010**, 11, 703-12 3.8 1
- 144 An efficient and practical EPC synthesis of α -amino acids from L-asparagine. *Archiv Der Pharmazie*, **2010**, 324, 551-557 4.3 16
- 143 Markierung und Glycosylierung von Peptiden mithilfe der Klick-Chemie: ein allgemeiner Ansatz zur Synthese von ¹⁸F-Glycopeptiden, leistungsstarken Tracern für die Positronenemissionstomographie. *Angewandte Chemie*, **2010**, 122, 988-992 3.6 18
- 142 Labeling and glycosylation of peptides using click chemistry: a general approach to (¹⁸F)-glycopeptides as effective imaging probes for positron emission tomography. *Angewandte Chemie - International Edition*, **2010**, 49, 976-9 16.4 103
- 141 Synthesis, biological evaluation and radiolabelling by ¹⁸F-fluoroarylation of a dopamine D3-selective ligand as prospective imaging probe for PET. *Bioorganic and Medicinal Chemistry Letters*, **2010**, 20, 6933-7 2.9 13
- 140 Dopamine Receptor Subtype-Selective Drugs: D2-Like Receptors **2010**, 101-135
- 139 Diarylpropane-1,3-dione derivatives as TetR-inducing tetracycline mimetics: Synthesis and biological investigations. *ChemBioChem*, **2009**, 10, 2924-33 3.8 7
- 138 The azulene framework as a novel arene bioisostere: design of potent dopamine D4 receptor ligands inducing penile erection. *ChemMedChem*, **2009**, 4, 325-8 3.7 23
- 137 Design, synthesis and dopamine D4 receptor binding activities of new N-heteroaromatic 5/6-ring Mannich bases. *Bioorganic and Medicinal Chemistry*, **2009**, 17, 4448-58 3.4 14
- 136 Structural origins for selectivity and specificity in an engineered bacterial repressor-inducer pair. *FEBS Journal*, **2009**, 276, 5610-21 5.7 4
- 135 Novel triazolo-peptides: chiro-specific synthesis and conformational studies of proline derived analogs. *Tetrahedron*, **2009**, 65, 6156-6168 2.4 11
- 134 Parallel synthesis of potent dopaminergic N-phenyltriazole carboxamides applying a novel click chemistry based phenol linker. *Bioorganic and Medicinal Chemistry*, **2009**, 17, 5482-7 3.4 10
- 133 Quinpramine is a novel compound effective in ameliorating brain autoimmune disease. *Experimental Neurology*, **2009**, 215, 397-400 5.7 8
- 132 Dopamine D2, D3, and D4 selective phenylpiperazines as molecular probes to explore the origins of subtype specific receptor binding. *Journal of Medicinal Chemistry*, **2009**, 52, 4923-35 8.3 75
- 131 1,1RDisubstituted ferrocenes as molecular hinges in mono- and bivalent dopamine receptor ligands. *Journal of Medicinal Chemistry*, **2009**, 52, 6860-70 8.3 69
- 130 Proline derived spirobarbiturates as highly effective beta-turn mimetics incorporating polar and functionalizable constraint elements. *Journal of Organic Chemistry*, **2008**, 73, 3608-11 4.2 51
- 129 Novel D3 selective dopaminergics incorporating enyne units as nonaromatic catechol bioisosteres: synthesis, bioactivity, and mutagenesis studies. *Journal of Medicinal Chemistry*, **2008**, 51, 6829-38 8.3 30

128	A simple chemical method for synthesizing malonyl hemiesters of 21-hydroxypregnanes, potential intermediates in cardenolide biosynthesis. <i>Steroids</i> , 2008 , 73, 458-65	2.8	6
127	Characterization of the molecular fragment that is responsible for agonism of pergolide at serotonin 5-Hydroxytryptamine _{2B} and 5-Hydroxytryptamine _{2A} receptors. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2008 , 324, 1136-45	4.7	13
126	¹⁸ F-Labeled FAUC 346 and BP 897 derivatives as subtype-selective potential PET radioligands for the dopamine D ₃ receptor. <i>ChemMedChem</i> , 2008 , 3, 788-93	3.7	23
125	Novel insights into GPCR-peptide interactions: mutations in extracellular loop 1, ligand backbone methylations and molecular modeling of neurotensin receptor 1. <i>Bioorganic and Medicinal Chemistry</i> , 2008 , 16, 9359-68	3.4	33
124	Discovery of a dopamine D ₄ selective PET ligand candidate taking advantage of a click chemistry based REM linker. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2008 , 18, 983-8	2.9	20
123	Peptide backbone modifications on the C-terminal hexapeptide of neurotensin. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2008 , 18, 2013-8	2.9	30
122	Synthesis, radiofluorination, and in vitro evaluation of pyrazolo[1,5-a]pyridine-based dopamine D ₄ receptor ligands: discovery of an inverse agonist radioligand for PET. <i>Journal of Medicinal Chemistry</i> , 2008 , 51, 1800-10	8.3	37
121	Structure-selectivity investigations of D ₂ -like receptor ligands by CoMFA and CoMSIA guiding the discovery of D ₃ selective PET radioligands. <i>Journal of Medicinal Chemistry</i> , 2007 , 50, 489-500	8.3	46
120	Molecular building kit of fused-proline-derived peptide mimetics allowing specific adjustment of the dihedral Psi angle. <i>Journal of Organic Chemistry</i> , 2007 , 72, 9102-13	4.2	29
119	3,4,6-Tri-O-acetyl-2-deoxy-2-[¹⁸ F]fluoroglucopyranosyl phenylthiosulfonate: a thiol-reactive agent for the chemoselective ¹⁸ F-glycosylation of peptides. <i>Bioconjugate Chemistry</i> , 2007 , 18, 254-62	6.3	58
118	In vitro affinities of various halogenated benzamide derivatives as potential radioligands for non-invasive quantification of D(2)-like dopamine receptors. <i>Bioorganic and Medicinal Chemistry</i> , 2007 , 15, 6819-29	3.4	27
117	Click chemistry based solid phase supported synthesis of dopaminergic phenylacetylenes. <i>Bioorganic and Medicinal Chemistry</i> , 2007 , 15, 7248-57	3.4	18
116	Optimization of Click Chemistry Using Azide and Alkyne Scavenger Resins. <i>QSAR and Combinatorial Science</i> , 2007 , 26, 1145-1150		17
115	Dopamine D ₃ receptor ligands: recent advances in the control of subtype selectivity and intrinsic activity. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2007 , 1768, 871-87	3.8	35
114	Synthesis and biological investigations of dopaminergic partial agonists preferentially recognizing the D ₄ receptor subtype. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2006 , 16, 2955-9	2.9	36
113	The structural evolution of dopamine D ₃ receptor ligands: structure-activity relationships and selected neuropharmacological aspects 2006 , 112, 281-333		74
112	A highly practical RCM approach towards a molecular building kit of spirocyclic reverse turn mimics. <i>Chemistry - A European Journal</i> , 2006 , 12, 6315-22	4.8	52
111	Subtype selective tetracycline agonists and their application for a two-stage regulatory system. <i>ChemBioChem</i> , 2006 , 7, 1320-4	3.8	21

- 110 A gene regulation system with four distinct expression levels. *Journal of Gene Medicine*, **2006**, 8, 1037-47^{3,5} 31
- 109 Syntheses and radiofluorination of two derivatives of 5-cyano-indole as selective ligands for the dopamine subtype-4 receptor. *Journal of Labelled Compounds and Radiopharmaceuticals*, **2006**, 49, 55-70^{1,9} 11
- 108 2-[(4-phenylpiperazin-1-yl)methyl]imidazo(di)azines as selective D4-ligands. Induction of penile erection by 2-[4-(2-methoxyphenyl)piperazin-1-ylmethyl]imidazo[1,2-a]pyridine (PIP3EA), a potent and selective D4 partial agonist. *Journal of Medicinal Chemistry*, **2006**, 49, 3938-47 8.3 47
- 107 Chiroselective synthesis of spirocyclic beta-lactams and their characterization as potent type II beta-turn inducing peptide mimetics. *Journal of Organic Chemistry*, **2006**, 71, 97-102 4.2 73
- 106 SCRF-DFT and NMR comparison of tetracycline and 5a,6-anhydrotetracycline in solution. *Journal of Physical Chemistry B*, **2006**, 110, 24766-74 3.4 12
- 105 A chimeric ligand approach leading to potent antiprion active acridine derivatives: design, synthesis, and biological investigations. *Journal of Medicinal Chemistry*, **2006**, 49, 6591-5 8.3 57
- 104 Fancy bioisosteres: novel paracyclophane derivatives as super-affinity dopamine D3 receptor antagonists. *Journal of Medicinal Chemistry*, **2006**, 49, 3628-35 8.3 25
- 103 Click chemistry on solid phase: parallel synthesis of N-benzyltriazole carboxamides as super-potent G-protein coupled receptor ligands. *ACS Combinatorial Science*, **2006**, 8, 252-61 49
- 102 Dopamine D2 and D3 receptors in human putamen, caudate nucleus, and globus pallidus. *Synapse*, **2006**, 60, 205-11 2.4 61
- 101 PIP3EA and PD-168077, two selective dopamine D4 receptor agonists, induce penile erection in male rats: site and mechanism of action in the brain. *European Journal of Neuroscience*, **2006**, 24, 2021-30^{3,5} 38
- 100 Tricyclic antidepressants, quinacrine and a novel, synthetic chimera thereof clear prions by destabilizing detergent-resistant membrane compartments. *Journal of Neurochemistry*, **2006**, 98, 748-59⁶ 67
- 99 Tricyclic antidepressants, quinacrine and a novel, synthetic chimera thereof clear prions by destabilizing detergent-resistant membrane compartments. *Journal of Neurochemistry*, **2006**, 98, 1696-1696⁶ 2
- 98 CoMFA and CoMSIA investigations of dopamine D3 receptor ligands leading to the prediction, synthesis, and evaluation of rigidized FAUC 365 analogues. *Bioorganic and Medicinal Chemistry*, **2006**, 14, 5898-912 3.4 19
- 97 Bicyclic melatonin receptor agonists containing a ring-junction nitrogen: Synthesis, biological evaluation, and molecular modeling of the putative bioactive conformation. *Bioorganic and Medicinal Chemistry*, **2006**, 14, 1949-58 3.4 27
- 96 A consecutive Diels-Alder approach toward a Tet repressor directed combinatorial library. *Tetrahedron*, **2006**, 62, 6899-6908 2.4 15
- 95 Triazolo-peptides: chiroselective synthesis and cis/trans prolyl ratios of structural isomers. *Tetrahedron*, **2006**, 62, 8919-8927 2.4 41
- 94 CoMFA and CoMSIA investigations revealing novel insights into the binding modes of dopamine D3 receptor agonists. *Journal of Medicinal Chemistry*, **2005**, 48, 2493-508 8.3 28
- 93 Modeling the similarity and divergence of dopamine D2-like receptors and identification of validated ligand-receptor complexes. *Journal of Medicinal Chemistry*, **2005**, 48, 694-709 8.3 41

92	Fancy bioisosteres: metallocene-derived G-protein-coupled receptor ligands with subnanomolar binding affinity and novel selectivity profiles. <i>Journal of Medicinal Chemistry</i> , 2005 , 48, 3696-9	8.3	62
91	Pharmacophore-guided drug discovery investigations leading to bioactive 5-aminotetrahydropyrazolopyridines. Implications for the binding mode of heterocyclic dopamine D3 receptor agonists. <i>Journal of Medicinal Chemistry</i> , 2005 , 48, 5771-9	8.3	38
90	Fancy bioisosteres: synthesis and dopaminergic properties of the endiynes FAUC 88 as a novel non-aromatic D3 agonist. <i>Bioorganic and Medicinal Chemistry</i> , 2005 , 13, 185-91	3.4	24
89	Fancy bioisosteres: synthesis, SAR, and pharmacological investigations of novel nonaromatic dopamine D3 receptor ligands. <i>Bioorganic and Medicinal Chemistry</i> , 2005 , 13, 4434-42	3.4	22
88	Synthesis and evaluation of 18F-labeled dopamine D3 receptor ligands as potential PET imaging agents. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2005 , 15, 4819-23	2.9	33
87	Parallel synthesis and biological screening of dopamine receptor ligands taking advantage of a click chemistry based BAL linker. <i>ACS Combinatorial Science</i> , 2005 , 7, 309-16		57
86	Chiroselective and subtype selective dopamine receptor binding of heterocyclic methoxynaphthamide analogs. <i>Archiv Der Pharmazie</i> , 2005 , 338, 276-80	4.3	4
85	Certain 1,4-disubstituted aromatic piperidines and piperazines with extreme selectivity for the dopamine D4 receptor interact with a common receptor microdomain. <i>Molecular Pharmacology</i> , 2004 , 66, 1491-9	4.3	36
84	Synthesis and radioiodination of selective ligands for the dopamine D3 receptor subtype. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2004 , 14, 3963-6	2.9	35
83	FAUC 213, a highly selective dopamine D4 receptor full antagonist, exhibits atypical antipsychotic properties in behavioural and neurochemical models of schizophrenia. <i>Psychopharmacology</i> , 2004 , 175, 7-17	4.7	31
82	Analogues of FAUC 73 revealing new insights into the structural requirements of nonaromatic dopamine D3 receptor agonists. <i>Bioorganic and Medicinal Chemistry</i> , 2004 , 12, 113-7	3.4	16
81	Clozapine derived 2,3-dihydro-1H-1,4- and 1,5-benzodiazepines with D4 receptor selectivity: synthesis and biological testing. <i>Bioorganic and Medicinal Chemistry</i> , 2004 , 12, 2625-37	3.4	22
80	Practical ex-chiral-pool methodology for the synthesis of dopaminergic tetrahydroindoles. <i>Tetrahedron</i> , 2004 , 60, 1197-1204	2.4	19
79	Click chemistry on solid support: synthesis of a new REM resin and application for the preparation of tertiary amines. <i>Tetrahedron</i> , 2004 , 60, 8699-8702	2.4	47
78	Structure-based design of Tet repressor to optimize a new inducer specificity. <i>Biochemistry</i> , 2004 , 43, 9512-8	3.2	41
77	Evaluation of lactam-bridged neurotensin analogues adjusting psi(Pro10) close to the experimentally derived bioactive conformation of NT(8-13). <i>Journal of Medicinal Chemistry</i> , 2004 , 47, 5587-90	8.3	39
76	Attenuation of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) neurotoxicity by the novel selective dopamine D3-receptor partial agonist FAUC 329 predominantly in the nucleus accumbens of mice. <i>Biochemical Pharmacology</i> , 2003 , 66, 1025-32	6	20
75	Ex-chiral pool synthesis and receptor binding studies of 4-substituted prolinol derivatives. <i>Tetrahedron: Asymmetry</i> , 2003 , 14, 3141-3152		17

74	Enantiospecific synthesis and receptor binding of novel dopamine receptor ligands employing natural 4-hydroxyproline as a practical and flexible building block. <i>Tetrahedron: Asymmetry</i> , 2003 , 14, 3153-3172		23
73	Synthesis of dihydrooxazole analogues derived from linezolid. <i>Tetrahedron</i> , 2003 , 59, 3403-3407	2.4	6
72	Cyclic amidines as benzamide bioisosteres: EPC synthesis and SAR studies leading to the selective dopamine D4 receptor agonist FAUC 312. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2003 , 13, 851-4	2.9	18
71	Stereocontrolled dopamine receptor binding and subtype selectivity of clebopride analogues synthesized from aspartic acid. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2003 , 13, 3293-6	2.9	11
70	A general approach to dehydro-Freidinger lactams: ex-chiral pool synthesis and spectroscopic evaluation as potential reverse turn inducers. <i>Journal of Organic Chemistry</i> , 2003 , 68, 62-9	4.2	49
69	Click linker: efficient and high-yielding synthesis of a new family of SPOS resins by 1,3-dipolar cycloaddition. <i>Organic Letters</i> , 2003 , 5, 1753-5	6.2	156
68	Analogs of the dopamine D4 receptor ligand FAUC 113 with planar- and central-chirality. <i>Tetrahedron: Asymmetry</i> , 2002 , 13, 2303-2310		26
67	Di- and trisubstituted pyrazolo[1,5-a]pyridine derivatives: synthesis, dopamine receptor binding and ligand efficacy. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2002 , 12, 633-6	2.9	29
66	Fused azaindole derivatives: molecular design, synthesis and in vitro pharmacology leading to the preferential dopamine D3 receptor agonist FAUC 725. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2002 , 12, 2377-80	2.9	44
65	2,4-Disubstituted pyrroles: synthesis, traceless linking and pharmacological investigations leading to the dopamine D4 receptor partial agonist FAUC 356. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2002 , 12, 1937-40	2.9	24
64	Phenylpiperazinylmethylindolecarboxylates and derivatives as selective D(4)-ligands. <i>Bioorganic and Medicinal Chemistry</i> , 2002 , 10, 1671-9	3.4	18
63	Interactive SAR studies: rational discovery of super-potent and highly selective dopamine D3 receptor antagonists and partial agonists. <i>Journal of Medicinal Chemistry</i> , 2002 , 45, 4594-7	8.3	166
62	Traceless Linking of Pyrroles: General Methodology and Solid Phase Supported Functionalizations. <i>Synthesis</i> , 2002 , 2002, 0274	2.9	6
61	Short and Efficient Synthesis of Homo-Freidinger Lactams: An Olefin Metathesis Approach Towards Conformationally Restricted β -Amino Acid Analogues. <i>Synlett</i> , 2002 , 2002, 1014-1016	2.2	15
60	Planar chiral indoles: synthesis and biological effects of the enantiomers. <i>Tetrahedron: Asymmetry</i> , 2001 , 12, 3205-3208		16
59	Benzamide bioisosteres incorporating dihydroheteroazole substructures: EPC synthesis and SAR leading to a selective dopamine D4 receptor partial agonist (FAUC 179). <i>Bioorganic and Medicinal Chemistry Letters</i> , 2001 , 11, 2533-6	2.9	26
58	Dopaminergic 7-aminotetrahydroindolizines: ex-chiral pool synthesis and preferential D3 receptor binding. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2001 , 11, 2863-6	2.9	20
57	Indoloparacyclophanes: Synthesis and Dopamine Receptor Binding of a Novel Arylbioisostere. <i>Angewandte Chemie</i> , 2001 , 113, 1323-1325	3.6	6

56	Rational Molecular Design and EPC Synthesis of a Type VI β -Turn Inducing Peptide Mimetic. <i>Angewandte Chemie</i> , 2001 , 113, 3465-3468	3.6	8
55	Indoloparacyclophanes: Synthesis and Dopamine Receptor Binding of a Novel Arylbioisostere This work was supported by the Fonds der Chemischen Industrie.. <i>Angewandte Chemie - International Edition</i> , 2001 , 40, 1283-1285	16.4	34
54	Rational Molecular Design and EPC Synthesis of a Type VI β -Turn Inducing Peptide Mimetic This work was supported by the BMBF and the Fonds der Chemischen Industrie. EPC=enantiomerically pure compound.. <i>Angewandte Chemie - International Edition</i> , 2001 , 40, 3361-3364	16.4	52
53	Rationally based efficacy tuning of selective dopamine d4 receptor ligands leading to the complete antagonist 2-[4-(4-chlorophenyl)piperazin-1-ylmethyl]pyrazolo[1,5-a]pyridine (FAUC 213). <i>Journal of Medicinal Chemistry</i> , 2001 , 44, 2691-4	8.3	88
52	Diethoxymethyl Protected Pyrroles: Synthesis and Regioselective Transformations. <i>Synthesis</i> , 2001 , 2001, 2281-2288	2.9	16
51	Comparative molecular field analysis of dopamine D4 receptor antagonists including 3-[4-(4-chlorophenyl)piperazin-1-ylmethyl]pyrazolo[1,5-a]pyridine (FAUC 113), 3-[4-(4-chlorophenyl)piperazin-1-ylmethyl]-1H-pyrrolo-[2,3-b]pyridine (L-745,870), and clozapine. <i>Journal of Medicinal Chemistry</i> , 2001 , 44, 1151-7	8.3	41
50	Azepino- and diazepinoindoles: synthesis and dopamine receptor binding profiles. <i>Archiv Der Pharmazie</i> , 2000 , 333, 287-92	4.3	16
49	Phenyloxazoles and phenylthiazoles as benzamide bioisosteres: synthesis and dopamine receptor binding profiles. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2000 , 10, 2041-4	2.9	32
48	Synthesis of Enantiopure 8-Aminomethylindolizines from Glutamine by Stereoelectronically Controlled Cationic Cyclization. <i>Heterocycles</i> , 2000 , 53, 1371	0.8	9
47	Effective and Variable Functionalization of Pyrazolo[1,5-a]pyridines Involving Palladium-Catalyzed Coupling Reactions. <i>Synthesis</i> , 2000 , 2000, 1727-1732	2.9	26
46	Regiocontrolled Annellation Reactions Starting from Diethoxymethyl Protected Indole. <i>Synthesis</i> , 2000 , 2000, 1081-1083	2.9	8
45	Conjugated enynes as nonaromatic catechol bioisosteres: synthesis, binding experiments, and computational studies of novel dopamine receptor agonists recognizing preferentially the D(3) subtype. <i>Journal of Medicinal Chemistry</i> , 2000 , 43, 756-62	8.3	157
44	Cyanoindole derivatives as highly selective dopamine D(4) receptor partial agonists: solid-phase synthesis, binding assays, and functional experiments. <i>Journal of Medicinal Chemistry</i> , 2000 , 43, 4563-9	8.3	57
43	Treatment of N,N-dibenzylamino alcohols with sulfonyl chloride leads to rearranged beta-chloro amines, precursors to beta-amino acids, and not to tetrahydroisoquinolines. <i>Organic Letters</i> , 2000 , 2, 647-9	6.2	42
42	Enantiopure 4- and 5-aminopiperidin-2-ones: regiocontrolled synthesis and conformational characterization as bioactive beta-turn mimetics. <i>Journal of Organic Chemistry</i> , 2000 , 65, 7406-16	4.2	38
41	Azaindole derivatives with high affinity for the dopamine D4 receptor: synthesis, ligand binding studies and comparison of molecular electrostatic potential maps. <i>Bioorganic and Medicinal Chemistry Letters</i> , 1999 , 9, 97-102	2.9	82
40	Enantio- and diastereocontrolled dopamine D1, D2, D3 and D4 receptor binding of N-(3-pyrrolidinylmethyl)benzamides synthesized from aspartic acid. <i>Bioorganic and Medicinal Chemistry Letters</i> , 1999 , 9, 841-6	2.9	16
39	2,2-Dicyanovinyl as a nonaromatic aryl bioisostere: synthesis, binding experiments and SAR studies of highly selective dopamine D4 receptor ligands. <i>Bioorganic and Medicinal Chemistry Letters</i> , 1999 , 9, 1969-72	2.9	19

38	Piperidinyppyroles: design, synthesis and binding properties of novel and selective dopamine D4 receptor ligands. <i>Bioorganic and Medicinal Chemistry Letters</i> , 1999 , 9, 3143-6	2.9	33
37	Ex-Chiral Pool Synthesis of Aminooxazepinones as Conformationally Restricted β -Amino Acid Analogs. <i>Heterocycles</i> , 1999 , 51, 365	0.8	7
36	Chemo- and Regioselective Syntheses of Enantiopure Aminopyrrolidinones as Building Blocks for Constrained Peptidomimetics. <i>Heterocycles</i> , 1999 , 51, 1389	0.8	11
35	Beta-analogs of PLG (L-prolyl-L-leucyl-glycinamide): ex-chiral pool syntheses and dopamine D2 receptor modulating effects. <i>Bioorganic and Medicinal Chemistry Letters</i> , 1998 , 8, 2885-90	2.9	19
34	Homo-Freidinger Lactams: Stereoselective Synthesis of 4-Aminopiperidin-2-one Derivatives from Aspartic Acid. <i>Synlett</i> , 1998 , 1998, 885-887	2.2	11
33	A Practical Ex-Chiral-Pool Synthesis of β -Proline and Homo- β -Proline. <i>Synthesis</i> , 1998 , 1998, 1491-1496	2.9	31
32	Diethoxymethyl Protected Indoles: Synthesis and Regioselective Transformations. <i>Synthesis</i> , 1996 , 1996, 1196-1198	2.9	15
31	Synthesis and stereochemistry of oxazolo[4?,5?:1,2]benz[5,4,3-c, d]-indole derivatives as intermediates on the way to a selective dopamine autoreceptor agonist. <i>Journal of Heterocyclic Chemistry</i> , 1996 , 33, 481-483	1.9	1
30	2-Dibenzylaminobutane-1,4-diol: A versatile intermediate for a chiroselective β -amino acid synthesis. <i>Tetrahedron Letters</i> , 1995 , 36, 381-382	2	16
29	2-Dibenzylaminobutane-1,4-diol: An Effective Chiral Building Block for the Synthesis of 1,2-Amino Alcohols. <i>Synthesis</i> , 1995 , 1995, 83-86	2.9	7
28	Diethoxymethyl: A Useful Nitrogen-Protecting Group for Lactams and Amides. <i>Synthesis</i> , 1995 , 1995, 168-170	2.9	9
27	Regioselective Transformation of Malic Acid: A Practical Method for the Construction of Enantiomerically Pure Indolizidines. <i>Journal of Organic Chemistry</i> , 1995 , 60, 3910-3915	4.2	13
26	Synthesis of optically active β -disubstituted β -amino nitriles and β -amino acids starting from asparagine. <i>Liebigs Annalen</i> , 1995 , 1995, 1987-1992		15
25	Synthesis of (S)-3,4-diaminobutanenitriles as precursors for 3-amino-GABA derivatives. <i>Archiv Der Pharmazie</i> , 1995 , 328, 265-8	4.3	4
24	Synthesis and dopamine receptor binding of 3-phenylazepino[5,4,3-c,d]indole derivatives. <i>Archiv Der Pharmazie</i> , 1995 , 328, 329-32	4.3	8
23	Synthesis, pharmacological investigation and computational studies on a tricyclic ergoline analog with selective dopamine autoreceptor activity. <i>Archiv Der Pharmazie</i> , 1995 , 328, 609-14	4.3	7
22	Synthesis and dopamine receptor binding studies of homochiral 8-aminopyrido[1,2-a]indoles. <i>Archiv Der Pharmazie</i> , 1995 , 328, 626-8	4.3	3
21	Diastereoselective Synthesis of the C ₂ -Symmetric 2,3-Diaminotetralin via Electrophilic Amination. <i>Synthesis</i> , 1994 , 1994, 1026-1028	2.9	7

20	Enantiomerically Pure Amino Alcohols and Diamino Alcohols from L-Aspartic Acid. Application to the Synthesis of Epi- and Diepislafamine. <i>Journal of Organic Chemistry</i> , 1994 , 59, 6766-6776	4.2	46
19	Novel HO-DPAT (hydroxy-2-dipropylaminotetralin) isomers: Stereoselective synthesis and receptor binding studies. <i>Bioorganic and Medicinal Chemistry Letters</i> , 1994 , 4, 2589-2590	2.9	2
18	Asymmetric synthesis of β -aminotetralins by electrophilic amination. <i>Tetrahedron</i> , 1994 , 50, 10909-10922	2.4	10
17	Practical EPC synthesis of 1,2- and 1,3-amino alcohols. <i>Tetrahedron Letters</i> , 1993 , 34, 4325-4326	2	15
16	Dopamine autoreceptor agonists: computational studies, synthesis and biological investigations. <i>Bioorganic and Medicinal Chemistry Letters</i> , 1993 , 3, 1477-1483	2.9	16
15	Enantiomerically pure aminoindolizines: bicyclic ergoline analogues with dopamine autoreceptor activity. <i>Archiv Der Pharmazie</i> , 1992 , 325, 57-60	4.3	19
14	Tricyclic azaergoline analogues: synthesis, structural modifications, and pharmacological studies. <i>Archiv Der Pharmazie</i> , 1992 , 325, 649-55	4.3	6
13	Synthesis of Amines and Amino Alcohols by Electrophilic Amination and Highly Stereoselective Reduction. <i>Liebigs Annalen Der Chemie</i> , 1992 , 1992, 273-278		16
12	Synthesis of Chiral Indolizines as Bicyclic Ergoline Analogues. <i>Heterocycles</i> , 1991 , 32, 1499	0.8	7
11	Stereoelectronically controlled cyclization reactions on the way to peri-fused tetrahydropyrazolo[1,5-a]pyridines as aza analogs of ergoline partial structures. <i>Liebigs Annalen Der Chemie</i> , 1991 , 1991, 921-927		3
10	Efficient EPC synthesis of β -aminobutanoic acid. <i>Liebigs Annalen Der Chemie</i> , 1991 , 1991, 501-502		5
9	Efficient methodology for the preparation of β -aminotetralin derivatives via electrophilic amination. <i>Tetrahedron Letters</i> , 1991 , 32, 5927-5930	2	11
8	General synthesis of enantiomerically pure β -amino acids. <i>Tetrahedron Letters</i> , 1990 , 31, 5717-5720	2	40
7	Synthesis of Tricyclic Azaergoline Analogues Synthese tricyclischer Azaergolin-Analoga. <i>Archiv Der Pharmazie</i> , 1990 , 323, 991-993	4.3	4
6	An efficient and practical total synthesis of (+)-vincamine from L-aspartic acid. <i>Journal of Organic Chemistry</i> , 1990 , 55, 3068-3074	4.2	88
5	New and Efficient Synthesis of 5,6,7,8-Tetrahydroindolizidines. Application to the Synthesis of Pharmacologically Relevant Chiral Aminoderivatives from L-Asparagine. <i>Heterocycles</i> , 1990 , 31, 9	0.8	22
4	Pyrane, CXXIII. Regio- und diastereoselektive Synthesen von Hexahydro-4a,9-propanoxanthenonen. <i>Liebigs Annalen Der Chemie</i> , 1988 , 1988, 125-132		1
3	Oxabenzomorphone: Synthese ZNS-wirksamer Hexahydro-2,7-methano-1,5-benzoxazonine. <i>Archiv Der Pharmazie</i> , 1988 , 321, 321-324	4.3	2

- | | | | |
|---|---|-----|---|
| 2 | Synthese von ZNS-Wirkstoffen: Pyran-, thiopyran- und piperidinanellierte Oxa-carbabenzomorphane. <i>Archiv Der Pharmazie</i> , 1988 , 321, 397-401 | 4-3 | 3 |
| 1 | Synthese anellierter Oxabenzomorphane. <i>Archiv Der Pharmazie</i> , 1986 , 319, 431-434 | 4-3 | 6 |