

# Hitoshi Sugiyama

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4372115/publications.pdf>

Version: 2024-02-01

132  
papers

4,017  
citations

126708

33  
h-index

143772

57  
g-index

136  
all docs

136  
docs citations

136  
times ranked

4379  
citing authors

#	ARTICLE	IF	CITATIONS
1	Prevalence of Chronic Kidney Disease and Variation of Its Risk Factors by the Regions in Okayama Prefecture. <i>Journal of Personalized Medicine</i> , 2022, 12, 97.	1.1	6
2	Plasma Globotriaosylsphingosine and $\beta$ -Galactosidase A Activity as a Combined Screening Biomarker for Fabry Disease in a Large Japanese Cohort. <i>Current Issues in Molecular Biology</i> , 2021, 43, 389-404.	1.0	2
3	Blood concentrations of tacrolimus upon conversion from rabeprazole to vonoprazan in renal transplant recipients: Correlation with cytochrome P450 gene polymorphisms. <i>Drug Metabolism and Pharmacokinetics</i> , 2021, 40, 100407.	1.1	3
4	Age-dependent survival in rapidly progressive glomerulonephritis: A nationwide questionnaire survey from children to the elderly. <i>PLoS ONE</i> , 2020, 15, e0236017.	1.1	3
5	Genetic Deletion of Vasohibin-2 Exacerbates Ischemia-Reperfusion-Induced Acute Kidney Injury. <i>International Journal of Molecular Sciences</i> , 2020, 21, 4545.	1.8	5
6	Podocyte autophagy is associated with foot process effacement and proteinuria in patients with minimal change nephrotic syndrome. <i>PLoS ONE</i> , 2020, 15, e0228337.	1.1	20
7	Prevalences of hyperuricemia and electrolyte abnormalities in patients with chronic kidney disease in Japan: A nationwide, cross-sectional cohort study using data from the Japan Chronic Kidney Disease Database (J-CKD-DB). <i>PLoS ONE</i> , 2020, 15, e0240402.	1.1	17
8	<p>Report of health checkup system for chronic kidney disease in general population in Okayama city: effect of health guidance intervention on chronic kidney disease outcome</p>. <i>International Journal of Nephrology and Renovascular Disease</i> , 2019, Volume 12, 143-152.	0.8	5
9	Urine 5MedC, a Marker of DNA Methylation, in the Progression of Chronic Kidney Disease. <i>Disease Markers</i> , 2019, 2019, 1-10.	0.6	11
10	The relationship between repeated measurement of casual and 24-h urinary sodium-to-potassium ratio in patients with chronic kidney disease. <i>Journal of Human Hypertension</i> , 2019, 33, 286-297.	1.0	7
11	Impaired mental health status in patients with chronic kidney disease is associated with estimated glomerular filtration rate decline. <i>Nephrology</i> , 2019, 24, 926-932.	0.7	3
12	Temporal change in life and renal prognosis of rapidly progressive glomerulonephritis in Japan via nationwide questionnaire survey. <i>Clinical and Experimental Nephrology</i> , 2019, 23, 573-575.	0.7	3
13	Histopathological classification of anti-neutrophil cytoplasmic antibody-associated glomerulonephritis in a nationwide Japanese prospective 2-year follow-up cohort study. <i>Clinical and Experimental Nephrology</i> , 2019, 23, 387-394.	0.7	9
14	Regional variations in immunosuppressive therapy in patients with primary nephrotic syndrome: the Japan nephrotic syndrome cohort study. <i>Clinical and Experimental Nephrology</i> , 2018, 22, 1266-1280.	0.7	21
15	Peripheral artery disease is associated with frailty in chronic hemodialysis patients. <i>Vascular</i> , 2018, 26, 425-431.	0.4	17
16	Estrogen-related receptor $\beta$ is essential for maintaining mitochondrial integrity in cisplatin-induced acute kidney injury. <i>Biochemical and Biophysical Research Communications</i> , 2018, 498, 918-924.	1.0	30
17	Clinical features and pathogenesis of membranoproliferative glomerulonephritis: a nationwide analysis of the Japan renal biopsy registry from 2007 to 2015. <i>Clinical and Experimental Nephrology</i> , 2018, 22, 797-807.	0.7	19
18	Diabetic nephropathy is associated with frailty in patients with chronic hemodialysis. <i>Geriatrics and Gerontology International</i> , 2018, 18, 1597-1602.	0.7	20

#	ARTICLE	IF	CITATIONS
19	Urine Trefoil Factors as Prognostic Biomarkers in Chronic Kidney Disease. <i>BioMed Research International</i> , 2018, 2018, 1-11.	0.9	14
20	Distinct characteristics and outcomes in elderly-onset IgA vasculitis (Henoch-Schönlein purpura) with nephritis: Nationwide cohort study of data from the Japan Renal Biopsy Registry (J-RBR). <i>PLoS ONE</i> , 2018, 13, e0196955.	1.1	22
21	The Prevalence of Frailty and its Associated Factors in Japanese Hemodialysis Patients. , 2018, 9, 192.		55
22	Renal expression of trefoil factor 3 mRNA in association with tubulointerstitial fibrosis in IgA nephropathy. <i>Nephrology</i> , 2018, 23, 855-862.	0.7	19
23	Deletion of pro-angiogenic factor vasohibin-2 ameliorates glomerular alterations in a mouse diabetic nephropathy model. <i>PLoS ONE</i> , 2018, 13, e0195779.	1.1	16
24	Serum cystatin C is an independent biomarker associated with the renal resistive index in patients with chronic kidney disease. <i>PLoS ONE</i> , 2018, 13, e0193695.	1.1	15
25	IgA Nephropathy Complicated with X-linked Thrombocytopenia. <i>Acta Medica Okayama</i> , 2018, 72, 301-307.	0.1	1
26	Performance in adolescents of the two Japanese serum creatinine based estimated glomerular filtration rate equations, for adults and paediatric patients: A study of the Japan Renal Biopsy Registry and Japan Kidney Disease Registry from 2007 to 2013. <i>Nephrology</i> , 2017, 22, 494-497.	0.7	8
27	Prediction of response to remission induction therapy by gene expression profiling of peripheral blood in Japanese patients with microscopic polyangiitis. <i>Arthritis Research and Therapy</i> , 2017, 19, 117.	1.6	10
28	Primary peritoneal carcinosarcoma in a dialysis patient. <i>Nephrology</i> , 2017, 22, 925-925.	0.7	0
29	Arterial Stiffness is an Independent Risk Factor for Anemia After Percutaneous Native Kidney Biopsy. <i>Kidney and Blood Pressure Research</i> , 2017, 42, 284-293.	0.9	11
30	A clinical evaluation of renal amyloidosis in the Japan renal biopsy registry: a cross-sectional study. <i>Clinical and Experimental Nephrology</i> , 2017, 21, 624-632.	0.7	7
31	Clinical and histological features of lupus nephritis in Japan: A cross-sectional analysis of the Japan Renal Biopsy Registry (J-RBR). <i>Nephrology</i> , 2017, 22, 885-891.	0.7	12
32	Sustained Tubulointerstitial Inflammation in Kidney with Severe Leptospirosis. <i>Internal Medicine</i> , 2017, 56, 1179-1184.	0.3	16
33	A retrospective observational study of glucocorticoid-induced diabetes mellitus with IgA nephropathy treated with tonsillectomy plus methylprednisolone pulse therapy. <i>PLoS ONE</i> , 2017, 12, e0178018.	1.1	10
34	Paratubular basement membrane insudative lesions predict renal prognosis in patients with type 2 diabetes and biopsy-proven diabetic nephropathy. <i>PLoS ONE</i> , 2017, 12, e0183190.	1.1	14
35	The possible involvement of intestine-derived IgA1: a case of IgA nephropathy associated with Crohn's disease. <i>BMC Nephrology</i> , 2016, 17, 122.	0.8	18
36	Comparison of severity classification in Japanese patients with antineutrophil cytoplasmic antibody-associated vasculitis in a nationwide, prospective, inception cohort study. <i>Modern Rheumatology</i> , 2016, 26, 730-737.	0.9	39

#	ARTICLE	IF	CITATIONS
37	Drug-induced kidney disease: a study of the Japan Renal Biopsy Registry from 2007 to 2015. <i>Clinical and Experimental Nephrology</i> , 2016, 20, 720-730.	0.7	18
38	Clinical manifestations of Henoch-Schönlein purpura nephritis and IgA nephropathy: comparative analysis of data from the Japan Renal Biopsy Registry (J-RBR). <i>Clinical and Experimental Nephrology</i> , 2016, 20, 552-560.	0.7	38
39	The Efficacy of Rituximab in High-risk Renal Transplant Recipients. <i>Acta Medica Okayama</i> , 2016, 70, 295-7.	0.1	0
40	Autoimmune pancreatitis and minimal change nephrotic syndrome: an unusual association?. <i>Nephrology</i> , 2015, 20, 225-226.	0.7	2
41	Pulmonary involvements of anti-neutrophil cytoplasmic autoantibody-associated renal vasculitis in Japan. <i>Nephrology Dialysis Transplantation</i> , 2015, 30, i83-i93.	0.4	47
42	Outcomes of primary nephrotic syndrome in elderly Japanese: retrospective analysis of the Japan Renal Biopsy Registry (J-RBR). <i>Clinical and Experimental Nephrology</i> , 2015, 19, 496-505.	0.7	23
43	Successful treatment by mycophenolate mofetil in a patient with focal segmental glomerulosclerosis associated with posterior reversible encephalopathy syndrome. <i>CEN Case Reports</i> , 2015, 4, 190-195.	0.5	3
44	The urinary levels of prostanoid metabolites predict acute kidney injury in heterogeneous adult Japanese ICU patients: a prospective observational study. <i>Clinical and Experimental Nephrology</i> , 2015, 19, 1024-1036.	0.7	5
45	The influences of larger physical constitutions including obesity on the amount of urine protein excretion in primary glomerulonephritis: research of the Japan Renal Biopsy Registry. <i>Clinical and Experimental Nephrology</i> , 2015, 19, 359-370.	0.7	8
46	ONO-1301, a sustained-release prostacyclin analog, ameliorates the renal alterations in a mouse type 2 diabetes model possibly through its protective effects on mesangial cells. <i>Acta Medica Okayama</i> , 2015, 69, 1-15.	0.1	3
47	Urinary and Plasma Levels of Vasohibin-1 Can Predict Renal Functional Deterioration in Patients with Renal Disorders. <i>PLoS ONE</i> , 2014, 9, e96932.	1.1	15
48	Exacerbation of Diabetic Renal Alterations in Mice Lacking Vasohibin-1. <i>PLoS ONE</i> , 2014, 9, e107934.	1.1	25
49	Renal distribution of Vasohibin-1 in patients with chronic kidney disease. <i>Acta Medica Okayama</i> , 2014, 68, 219-33.	0.1	5
50	Japan Renal Biopsy Registry and Japan Kidney Disease Registry: Committee Report for 2009 and 2010. <i>Clinical and Experimental Nephrology</i> , 2013, 17, 155-173.	0.7	111
51	Clinical findings on ANCA-associated renal vasculitis from the Japan RPGN registry obtained via a questionnaire survey. <i>Clinical and Experimental Nephrology</i> , 2013, 17, 646-649.	0.7	5
52	The Selection of Peritoneal Mesothelial Cells Is Important for Cell Therapy to Prevent Peritoneal Fibrosis. <i>Tissue Engineering - Part A</i> , 2013, 20, 131203154812003.	1.6	9
53	Effect of vitamin E on alloxan-induced mouse diabetes. <i>Clinical Biochemistry</i> , 2013, 46, 795-798.	0.8	23
54	A Decreased Level of Serum Soluble Klotho Is an Independent Biomarker Associated with Arterial Stiffness in Patients with Chronic Kidney Disease. <i>PLoS ONE</i> , 2013, 8, e56695.	1.1	167

#	ARTICLE	IF	CITATIONS
55	Reversible Posterior Leukoencephalopathy Syndrome in a Young Adult Patient Receiving Peritoneal Dialysis. <i>Peritoneal Dialysis International</i> , 2012, 32, 587-589.	1.1	12
56	Effluent Free Radicals Are Associated with Residual Renal Function and Predict Technique Failure in Peritoneal Dialysis Patients. <i>Peritoneal Dialysis International</i> , 2012, 32, 453-461.	1.1	22
57	Sustained-release prostacyclin analog ONO-1301 ameliorates tubulointerstitial alterations in a mouse obstructive nephropathy model. <i>American Journal of Physiology - Renal Physiology</i> , 2012, 302, F1616-F1629.	1.3	30
58	Right hypoplastic kidney. <i>Kidney International</i> , 2012, 82, 1037.	2.6	1
59	Peritoneovenous shunting for refractory ascites results in worsening of nephrotic syndrome. <i>Hepatology Research</i> , 2012, 42, 1048-1053.	1.8	4
60	ANCA-associated systemic vasculitis in Japan: clinical features and prognostic changes. <i>Clinical and Experimental Nephrology</i> , 2012, 16, 580-588.	0.7	54
61	Membranous nephropathy in Japan: analysis of the Japan Renal Biopsy Registry (J-RBR). <i>Clinical and Experimental Nephrology</i> , 2012, 16, 557-563.	0.7	67
62	Renal disease in the elderly and the very elderly Japanese: analysis of the Japan Renal Biopsy Registry (J-RBR). <i>Clinical and Experimental Nephrology</i> , 2012, 16, 903-920.	0.7	91
63	Acatalasemic mice are mildly susceptible to adriamycin nephropathy and exhibit increased albuminuria and glomerulosclerosis. <i>BMC Nephrology</i> , 2012, 13, 14.	0.8	13
64	A case of focal segmental glomerulosclerosis in an adult patient with hypogammaglobulinemia superimposed on membranoproliferative glomerulonephritis in childhood. <i>BMC Nephrology</i> , 2012, 13, 46.	0.8	2
65	Suppression of Adiponectin by Aberrantly Glycosylated IgA1 in Glomerular Mesangial Cells In Vitro and In Vivo. <i>PLoS ONE</i> , 2012, 7, e33965.	1.1	14
66	The therapeutic potential of synthetic human atrial natriuretic peptide in nephrotic syndrome: a randomized controlled trial. <i>International Journal of Nephrology and Renovascular Disease</i> , 2012, 5, 91.	0.8	2
67	The frequency of Fabry disease with the E66Q variant in the $\beta$ -galactosidase A gene in Japanese dialysis patients: a case report and a literature review. <i>Clinical Nephrology</i> , 2012, 78, 224-229.	0.4	11
68	Intermittent administration of a sustained-release prostacyclin analog ONO-1301 ameliorates renal alterations in a rat type 1 diabetes model. <i>Prostaglandins Leukotrienes and Essential Fatty Acids</i> , 2011, 84, 99-107.	1.0	6
69	Development of Angiotensin II-induced Abdominal Aortic Aneurysms Is Independent of Catalase in Mice. <i>Journal of Cardiovascular Pharmacology</i> , 2011, 58, 633-638.	0.8	10
70	Japan Renal Biopsy Registry: the first nationwide, web-based, and prospective registry system of renal biopsies in Japan. <i>Clinical and Experimental Nephrology</i> , 2011, 15, 493-503.	0.7	127
71	Abnormalities of Glycogenes in Tonsillar Lymphocytes in IgA Nephropathy. <i>Advances in Oto-Rhino-Laryngology</i> , 2011, 72, 71-74.	1.6	12
72	Serum High-Sensitivity Cardiac Troponin T Is a Significant Biomarker of Left-Ventricular Diastolic Dysfunction in Subjects with Non-Diabetic Chronic Kidney Disease. <i>Nephron Extra</i> , 2011, 1, 166-177.	1.1	21

#	ARTICLE	IF	CITATIONS
73	Amelioration of renal alterations in obese type 2 diabetic mice by vasohibin-1, a negative feedback regulator of angiogenesis. <i>American Journal of Physiology - Renal Physiology</i> , 2011, 300, F873-F886.	1.3	48
74	Icodextrin Increases Technique Survival Rate in Peritoneal Dialysis Patients with Diabetic Nephropathy by Improving Body Fluid Management: A Randomized Controlled Trial. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2011, 6, 1337-1344.	2.2	77
75	Fabry Disease Exhibiting Recurrent Stroke and Persistent Inflammation. <i>Internal Medicine</i> , 2010, 49, 2247-2252.	0.3	13
76	Differential expression of glycogenes in tonsillar B lymphocytes in association with proteinuria and renal dysfunction in IgA nephropathy. <i>Clinical Immunology</i> , 2010, 136, 447-455.	1.4	44
77	Effects of icodextrin peritoneal dialysis solution on the peritoneal membrane in the STZ-induced diabetic rat model with partial nephrectomy. <i>Nephrology Dialysis Transplantation</i> , 2010, 25, 1479-1488.	0.4	16
78	C-reactive protein is associated with cigarette smoking-induced hyperfiltration and proteinuria in an apparently healthy population. <i>Hypertension Research</i> , 2010, 33, 1129-1136.	1.5	25
79	Sensitization to alloxan-induced diabetes and pancreatic cell apoptosis in acatalasemic mice. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2010, 1802, 240-246.	1.8	43
80	Vasohibin-1, a Negative Feedback Regulator of Angiogenesis, Ameliorates Renal Alterations in a Mouse Model of Diabetic Nephropathy. <i>Diabetes</i> , 2009, 58, 2365-2375.	0.3	65
81	Clinical Usefulness of a Prognostic Score in Histological Analysis of Renal Biopsy in Patients with Lupus Nephritis. <i>Journal of Rheumatology</i> , 2009, 36, 2218-2223.	1.0	30
82	A case of immunotactoid glomerulopathy exhibiting nephrotic syndrome successfully treated with corticosteroids and antihypertensive therapy. <i>Clinical and Experimental Nephrology</i> , 2009, 13, 378-384.	0.7	7
83	Unique microstructures and podocytic infolding in glomerular basement membrane associated with collagen diseases: a report of three cases. <i>Clinical and Experimental Nephrology</i> , 2008, 12, 450-454.	0.7	11
84	Proposal of podocytic infolding glomerulopathy as a new disease entity: a review of 25 cases from nationwide research in Japan. <i>Clinical and Experimental Nephrology</i> , 2008, 12, 421-431.	0.7	36
85	Novel mutations of the GLA gene in Japanese patients with Fabry disease and their functional characterization by active site specific chaperone. <i>Human Mutation</i> , 2008, 29, 331-331.	1.1	49
86	Increased Susceptibility to Oxidant-Mediated Tissue Injury and Peritoneal Fibrosis in Acatlasemic Mice. <i>American Journal of Nephrology</i> , 2008, 28, 661-668.	1.4	24
87	Amelioration of Cisplatin-Induced Acute Renal Injury by Renal Progenitor-Like Cells Derived from the Adult Rat Kidney. <i>Cell Transplantation</i> , 2008, 17, 143-158.	1.2	43
88	A novel variant apolipoprotein E Okayama in a patient with lipoprotein glomerulopathy. <i>Nephrology Dialysis Transplantation</i> , 2007, 23, 751-756.	0.4	31
89	Endostatin peptide, an inhibitor of angiogenesis, prevents the progression of peritoneal sclerosis in a mouse experimental model. <i>Kidney International</i> , 2007, 71, 227-238.	2.6	75
90	Elevated Serum sFlt-1/Ang-2 Ratio in Women with Preeclampsia. <i>Nephron Clinical Practice</i> , 2007, 106, c43-c50.	2.3	29

#	ARTICLE	IF	CITATIONS
91	Downregulation of the $\beta$ 1,3-Galactosyltransferase Gene in Tonsillar B Lymphocytes and Aberrant Lectin Bindings to Tonsillar IgA as a Pathogenesis of IgA Nephropathy. , 2007, 157, 120-124.		15
92	Enhanced TGF- $\beta$ /Smad signaling in the early stage of diabetic nephropathy is independent of the AT1a receptor. Clinical and Experimental Nephrology, 2007, 11, 77-87.	0.7	21
93	Desmin as a marker of proteinuria in early stages of membranous nephropathy in elderly patients. Clinical Nephrology, 2007, 68, 73-80.	0.4	12
94	Regulation of Angiogenic Factors in Angiotensin II Infusion Model in Association With Tubulointerstitial Injuries. American Journal of Hypertension, 2006, 19, 718-727.	1.0	41
95	An Elderly Patient with Severe Acute Renal Failure Due to Sodium Bromate Intoxication. Internal Medicine, 2006, 45, 151-154.	0.3	11
96	Steroid pulse therapy impaired endothelial function while increasing plasma high molecule adiponectin concentration in patients with IgA nephropathy. Nephrology Dialysis Transplantation, 2006, 21, 3475-3480.	0.4	30
97	Characterization of Acatlasemic Erythrocytes Treated with Low and High Dose Hydrogen Peroxide. Journal of Biological Chemistry, 2006, 281, 21728-21734.	1.6	15
98	Good response of membranous lupus nephritis to tacrolimus. Clinical Nephrology, 2006, 65, 276-279.	0.4	16
99	Successful Treatment of Progressive Henoch-Schoenlein Purpura Nephritis with Tonsillectomy and Steroid Pulse Therapy. Internal Medicine, 2005, 44, 611-615.	0.3	23
100	Infusion of angiotensin II reduces loss of glomerular capillary area in the early phase of anti-Thy-1.1 nephritis possibly via regulating angiogenesis-associated factors. Kidney International, 2005, 68, 704-722.	2.6	18
101	Catalase deficiency renders remnant kidneys more susceptible to oxidant tissue injury and renal fibrosis in mice. Kidney International, 2005, 68, 1018-1031.	2.6	77
102	Establishment and characterization of renal progenitor like cells from S3 segment of nephron in rat adult kidney. FASEB Journal, 2005, 19, 1789-1797.	0.2	178
103	Antiangiogenic Endostatin Peptide Ameliorates Renal Alterations in the Early Stage of a Type 1 Diabetic Nephropathy Model. Diabetes, 2005, 54, 2891-2903.	0.3	129
104	Telmisartan inhibits both oxidative stress and renal fibrosis after unilateral ureteral obstruction in acatalasemic mice. Nephrology Dialysis Transplantation, 2005, 20, 2670-2680.	0.4	80
105	Increase of Serum Angiopoietin-2 During Pregnancy Is Suppressed in Women With Preeclampsia. American Journal of Hypertension, 2005, 18, 1181-1188.	1.0	44
106	A case of monoclonal immunoglobulin light- and heavy-chain deposition disease exhibiting atypical deposition with fibrillary structures, successfully treated with chemotherapy. Clinical Nephrology, 2005, 64, 221-227.	0.4	6
107	Overexpression of angiotensin type 2 receptor ameliorates glomerular injury in a mouse remnant kidney model. American Journal of Physiology - Renal Physiology, 2004, 286, F516-F525.	1.3	62
108	Tumstatin Peptide, an Inhibitor of Angiogenesis, Prevents Glomerular Hypertrophy in the Early Stage of Diabetic Nephropathy. Diabetes, 2004, 53, 1831-1840.	0.3	169

#	ARTICLE	IF	CITATIONS
109	Acatalasemia sensitizes renal tubular epithelial cells to apoptosis and exacerbates renal fibrosis after unilateral ureteral obstruction. <i>American Journal of Physiology - Renal Physiology</i> , 2004, 286, F1030-F1038.	1.3	59
110	Mesangial cell Fas ligand: upregulation in human lupus nephritis and NF- $\kappa$ B-mediated expression in cultured human mesangial cells. <i>Clinical and Experimental Nephrology</i> , 2004, 8, 196-205.	0.7	25
111	Glomerular cell apoptosis in human lupus nephritis. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2003, 443, 67-77.	1.4	36
112	Pathogenesis of IgA nephropathy. <i>Seminars in Nephrology</i> , 2003, 23, 556-563.	0.6	32
113	Transforming Growth Factor- $\beta$ 1 Induces Vascular Endothelial Growth Factor Expression in Murine Proximal Tubular Epithelial Cells. <i>Nephron Experimental Nephrology</i> , 2003, 95, e79-e86.	2.4	29
114	Implication of Apoptosis in Progression of Renal Diseases. , 2003, 139, 156-172.		15
115	Cyanotic congenital heart disease associated with glomerulomegaly and focal segmental glomerulosclerosis: remission of nephrotic syndrome with angiotensin converting enzyme inhibitor. <i>Nephrology Dialysis Transplantation</i> , 2002, 17, 144-147.	0.4	17
116	Fulminant necrotising fasciitis developing during long term corticosteroid treatment of systemic lupus erythematosus. <i>Annals of the Rheumatic Diseases</i> , 2002, 61, 848-849.	0.5	15
117	Minimal Change Nephrotic Syndrome Developing during Postoperative Interferon-Beta Therapy for Malignant Melanoma. <i>Nephron</i> , 2002, 90, 498-500.	0.9	22
118	Methylprednisolone Accelerates the Resolution of Glomerulonephritis by Sensitizing Mesangial Cells to Apoptosis. <i>Nephron Experimental Nephrology</i> , 2001, 9, 317-326.	2.4	8
119	Expression, regulation, and function of inhibitor of apoptosis family genes in rat mesangial cells. <i>Kidney International</i> , 2001, 60, 579-586.	2.6	11
120	Apoptosis and extracellular matrix-cell interactions in kidney disease. <i>Kidney International</i> , 2000, 58, S67-S75.	2.6	62
121	Lewis <sup>x</sup> Expression and Renal Tubular Atrophy in IgA Nephritis. <i>Nephron</i> , 2000, 84, 274-275.	0.9	0
122	Caldesmon Isoform Associated with Phenotypic Modulation of Mesangial Cells. <i>Nephron Experimental Nephrology</i> , 2000, 8, 20-27.	2.4	4
123	Mechanisms of induction of apoptosis and glomerular disease. <i>Nephrology Dialysis Transplantation</i> , 1999, 14, 52-54.	0.4	9
124	Selective Sensitization to Tumor Necrosis Factor- $\alpha$ -induced Apoptosis by Blockade of NF- $\kappa$ B in Primary Glomerular Mesangial Cells. <i>Journal of Biological Chemistry</i> , 1999, 274, 19532-19537.	1.6	63
125	Regulation of survival and death of mesangial cells by extracellular matrix. <i>Kidney International</i> , 1998, 54, 1188-1196.	2.6	44
126	Inhibition of mesangial cell proliferation by E2F decoy oligodeoxynucleotide in vitro and in vivo.. <i>Journal of Clinical Investigation</i> , 1998, 101, 2589-2597.	3.9	86



#	ARTICLE	IF	CITATIONS
127	Treatment of Severe Thrombocytopenia with Intravenous Immunoglobulins and Corticosteroids in a Patient Receiving Continuous Ambulatory Peritoneal Dialysis. <i>Nephron</i> , 1997, 77, 371-372.	0.6	1
128	Role of Apoptosis in the Progression of Glomerulosclerosis. <i>Contributions To Nephrology</i> , 1996, 118, 41-47.	1.1	9
129	Apoptosis in glomerular sclerosis. <i>Kidney International</i> , 1996, 49, 103-111.	2.6	235
130	Phenotypic modulation of the mesangium reflected by contractile proteins in diabetes. <i>Diabetes</i> , 1996, 45, 488-495.	0.3	27
131	Transition of Morphologic Features in Lupus Nephritis: Does Steroid Therapy Accelerate Glomerulosclerosis?. <i>Internal Medicine</i> , 1995, 34, 982-987.	0.3	4
132	Phenotypic changes of the mesangium in diabetic nephropathy. <i>Journal of Diabetes and Its Complications</i> , 1995, 9, 282-284.	1.2	22