

# Cindy Sander

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4376176/publications.pdf>

Version: 2024-02-01

15  
papers

2,724  
citations

686830

13  
h-index

996533

15  
g-index

15  
all docs

15  
docs citations

15  
times ranked

5050  
citing authors

#	ARTICLE	IF	CITATIONS
1	Tim-3 mediates T cell trogocytosis to limit antitumor immunity. <i>Journal of Clinical Investigation</i> , 2022, 132, .	3.9	25
2	Predicting Metastasis in Melanoma by Enumerating Circulating Tumor Cells Using Photoacoustic Flow Cytometry. <i>Lasers in Surgery and Medicine</i> , 2021, 53, 578-586.	1.1	10
3	Neoadjuvant Pembrolizumab and High-Dose IFN $\alpha$ -2b in Resectable Regionally Advanced Melanoma. <i>Clinical Cancer Research</i> , 2021, 27, 4195-4204.	3.2	18
4	IL15 Stimulation with TIGIT Blockade Reverses CD155-mediated NK-Cell Dysfunction in Melanoma. <i>Clinical Cancer Research</i> , 2020, 26, 5520-5533.	3.2	88
5	Neratinib and entinostat combine to rapidly reduce the expression of K-RAS, N-RAS, G $\alpha$ <sub>q</sub> and G $\alpha$ <sub>11</sub> and kill uveal melanoma cells. <i>Cancer Biology and Therapy</i> , 2019, 20, 700-710.	1.5	37
6	Tumor cell oxidative metabolism as a barrier to PD-1 blockade immunotherapy in melanoma. <i>JCI Insight</i> , 2019, 4, .	2.3	148
7	The levels of mutant K-RAS and mutant N-RAS are rapidly reduced in a Beclin1 / ATG5 -dependent fashion by the irreversible ERBB1/2/4 inhibitor neratinib. <i>Cancer Biology and Therapy</i> , 2018, 19, 132-137.	1.5	32
8	Phase Ib/II Study of Pembrolizumab and Pegylated-Interferon Alfa-2b in Advanced Melanoma. <i>Journal of Clinical Oncology</i> , 2018, 36, 3450-3458.	0.8	55
9	CD226 opposes TIGIT to disrupt Tregs in melanoma. <i>JCI Insight</i> , 2018, 3, .	2.3	137
10	Expression profiles of immune-related genes are associated with neoadjuvant ipilimumab clinical benefit. <i>Oncolmmunology</i> , 2017, 6, e1231291.	2.1	29
11	The HDAC inhibitor AR42 interacts with pazopanib to kill trametinib/dabrafenib-resistant melanoma cells <i>in vitro</i> and <i>in vivo</i> . <i>Oncotarget</i> , 2017, 8, 16367-16386.	0.8	55
12	TIGIT and PD-1 impair tumor antigen-specific CD8+ T cells in melanoma patients. <i>Journal of Clinical Investigation</i> , 2015, 125, 2046-2058.	3.9	603
13	Immune Monitoring of the Circulation and the Tumor Microenvironment in Patients with Regionally Advanced Melanoma Receiving Neoadjuvant Ipilimumab. <i>PLoS ONE</i> , 2014, 9, e87705.	1.1	261
14	Upregulation of Tim-3 and PD-1 expression is associated with tumor antigen-specific CD8+ T cell dysfunction in melanoma patients. <i>Journal of Experimental Medicine</i> , 2010, 207, 2175-2186.	4.2	1,118
15	Immunization With Analog Peptide in Combination With CpG and Montanide Expands Tumor Antigen-specific CD8+ T Cells in Melanoma Patients. <i>Journal of Immunotherapy</i> , 2008, 31, 781-791.	1.2	108