

Yong J Lee

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/437842/yong-j-lee-publications-by-year.pdf>
Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

114 papers	8,879 citations	40 h-index	93 g-index
118 ext. papers	9,843 ext. citations	5.8 avg, IF	5.44 L-index

#	Paper	IF	Citations
114	Improved chemosensitivity following mucolytic therapy in patient-derived models of mucinous appendix cancer. <i>Translational Research</i> , 2021 , 229, 100-114	11	2
113	Glucose deprivation-induced endoplasmic reticulum stress response plays a pivotal role in enhancement of TRAIL cytotoxicity. <i>Journal of Cellular Physiology</i> , 2021 , 236, 6666-6677	7	2
112	The anti-fibrotic drug pirfenidone inhibits liver fibrosis by targeting the small oxidoreductase glutaredoxin-1. <i>Science Advances</i> , 2021 , 7, eabg9241	14.3	3
111	The emerging role of selenium metabolic pathways in cancer: New therapeutic targets for cancer.. <i>Journal of Cellular Biochemistry</i> , 2021 ,	4.7	3
110	The interplay between apoptosis and ferroptosis mediated by ER stress. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2020 , 25, 783	5.4	0
109	Ferroptotic agent-induced endoplasmic reticulum stress response plays a pivotal role in the autophagic process outcome. <i>Journal of Cellular Physiology</i> , 2020 , 235, 6767-6778	7	14
108	BAX-dependent mitochondrial pathway mediates the crosstalk between ferroptosis and apoptosis. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2020 , 25, 625-631	5.4	20
107	Synergistic apoptosis following endoplasmic reticulum stress aggravation in mucinous colon cancer. <i>Orphanet Journal of Rare Diseases</i> , 2020 , 15, 211	4.2	3
106	Biological Aspects of Endoplasmic Reticulum Stress in Ferroptosis 2019 , 83-98		
105	Ferroptosis-inducing agents enhance TRAIL-induced apoptosis through upregulation of death receptor 5. <i>Journal of Cellular Biochemistry</i> , 2019 , 120, 928-939	4.7	26
104	Crosstalk Between Apoptosis and Autophagy Is Regulated by the Arginylated BiP/Beclin-1/p62 Complex. <i>Molecular Cancer Research</i> , 2018 , 16, 1077-1091	6.6	26
103	Ferroptosis-Induced Endoplasmic Reticulum Stress: Cross-talk between Ferroptosis and Apoptosis. <i>Molecular Cancer Research</i> , 2018 , 16, 1073-1076	6.6	105
102	PARK7 modulates autophagic proteolysis through binding to the N-terminally arginylated form of the molecular chaperone HSPA5. <i>Autophagy</i> , 2018 , 14, 1870-1885	10.2	14
101	Glioma-derived cancer stem cells are hypersensitive to proteasomal inhibition. <i>EMBO Reports</i> , 2017 , 18, 150-168	6.5	19
100	Molecular crosstalk between ferroptosis and apoptosis: emerging role of ER stress-induced p53-independent PUMA expression. <i>Oncotarget</i> , 2017 , 8, 115164-115178	3.3	71
99	TRAIL-Induced Caspase Activation Is a Prerequisite for Activation of the Endoplasmic Reticulum Stress-Induced Signal Transduction Pathways. <i>Journal of Cellular Biochemistry</i> , 2016 , 117, 1078-91	4.7	10
98	Cancer Stem Cells Protect Non-Stem Cells From Anoikis: Bystander Effects. <i>Journal of Cellular Biochemistry</i> , 2016 , 117, 2289-301	4.7	26

97	Guidelines for the use and interpretation of assays for monitoring autophagy (3rd edition). <i>Autophagy</i> , 2016 , 12, 1-222	10.2	3838
96	Secretory TRAIL-Armed Natural Killer Cell-Based Therapy: In Vitro and In Vivo Colorectal Peritoneal Carcinomatosis Xenograft. <i>Molecular Cancer Therapeutics</i> , 2016 , 15, 1591-601	6.1	9
95	Hypoxia Promotes Synergy between Mitomycin C and Bortezomib through a Coordinated Process of Bcl-xL Phosphorylation and Mitochondrial Translocation of p53. <i>Molecular Cancer Research</i> , 2015 , 13, 1533-43	6.6	5
94	HSP90 inhibitor NVP-AUY922 enhances TRAIL-induced apoptosis by suppressing the JAK2-STAT3-Mcl-1 signal transduction pathway in colorectal cancer cells. <i>Cellular Signalling</i> , 2015 , 27, 293-305	4.9	39
93	Gingerol sensitizes TRAIL-induced apoptotic cell death of glioblastoma cells. <i>Toxicology and Applied Pharmacology</i> , 2014 , 279, 253-265	4.6	50
92	Role of Bcl-xL/Beclin-1 in synergistic apoptotic effects of secretory TRAIL-armed adenovirus in combination with mitomycin C and hyperthermia on colon cancer cells. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2014 , 19, 1603-15	5.4	9
91	Role of Bcl-xL/Beclin-1 in interplay between apoptosis and autophagy in oxaliplatin and bortezomib-induced cell death. <i>Biochemical Pharmacology</i> , 2014 , 88, 178-88	6	43
90	Role of the IL-6-JAK1-STAT3-Oct-4 pathway in the conversion of non-stem cancer cells into cancer stem-like cells. <i>Cellular Signalling</i> , 2013 , 25, 961-9	4.9	199
89	Evidence for two modes of synergistic induction of apoptosis by mapatumumab and oxaliplatin in combination with hyperthermia in human colon cancer cells. <i>PLoS ONE</i> , 2013 , 8, e73654	3.7	12
88	The role of Bcl-xL in synergistic induction of apoptosis by mapatumumab and oxaliplatin in combination with hyperthermia on human colon cancer. <i>Molecular Cancer Research</i> , 2012 , 10, 1567-79	6.6	24
87	Hyperthermia-enhanced TRAIL- and mapatumumab-induced apoptotic death is mediated through mitochondria in human colon cancer cells. <i>Journal of Cellular Biochemistry</i> , 2012 , 113, 1547-58	4.7	12
86	Breast cancer stem cell-like cells are more sensitive to ionizing radiation than non-stem cells: role of ATM. <i>PLoS ONE</i> , 2012 , 7, e50423	3.7	20
85	Astaxanthin protects against MPTP/MPP+-induced mitochondrial dysfunction and ROS production in vivo and in vitro. <i>Food and Chemical Toxicology</i> , 2011 , 49, 271-80	4.7	150
84	Preferential accumulation within tumors and in vivo imaging by functionalized luminescent dendrimer lanthanide complexes. <i>Biomaterials</i> , 2011 , 32, 9343-52	15.6	24
83	MEKK1/MEKK4 are responsible for TRAIL-induced JNK/p38 phosphorylation. <i>Oncology Reports</i> , 2011 , 25, 537-44	3.5	20
82	Role of Bim in diallyl trisulfide-induced cytotoxicity in human cancer cells. <i>Journal of Cellular Biochemistry</i> , 2011 , 112, 118-27	4.7	28
81	Luminescence targeting and imaging using a nanoscale generation 3 dendrimer in an in vivo colorectal metastatic rat model. <i>Nanomedicine: Nanotechnology, Biology, and Medicine</i> , 2011 , 7, 249-58	6	25
80	TRAIL-induced caspase/p38 activation is responsible for the increased catalytic and invasive activities of Akt. <i>International Journal of Oncology</i> , 2011 , 38, 249-56	4.4	5

79	Quercetin enhances TRAIL-induced apoptosis in prostate cancer cells via increased protein stability of death receptor 5. <i>Life Sciences</i> , 2010 , 86, 351-7	6.8	78
78	c-Cbl acts as a mediator of Src-induced activation of the PI3K-Akt signal transduction pathway during TRAIL treatment. <i>Cellular Signalling</i> , 2010 , 22, 377-85	4.9	32
77	c-Cbl-mediated degradation of TRAIL receptors is responsible for the development of the early phase of TRAIL resistance. <i>Cellular Signalling</i> , 2010 , 22, 553-63	4.9	42
76	Effect of hyperthermia in combination with TRAIL on the JNK-Bim signal transduction pathway and growth of xenograft tumors. <i>Journal of Cellular Biochemistry</i> , 2010 , 110, 1073-81	4.7	15
75	Effects of low dose quercetin: cancer cell-specific inhibition of cell cycle progression. <i>Journal of Cellular Biochemistry</i> , 2009 , 106, 73-82	4.7	224
74	Magnolol induces apoptosis via inhibiting the EGFR/PI3K/Akt signaling pathway in human prostate cancer cells. <i>Journal of Cellular Biochemistry</i> , 2009 , 106, 1113-22	4.7	90
73	Reactive oxygen species up-regulate p53 and Puma; a possible mechanism for apoptosis during combined treatment with TRAIL and wogonin. <i>British Journal of Pharmacology</i> , 2009 , 157, 1189-202	8.6	70
72	Quercetin augments TRAIL-induced apoptotic death: involvement of the ERK signal transduction pathway. <i>Biochemical Pharmacology</i> , 2008 , 75, 1946-58	6	135
71	Role of p53, PUMA, and Bax in wogonin-induced apoptosis in human cancer cells. <i>Biochemical Pharmacology</i> , 2008 , 75, 2020-33	6	97
70	Role of Bax in quercetin-induced apoptosis in human prostate cancer cells. <i>Biochemical Pharmacology</i> , 2008 , 75, 2345-55	6	82
69	Differential cleavage of Mst1 by caspase-7/-3 is responsible for TRAIL-induced activation of the MAPK superfamily. <i>Cellular Signalling</i> , 2008 , 20, 892-906	4.9	57
68	Effect of hyperthermia and chemotherapeutic agents on TRAIL-induced cell death in human colon cancer cells. <i>Journal of Cellular Biochemistry</i> , 2008 , 103, 98-109	4.7	17
67	Flavonoids-induced accumulation of hypoxia-inducible factor (HIF)-1alpha/2alpha is mediated through chelation of iron. <i>Journal of Cellular Biochemistry</i> , 2008 , 103, 1989-98	4.7	63
66	Effect of UV irradiation on colorectal cancer cells with acquired TRAIL resistance. <i>Journal of Cellular Biochemistry</i> , 2008 , 104, 1172-80	4.7	7
65	Pretreatment of docetaxel enhances TRAIL-mediated apoptosis in prostate cancer cells. <i>Journal of Cellular Biochemistry</i> , 2008 , 104, 1636-46	4.7	24
64	Quercetin suppresses hypoxia-induced accumulation of hypoxia-inducible factor-1alpha (HIF-1alpha) through inhibiting protein synthesis. <i>Journal of Cellular Biochemistry</i> , 2008 , 105, 546-53	4.7	53
63	Quercetin-induced ubiquitination and down-regulation of Her-2/neu. <i>Journal of Cellular Biochemistry</i> , 2008 , 105, 585-95	4.7	58
62	TRAIL apoptosis is enhanced by quercetin through Akt dephosphorylation. <i>Journal of Cellular Biochemistry</i> , 2007 , 100, 998-1009	4.7	75

61	Effect of hyperthermia on TRAIL-induced apoptotic death in human colon cancer cells: development of a novel strategy for regional therapy. <i>Journal of Cellular Biochemistry</i> , 2007 , 101, 619-30	4.7	9
60	Evidence for two modes of development of acquired tumor necrosis factor-related apoptosis-inducing ligand resistance. Involvement of Bcl-xL. <i>Journal of Biological Chemistry</i> , 2007 , 282, 319-28	5.4	53
59	Time sequence of tumor necrosis factor-related apoptosis-inducing ligand (TRAIL) and cisplatin treatment is responsible for a complex pattern of synergistic cytotoxicity. <i>Journal of Cellular Biochemistry</i> , 2006 , 98, 1284-95	4.7	16
58	Hyperthermia enhances tumour necrosis factor-related apoptosis-inducing ligand (TRAIL)-induced apoptosis in human cancer cells. <i>International Journal of Hyperthermia</i> , 2006 , 22, 713-28	3.7	12
57	Sulforaphane-induced cell death in human prostate cancer cells is initiated by reactive oxygen species. <i>Journal of Biological Chemistry</i> , 2005 , 280, 19911-24	5.4	290
56	Amiloride augments TRAIL-induced apoptotic death by inhibiting phosphorylation of kinases and phosphatases associated with the P13K-Akt pathway. <i>Oncogene</i> , 2005 , 24, 355-66	9.2	39
55	Hypoxia and low glucose differentially augments TRAIL-induced apoptotic death. <i>Molecular and Cellular Biochemistry</i> , 2005 , 270, 89-97	4.2	15
54	Dissociation of Akt1 from its negative regulator JIP1 is mediated through the ASK1-MEK-JNK signal transduction pathway during metabolic oxidative stress: a negative feedback loop. <i>Journal of Cell Biology</i> , 2005 , 170, 61-72	7.3	61
53	Signal pathway of hypoxia-inducible factor-1alpha phosphorylation and its interaction with von Hippel-Lindau tumor suppressor protein during ischemia in MiaPaCa-2 pancreatic cancer cells. <i>Clinical Cancer Research</i> , 2005 , 11, 7607-13	12.9	56
52	Pretreatment of acetylsalicylic acid promotes tumor necrosis factor-related apoptosis-inducing ligand-induced apoptosis by down-regulating BCL-2 gene expression. <i>Journal of Biological Chemistry</i> , 2005 , 280, 41047-56	5.4	45
51	TRAIL and ceramide. <i>Vitamins and Hormones</i> , 2004 , 67, 229-55	2.5	7
50	Reconstitution of caspase-3 confers low glucose-enhanced tumor necrosis factor-related apoptosis-inducing ligand cytotoxicity and Akt cleavage. <i>Clinical Cancer Research</i> , 2004 , 10, 1894-900	12.9	13
49	Diallyl trisulfide-induced apoptosis in human prostate cancer cells involves c-Jun N-terminal kinase and extracellular-signal regulated kinase-mediated phosphorylation of Bcl-2. <i>Oncogene</i> , 2004 , 23, 5594-606	8.2	234
48	Daxx deletion mutant (amino acids 501-625)-induced apoptosis occurs through the JNK/p38-Bax-dependent mitochondrial pathway. <i>Journal of Cellular Biochemistry</i> , 2004 , 92, 1257-70	4.7	26
47	Low extracellular pH augments TRAIL-induced apoptotic death through the mitochondria-mediated caspase signal transduction pathway. <i>Experimental Cell Research</i> , 2004 , 293, 129-43	4.2	29
46	Role of the ASK1-SEK1-JNK1-HIPK1 signal in Daxx trafficking and ASK1 oligomerization. <i>Journal of Biological Chemistry</i> , 2003 , 278, 47245-52	5.4	65
45	Differential role of glutaredoxin and thioredoxin in metabolic oxidative stress-induced activation of apoptosis signal-regulating kinase 1. <i>Biochemical Journal</i> , 2003 , 373, 845-53	3.8	149
44	Catalase, but not MnSOD, inhibits glucose deprivation-activated ASK1-MEK-MAPK signal transduction pathway and prevents relocalization of Daxx: hydrogen peroxide as a major second messenger of metabolic oxidative stress. <i>Journal of Cellular Biochemistry</i> , 2003 , 90, 304-14	4.7	38

43	Reconstitution of galectin-3 alters glutathione content and potentiates TRAIL-induced cytotoxicity by dephosphorylation of Akt. <i>Experimental Cell Research</i> , 2003 , 288, 21-34	4.2	40
42	Gene transfer into human prostate adenocarcinoma cells with an adenoviral vector: Hyperthermia enhances a double suicide gene expression, cytotoxicity and radiotoxicity. <i>Cancer Gene Therapy</i> , 2002 , 9, 267-74	5.4	13
41	Low glucose-enhanced TRAIL cytotoxicity is mediated through the ceramide-Akt-FLIP pathway. <i>Oncogene</i> , 2002 , 21, 337-46	9.2	61
40	Enhancement of metabolic oxidative stress-induced cytotoxicity by the thioredoxin inhibitor 1-methylpropyl 2-imidazolyl disulfide is mediated through the ASK1-SEK1-JNK1 pathway. <i>Molecular Pharmacology</i> , 2002 , 62, 1409-17	4.3	16
39	Role of glutaredoxin in metabolic oxidative stress. Glutaredoxin as a sensor of oxidative stress mediated by H ₂ O ₂ . <i>Journal of Biological Chemistry</i> , 2002 , 277, 46566-75	5.4	209
38	Analysis of heat-shock transcription factor and element-binding activity. <i>Methods in Molecular Biology</i> , 2002 , 196, 131-8	1.4	
37	Role of galectin-3 in breast cancer metastasis: involvement of nitric oxide. <i>American Journal of Pathology</i> , 2002 , 160, 1069-75	5.8	69
36	Cooperative interaction between interleukin 10 and galectin-3 against liver ischemia-reperfusion injury. <i>Clinical Cancer Research</i> , 2002 , 8, 217-20	12.9	9
35	Replicating adenoviral vector-mediated transfer of a heat-inducible double suicide gene for gene therapy. <i>Cancer Gene Therapy</i> , 2001 , 8, 397-404	5.4	33
34	Sodium nitroprusside enhances TRAIL-induced apoptosis via a mitochondria-dependent pathway in human colorectal carcinoma CX-1 cells. <i>Oncogene</i> , 2001 , 20, 1476-85	9.2	64
33	Glucose deprivation-induced oxidative stress in human tumor cells. A fundamental defect in metabolism?. <i>Annals of the New York Academy of Sciences</i> , 2000 , 899, 349-62	6.5	243
32	Role of small heat shock protein HSP25 in radioresistance and glutathione-redox cycle. <i>Journal of Cellular Physiology</i> , 2000 , 183, 100-7	7	79
31	Dominant-negative Jun N-terminal protein kinase (JNK-1) inhibits metabolic oxidative stress during glucose deprivation in a human breast carcinoma cell line. <i>Free Radical Biology and Medicine</i> , 2000 , 28, 575-84	7.8	23
30	Hypoxia-induced bFGF gene expression is mediated through the JNK signal transduction pathway. <i>Molecular and Cellular Biochemistry</i> , 1999 , 202, 1-8	4.2	41
29	Adenoviral transduction of a cytosine deaminase/thymidine kinase fusion gene into prostate carcinoma cells enhances prodrug and radiation sensitivity. <i>International Journal of Cancer</i> , 1999 , 82, 293-7	7.5	24
28	The role of protein kinase Calpha in U-87 glioma invasion. <i>International Journal of Developmental Neuroscience</i> , 1999 , 17, 447-61	2.7	32
27	Differential induction of cell death in human glioma cell lines by sodium nitroprusside. <i>Cancer</i> , 1998 , 82, 1137-45	6.4	19
26	Overexpression of HSP25 reduces the level of TNF alpha-induced oxidative DNA damage biomarker, 8-hydroxy-2'-deoxyguanosine, in L929 cells. <i>Journal of Cellular Physiology</i> , 1998 , 174, 27-34	7	43

25	Glucose deprivation-induced cytotoxicity and alterations in mitogen-activated protein kinase activation are mediated by oxidative stress in multidrug-resistant human breast carcinoma cells. <i>Journal of Biological Chemistry</i> , 1998 , 273, 5294-9	5.4	176
24	Hypoglycemia-induced c-Jun phosphorylation is mediated by c-Jun N-terminal kinase 1 and Lyn kinase in drug-resistant human breast carcinoma MCF-7/ADR cells. <i>Journal of Biological Chemistry</i> , 1997 , 272, 11690-3	5.4	37
23	Examination of the molecular basis for the lack of alphaB-crystallin expression in L929 cells. <i>Molecular and Cellular Biochemistry</i> , 1997 , 170, 31-42	4.2	3
22	Differential effect of glucose deprivation on MAPK activation in drug sensitive human breast carcinoma MCF-7 and multidrug resistant MCF-7/ADR cells. <i>Molecular and Cellular Biochemistry</i> , 1997 , 170, 23-30	4.2	21
21	Comparison of tumor growth between hsp25- and hsp27-transfected murine L929 cells in nude mice. <i>International Journal of Cancer</i> , 1997 , 72, 871-7	7.5	24
20	Excess protein in nuclei isolated from heat-shocked cells results from a reduced extractability of nuclear proteins. <i>Journal of Cellular Physiology</i> , 1996 , 167, 369-79	7	25
19	Thermotolerance expression in mitotic CHO cells without increased translation of heat shock proteins. <i>Journal of Cellular Physiology</i> , 1996 , 169, 420-8	7	13
18	Thermal response in murine L929 cells lacking alpha B-crystallin expression and alpha B-crystallin expressing L929 transfectants. <i>Molecular and Cellular Biochemistry</i> , 1996 , 155, 51-60	4.2	15
17	Hypoglycemia-induced AP-1 transcription factor and basic fibroblast growth factor gene expression in multidrug resistant human breast carcinoma MCF-7/ADR cells. <i>Molecular and Cellular Biochemistry</i> , 1996 , 155, 163-71	4.2	15
16	Lack of radiosensitization after paclitaxel treatment of three human carcinoma cell lines. <i>Cancer</i> , 1995 , 75, 2262-8	6.4	41
15	Heat-induced bFGF gene expression in the absence of heat shock element correlates with enhanced AP-1 binding activity. <i>Journal of Cellular Physiology</i> , 1995 , 164, 404-13	7	22
14	Synergistic effects of cytokine and hyperthermia on cytotoxicity in HT-29 cells are not mediated by alteration of induced protein levels. <i>Journal of Cellular Physiology</i> , 1993 , 155, 27-35	7	15
13	Effect of thermotolerance on heat-induced excess nuclear-associated proteins. <i>Journal of Cellular Physiology</i> , 1993 , 156, 171-81	7	9
12	Expression, synthesis, and phosphorylation of HSP28 family during development and decay of thermotolerance in CHO plateau-phase cells. <i>Journal of Cellular Physiology</i> , 1992 , 150, 441-6	7	12
11	Heat-resistant variants of the Chinese hamster ovary cell: alteration of cellular structure and expression of vimentin. <i>Journal of Cellular Physiology</i> , 1992 , 151, 138-46	7	13
10	Development of acute thermotolerance in L929 cells: lack of HSP28 synthesis and phosphorylation. <i>Journal of Cellular Physiology</i> , 1992 , 152, 118-25	7	20
9	Constitutive HSP70: oligomerization and its dependence on ATP binding. <i>Journal of Cellular Physiology</i> , 1992 , 153, 353-61	7	42
8	Differences in preferential synthesis and redistribution of HSP70 and HSP28 families by heat or sodium arsenite in Chinese hamster ovary cells. <i>Journal of Cellular Physiology</i> , 1991 , 149, 77-87	7	23

7	Effect of tunicamycin on glycosylation of a 50 kDa protein and thermotolerance development. <i>Journal of Cellular Physiology</i> , 1991 , 149, 202-7	7	4
6	Inhibition of protein synthesis and heat protection: histidinol-resistant mutant cell lines. <i>Journal of Cellular Physiology</i> , 1991 , 149, 396-402	7	5
5	Effect of histidine on histidinol-induced heat protection in Chinese hamster ovary cells. <i>Journal of Cellular Physiology</i> , 1990 , 144, 401-7	7	16
4	Correlation between redistribution of a 26 kDa protein and development of chronic thermotolerance in various mammalian cell lines. <i>Journal of Cellular Physiology</i> , 1990 , 145, 324-32	7	12
3	Heat protectors and heat-induced preferential redistribution of 26 and 70 kDa proteins in Chinese hamster ovary cells. <i>Journal of Cellular Physiology</i> , 1989 , 141, 510-6	7	11
2	Protection of Chinese Hamster Ovary Cells from Hyperthermic Killing by Cycloheximide or Puromycin. <i>Radiation Research</i> , 1986 , 106, 98	3.1	48
1	Low glucose-enhanced TRAIL cytotoxicity is mediated through the ceramide/Akt/JIP pathway		1