In Kook Chun

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4380116/publications.pdf

Version: 2024-02-01

933447 940533 18 572 10 16 citations h-index g-index papers 18 18 18 1192 citing authors docs citations times ranked all docs

#	Article	IF	CITATIONS
1	Total lesion glycolysis in positron emission tomography is a better predictor of outcome than the International Prognostic Index for patients with diffuse large B cell lymphoma. Cancer, 2013, 119, 1195-1202.	4.1	136
2	Oligomeric forms of amyloid-β protein in plasma as a potential blood-based biomarker for Alzheimer's disease. Alzheimer's Research and Therapy, 2017, 9, 98.	6.2	108
3	Imaging sensitivity of dedicated positron emission mammography in relation to tumor size. Breast, 2012, 21, 66-71.	2.2	72
4	Comparison of 4D CT, Ultrasonography, and ^{99m} Tc Sestamibi SPECT/CT in Localizing Singleâ€Gland Primary Hyperparathyroidism. Otolaryngology - Head and Neck Surgery, 2015, 152, 438-443.	1.9	72
5	Correlation of breast cancer subtypes, based on estrogen receptor, progesterone receptor, and HER2, with functional imaging parameters from 68Ga-RGD PET/CT and 18F-FDG PET/CT. European Journal of Nuclear Medicine and Molecular Imaging, 2014, 41, 1534-1543.	6.4	65
6	Detection and Characterization of Parathyroid Adenoma/Hyperplasia for Preoperative Localization: Comparison Between 11C-Methionine PET/CT and 99mTc-Sestamibi Scintigraphy. Nuclear Medicine and Molecular Imaging, 2013, 47, 166-172.	1.0	24
7	Background 18F-FDG uptake in positron emission mammography (PEM): Correlation with mammographic density and background parenchymal enhancement in breast MRI. European Journal of Radiology, 2013, 82, 1738-1742.	2.6	24
8	Analysis of Cerebrospinal Fluid and [11C]PIB PET Biomarkers for Alzheimer's Disease with Updated Protocols. Journal of Alzheimer's Disease, 2016, 52, 1403-1413.	2.6	17
9	Early prediction of response to neoadjuvant chemotherapy in breast cancer patients: comparison of single-voxel 1H-magnetic resonance spectroscopy and 18F-fluorodeoxyglucose positron emission tomography. European Radiology, 2016, 26, 2279-2290.	4.5	14
10	Longitudinal Cerebral Perfusion Change in Transient Global Amnesia Related to Left Posterior Medial Network Disruption. PLoS ONE, 2015, 10, e0145658.	2.5	12
11	¹¹ C-PIB PET imaging reveals that amyloid deposition in cases with early-onset Alzheimer's disease in the absence of known mutations retains higher levels of PIB in the basal ganglia. Clinical Interventions in Aging, 2017, Volume 12, 1041-1048.	2.9	9
12	Pulmonary Artery Sarcoma Detected on F-18 FDG PET/CT as Origin of Multiple Spinal Metastases. Clinical Nuclear Medicine, 2011, 36, e87-e89.	1.3	6
13	Decreased Metabolism in the Posterior Medial Network with Concomitantly Increased Metabolism in the Anterior Temporal Network During Transient Global Amnesia. Brain Topography, 2018, 31, 468-476.	1.8	5
14	Clinical Significance of the Circle of Willis in Patients with Symptomatic Internal Carotid Artery Occlusion. World Neurosurgery, 2018, 115, e585-e591.	1.3	5
15	A Case of Enterocutaneous Fistula Diagnosed with Tc-99m DTPA Fistulography Using Hybrid SPECT/CT. Nuclear Medicine and Molecular Imaging, 2012, 46, 111-114.	1.0	1
16	Photo-Guided Sentinel Node Mapping in Breast Cancer Using Marker-Free Photo-Gamma Fusion Lymphoscintigraphy. Nuclear Medicine and Molecular Imaging, 2013, 47, 9-17.	1.0	1
17	P2-185: IMPROVED QUANTIFICATION METHODS FOR BRAIN \hat{l}^2 -AMYLOID BURDEN ON 11C-PIB PET IN PATIENTS WITH ALZHEIMER'S DISEASE. , 2014, 10, P538-P538.		1
18	Incidentally Found Soft Tissue 99mTc-DPD Uptake on Bone Scintigraphy Was Useful in an Early Diagnosis of Peripheral Arterial Disease. Nuclear Medicine and Molecular Imaging, 0, , .	1.0	0