

In Kook Chun

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4380116/publications.pdf>

Version: 2024-02-01

18
papers

572
citations

933447

10
h-index

940533

16
g-index

18
all docs

18
docs citations

18
times ranked

1192
citing authors

#	ARTICLE	IF	CITATIONS
1	Decreased Metabolism in the Posterior Medial Network with Concomitantly Increased Metabolism in the Anterior Temporal Network During Transient Global Amnesia. <i>Brain Topography</i> , 2018, 31, 468-476.	1.8	5
2	Clinical Significance of the Circle of Willis in Patients with Symptomatic Internal Carotid Artery Occlusion. <i>World Neurosurgery</i> , 2018, 115, e585-e591.	1.3	5
3	¹¹ C-PIB PET imaging reveals that amyloid deposition in cases with early-onset Alzheimer's disease in the absence of known mutations retains higher levels of PIB in the basal ganglia. <i>Clinical Interventions in Aging</i> , 2017, Volume 12, 1041-1048.	2.9	9
4	Oligomeric forms of amyloid- β protein in plasma as a potential blood-based biomarker for Alzheimer's disease. <i>Alzheimer's Research and Therapy</i> , 2017, 9, 98.	6.2	108
5	Analysis of Cerebrospinal Fluid and [11C]PIB PET Biomarkers for Alzheimer's Disease with Updated Protocols. <i>Journal of Alzheimer's Disease</i> , 2016, 52, 1403-1413.	2.6	17
6	Early prediction of response to neoadjuvant chemotherapy in breast cancer patients: comparison of single-voxel 1H-magnetic resonance spectroscopy and 18F-fluorodeoxyglucose positron emission tomography. <i>European Radiology</i> , 2016, 26, 2279-2290.	4.5	14
7	Comparison of 4D CT, Ultrasonography, and ^{99m} Tc Sestamibi SPECT/CT in Localizing Single Gland Primary Hyperparathyroidism. <i>Otolaryngology - Head and Neck Surgery</i> , 2015, 152, 438-443.	1.9	72
8	Longitudinal Cerebral Perfusion Change in Transient Global Amnesia Related to Left Posterior Medial Network Disruption. <i>PLoS ONE</i> , 2015, 10, e0145658.	2.5	12
9	P2-185: IMPROVED QUANTIFICATION METHODS FOR BRAIN β -AMYLOID BURDEN ON 11C-PIB PET IN PATIENTS WITH ALZHEIMER'S DISEASE. , 2014, 10, P538-P538.		1
10	Correlation of breast cancer subtypes, based on estrogen receptor, progesterone receptor, and HER2, with functional imaging parameters from 68Ga-RGD PET/CT and 18F-FDG PET/CT. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2014, 41, 1534-1543.	6.4	65
11	Detection and Characterization of Parathyroid Adenoma/Hyperplasia for Preoperative Localization: Comparison Between 11C-Methionine PET/CT and 99mTc-Sestamibi Scintigraphy. <i>Nuclear Medicine and Molecular Imaging</i> , 2013, 47, 166-172.	1.0	24
12	Photo-Guided Sentinel Node Mapping in Breast Cancer Using Marker-Free Photo-Gamma Fusion Lymphoscintigraphy. <i>Nuclear Medicine and Molecular Imaging</i> , 2013, 47, 9-17.	1.0	1
13	Total lesion glycolysis in positron emission tomography is a better predictor of outcome than the International Prognostic Index for patients with diffuse large B cell lymphoma. <i>Cancer</i> , 2013, 119, 1195-1202.	4.1	136
14	Background 18F-FDG uptake in positron emission mammography (PEM): Correlation with mammographic density and background parenchymal enhancement in breast MRI. <i>European Journal of Radiology</i> , 2013, 82, 1738-1742.	2.6	24
15	A Case of Enterocutaneous Fistula Diagnosed with Tc-99m DTPA Fistulography Using Hybrid SPECT/CT. <i>Nuclear Medicine and Molecular Imaging</i> , 2012, 46, 111-114.	1.0	1
16	Imaging sensitivity of dedicated positron emission mammography in relation to tumor size. <i>Breast</i> , 2012, 21, 66-71.	2.2	72
17	Pulmonary Artery Sarcoma Detected on F-18 FDG PET/CT as Origin of Multiple Spinal Metastases. <i>Clinical Nuclear Medicine</i> , 2011, 36, e87-e89.	1.3	6
18	Incidentally Found Soft Tissue 99mTc-DPD Uptake on Bone Scintigraphy Was Useful in an Early Diagnosis of Peripheral Arterial Disease. <i>Nuclear Medicine and Molecular Imaging</i> , 0, , .	1.0	0