

# Naoyuki Kawao

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4382148/publications.pdf>

Version: 2024-02-01

93  
papers

2,187  
citations

249298

26  
h-index

312153

41  
g-index

93  
all docs

93  
docs citations

93  
times ranked

2333  
citing authors

#	ARTICLE	IF	CITATIONS
1	Renal failure suppresses muscle irisin expression, and irisin blunts cortical bone loss in mice. <i>Journal of Cachexia, Sarcopenia and Muscle</i> , 2022, 13, 758-771.	2.9	18
2	Role of Macrophages and Plasminogen Activator Inhibitor-1 in Delayed Bone Repair Induced by Glucocorticoids in Mice. <i>International Journal of Molecular Sciences</i> , 2022, 23, 478.	1.8	3
3	Role of peripheral myelin protein 22 in chronic exercise-induced interactions of muscle and bone in mice. <i>Journal of Cellular Physiology</i> , 2022, 237, 2492-2502.	2.0	3
4	MicroRNA-196a-5p in Extracellular Vesicles Secreted from Myoblasts Suppresses Osteoclast-like Cell Formation in Mouse Cells. <i>Calcified Tissue International</i> , 2021, 108, 364-376.	1.5	24
5	Role of irisin in effects of chronic exercise on muscle and bone in ovariectomized mice. <i>Journal of Bone and Mineral Metabolism</i> , 2021, 39, 547-557.	1.3	14
6	Influence of Angptl1 on osteoclast formation and osteoblastic phenotype in mouse cells. <i>BMC Musculoskeletal Disorders</i> , 2021, 22, 398.	0.8	2
7	Effects of fluid flow shear stress to mouse muscle cells on the bone actions of muscle cell-derived extracellular vesicles. <i>PLoS ONE</i> , 2021, 16, e0250741.	1.1	12
8	Serpinb1a suppresses osteoclast formation. <i>Biochemistry and Biophysics Reports</i> , 2021, 26, 101004.	0.7	3
9	Role of plasminogen activator inhibitor-1 in muscle wasting induced by a diabetic state in female mice. <i>Endocrine Journal</i> , 2021, 68, 1421-1428.	0.7	1
10	Role of Dkk2 in the Muscle/bone Interaction of Androgen-Deficient Mice. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2021, 129, 770-775.	0.6	3
11	Role of tissue factor in delayed bone repair induced by diabetic state in mice. <i>PLoS ONE</i> , 2021, 16, e0260754.	1.1	1
12	Role of irisin in androgen-deficient muscle wasting and osteopenia in mice. <i>Journal of Bone and Mineral Metabolism</i> , 2020, 38, 161-171.	1.3	25
13	Myonectin inhibits the differentiation of osteoblasts and osteoclasts in mouse cells. <i>Heliyon</i> , 2020, 6, e03967.	1.4	3
14	Roles of Olfactomedin 1 in Muscle and Bone Alterations Induced by Gravity Change in Mice. <i>Calcified Tissue International</i> , 2020, 107, 180-190.	1.5	10
15	PAI-1 is involved in delayed bone repair induced by glucocorticoids in mice. <i>Bone</i> , 2020, 134, 115310.	1.4	11
16	Extracellular vesicles secreted from mouse muscle cells suppress osteoclast formation: Roles of mitochondrial energy metabolism. <i>Bone</i> , 2020, 134, 115298.	1.4	28
17	Roles of Dkk2 in the Linkage from Muscle to Bone during Mechanical Unloading in Mice. <i>International Journal of Molecular Sciences</i> , 2020, 21, 2547.	1.8	17
18	Roles of the vestibular system in obesity and impaired glucose metabolism in high-fat diet-fed mice. <i>PLoS ONE</i> , 2020, 15, e0228685.	1.1	7

#	ARTICLE	IF	CITATIONS
19	Roles of leptin in the recovery of muscle and bone by reloading after mechanical unloading in high fat diet-fed obese mice. <i>PLoS ONE</i> , 2019, 14, e0224403.	1.1	6
20	Plasminogen activator inhibitor-1 is involved in interleukin-1 $\beta$ -induced matrix metalloproteinase expression in murine chondrocytes. <i>Modern Rheumatology</i> , 2019, 29, 959-963.	0.9	3
21	Plasminogen activator inhibitor-1 deficiency suppresses osteoblastic differentiation of mesenchymal stem cells in mice. <i>Journal of Cellular Physiology</i> , 2019, 234, 9687-9697.	2.0	17
22	Roles of Irisin in the Linkage from Muscle to Bone During Mechanical Unloading in Mice. <i>Calcified Tissue International</i> , 2018, 103, 24-34.	1.5	50
23	Role of Macrophages and Plasminogen Activator Inhibitor-1 in Delayed Bone Repair in Diabetic Female Mice. <i>Endocrinology</i> , 2018, 159, 1875-1885.	1.4	15
24	Role of follistatin in muscle and bone alterations induced by gravity change in mice. <i>Journal of Cellular Physiology</i> , 2018, 233, 1191-1201.	2.0	35
25	Role of plasminogen activator inhibitor-1 in glucocorticoid-induced muscle change in mice. <i>Journal of Bone and Mineral Metabolism</i> , 2018, 36, 148-156.	1.3	18
26	Effects of hypergravity on gene levels in anti-gravity muscle and bone through the vestibular system in mice. <i>Journal of Physiological Sciences</i> , 2018, 68, 609-616.	0.9	6
27	Serpina3n, dominantly expressed in female osteoblasts, suppresses the phenotypes of differentiated osteoblasts in mice. <i>Endocrinology</i> , 2018, 159, 3775-3790.	1.4	15
28	Roles of plasminogen in the alterations in bone marrow hematopoietic stem cells during bone repair. <i>Bone Reports</i> , 2018, 8, 195-203.	0.2	4
29	A synthetic peptide derived from staphylokinase enhances FGF-2-induced skin wound healing in mice. <i>Thrombosis Research</i> , 2017, 157, 7-8.	0.8	2
30	Vitamin D deficiency aggravates diabetes-induced muscle wasting in female mice. <i>Diabetology International</i> , 2017, 8, 52-58.	0.7	11
31	Plasminogen activator inhibitor-1 deficiency enhances subchondral osteopenia after induction of osteoarthritis in mice. <i>BMC Musculoskeletal Disorders</i> , 2017, 18, 392.	0.8	12
32	Tissue fibrinolytic system in bone metabolism. <i>Japanese Journal of Thrombosis and Hemostasis</i> , 2017, 28, 597-602.	0.1	0
33	The vestibular system is critical for the changes in muscle and bone induced by hypergravity in mice. <i>Physiological Reports</i> , 2016, 4, e12979.	0.7	28
34	Novel roles of FKBP5 in muscle alteration induced by gravity change in mice. <i>Biochemical and Biophysical Research Communications</i> , 2016, 479, 602-606.	1.0	20
35	Stromal cell-derived factor-1 mediates changes of bone marrow stem cells during the bone repair process. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2016, 310, E15-E23.	1.8	29
36	Role of osteoclasts in heterotopic ossification enhanced by fibrodysplasia ossificans progressiva-related activin-like kinase 2 mutation in mice. <i>Journal of Bone and Mineral Metabolism</i> , 2016, 34, 517-525.	1.3	9

#	ARTICLE	IF	CITATIONS
37	The Tissue Fibrinolytic System Contributes to the Induction of Macrophage Function and CCL3 during Bone Repair in Mice. <i>PLoS ONE</i> , 2015, 10, e0123982.	1.1	22
38	Role of Plasminogen Activator Inhibitor-1 in Glucocorticoid-Induced Diabetes and Osteopenia in Mice. <i>Diabetes</i> , 2015, 64, 2194-2206.	0.3	55
39	$\hat{\pm}2$ -Antiplasmin is involved in bone loss induced by ovariectomy in mice. <i>Bone</i> , 2015, 79, 233-241.	1.4	15
40	Interactions Between Muscle Tissues and Bone Metabolism. <i>Journal of Cellular Biochemistry</i> , 2015, 116, 687-695.	1.2	176
41	Plasminogen Activator Inhibitor-1 Is Involved in Impaired Bone Repair Associated with Diabetes in Female Mice. <i>PLoS ONE</i> , 2014, 9, e92686.	1.1	46
42	Fibrodysplasia Ossificans Progressiva-related Activated Activin-like Kinase Signaling Enhances Osteoclast Formation during Heterotopic Ossification in Muscle Tissues. <i>Journal of Biological Chemistry</i> , 2014, 289, 16966-16977.	1.6	26
43	Tissue-type plasminogen activator deficiency delays bone repair: roles of osteoblastic proliferation and vascular endothelial growth factor. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2014, 307, E278-E288.	1.8	31
44	Plasminogen Activator Inhibitor-1 Deficiency Ameliorates Insulin Resistance and Hyperlipidemia But Not Bone Loss in Obese Female Mice. <i>Endocrinology</i> , 2014, 155, 1708-1717.	1.4	29
45	Enhanced pre-operative thrombolytic status is associated with the incidence of deep venous thrombosis in patients undergoing total knee arthroplasty. <i>Thrombosis Journal</i> , 2014, 12, 11.	0.9	1
46	Influence of diabetic state and vitamin D deficiency on bone repair in female mice. <i>Bone</i> , 2014, 61, 102-108.	1.4	23
47	Plasminogen Activator Inhibitor-1 Is Involved in Streptozotocin-Induced Bone Loss in Female Mice. <i>Diabetes</i> , 2013, 62, 3170-3179.	0.3	46
48	Lack of both $\hat{\pm}2$ -antiplasmin and plasminogen activator inhibitor type-1 induces high IgE production. <i>Life Sciences</i> , 2013, 93, 89-95.	2.0	9
49	Enzamin ameliorates adipose tissue inflammation with impaired adipocytokine expression and insulin resistance in db/db mice. <i>Journal of Nutritional Science</i> , 2013, 2, e37.	0.7	4
50	Plasminogen Plays a Crucial Role in Bone Repair. <i>Journal of Bone and Mineral Research</i> , 2013, 28, 1561-1574.	3.1	62
51	Role of matrix metalloproteinase-10 in the BMP-2 inducing osteoblastic differentiation. <i>Endocrine Journal</i> , 2013, 60, 1309-1319.	0.7	20
52	Role of the fibrinolytic system in recovery responses after liver injury. <i>Japanese Journal of Thrombosis and Hemostasis</i> , 2013, 24, 501-506.	0.1	0
53	In Vivo Diagnostic Imaging Using Micro-CT: Sequential and Comparative Evaluation of Rodent Models for Hepatic/Brain Ischemia and Stroke. <i>PLoS ONE</i> , 2012, 7, e32342.	1.1	22
54	Urokinase-type plasminogen activator and plasminogen mediate activation of macrophage phagocytosis during liver repair in vivo. <i>Thrombosis and Haemostasis</i> , 2012, 107, 749-759.	1.8	16

#	ARTICLE	IF	CITATIONS
55	Urokinase-type plasminogen activator contributes to heterogeneity of macrophages at the border of damaged site during liver repair in mice. <i>Thrombosis and Haemostasis</i> , 2011, 105, 892-900.	1.8	15
56	Spatiotemporal differences in vascular permeability after ischaemic brain damage. <i>NeuroReport</i> , 2011, 22, 424-427.	0.6	3
57	Profibrinolytic effect of Enzamin, an extract of metabolic products from <i>Bacillus subtilis</i> AK and <i>Lactobacillus</i> . <i>Journal of Thrombosis and Thrombolysis</i> , 2011, 32, 195-200.	1.0	6
58	Systemic transplantation of embryonic stem cells accelerates brain lesion decrease and angiogenesis. <i>NeuroReport</i> , 2010, 21, 575-579.	0.6	26
59	Enhancement of fibrinolytic activity in vascular endothelial cells by heterologous expression of adenine nucleotide translocase-1. <i>Blood Coagulation and Fibrinolysis</i> , 2010, 21, 272-278.	0.5	0
60	Initial brain lesion size affects the extent of subsequent pathophysiological responses. <i>Brain Research</i> , 2010, 1322, 109-117.	1.1	19
61	Plasminogen is essential for granulation tissue formation during the recovery process after liver injury in mice. <i>Journal of Thrombosis and Haemostasis</i> , 2010, 8, 1555-1566.	1.9	12
62	Role of plasminogen in macrophage accumulation during liver repair. <i>Thrombosis Research</i> , 2010, 125, e214-e221.	0.8	15
63	Urokinase-type plasminogen activator receptor (uPAR) augments brain damage in a murine model of ischemic stroke. <i>Neuroscience Letters</i> , 2008, 432, 46-49.	1.0	25
64	Binding of plasminogen to hepatocytes isolated from injured mice liver and nonparenchymal cell-dependent proliferation of hepatocytes. <i>Blood Coagulation and Fibrinolysis</i> , 2008, 19, 503-511.	0.5	8
65	Effect of staphylokinase-derived nonadecapeptide on the activation of plasminogen. <i>Thrombosis and Haemostasis</i> , 2007, 97, 795-802.	1.8	6
66	Plasmin decreases the BH3-only protein BimEL via the ERK1/2 signaling pathway in hepatocytes. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2007, 1773, 718-727.	1.9	11
67	Colonic hyperalgesia triggered by proteinase-activated receptor-2 in mice: Involvement of endogenous bradykinin. <i>Neuroscience Letters</i> , 2006, 402, 167-172.	1.0	31
68	Antiallodynic effect of etidronate, a bisphosphonate, in rats with adjuvant-induced arthritis: Involvement of ATP-sensitive K <sup>+</sup> channels. <i>Neuropharmacology</i> , 2006, 51, 182-190.	2.0	18
69	Suppression of pancreatitis-related allodynia/hyperalgesia by proteinase-activated receptor-2 in mice. <i>British Journal of Pharmacology</i> , 2006, 148, 54-60.	2.7	47
70	Physiology and Pathophysiology of Proteinase-Activated Receptors (PARs): PARs in the Respiratory System: Cellular Signaling and Physiological/Pathological Roles. <i>Journal of Pharmacological Sciences</i> , 2005, 97, 20-24.	1.1	60
71	Signal Transduction for Proteinase-Activated Receptor-2-Triggered Prostaglandin E <sub>2</sub> Formation in Human Lung Epithelial Cells. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2005, 315, 576-589.	1.3	49
72	Proteinase-Activated Receptor-2-Mediated Relaxation in Mouse Tracheal and Bronchial Smooth Muscle: Signal Transduction Mechanisms and Distinct Agonist Sensitivity. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2004, 311, 402-410.	1.3	37

#	ARTICLE	IF	CITATIONS
73	Distinct roles for protease-activated receptors 1 and 2 in vasomotor modulation in rat superior mesenteric artery. <i>Cardiovascular Research</i> , 2004, 61, 683-692.	1.8	25
74	A protective role of protease-activated receptor 1 in rat gastric mucosa. <i>Gastroenterology</i> , 2004, 126, 208-219.	0.6	45
75	The potent inducible nitric oxide synthase inhibitor ONO-1714 inhibits neuronal NOS and exerts antinociception in rats. <i>Neuroscience Letters</i> , 2004, 365, 111-115.	1.0	19
76	Receptor-activating peptides for PAR-1 and PAR-2 relax rat gastric artery via multiple mechanisms. <i>Life Sciences</i> , 2004, 75, 2689-2702.	2.0	16
77	Activation of trigeminal nociceptive neurons by parotid PAR-2 activation in rats. <i>NeuroReport</i> , 2004, 15, 1617-1621.	0.6	17
78	Modulation of Capsaicin-Evoked Visceral Pain and Referred Hyperalgesia by Protease-Activated Receptors 1 and 2. <i>Journal of Pharmacological Sciences</i> , 2004, 94, 277-285.	1.1	58
79	Effect of a potent iNOS inhibitor (ONO-1714) on acetaminophen-induced hepatotoxicity in the rat. <i>Life Sciences</i> , 2003, 74, 793-802.	2.0	35
80	The PAR-1-activating peptide facilitates pepsinogen secretion in rats. <i>Peptides</i> , 2003, 24, 1449-1451.	1.2	13
81	Capsazepine Inhibits Thermal Hyperalgesia but Not Nociception Triggered by Protease-Activated Receptor-2 in Rats. <i>The Japanese Journal of Pharmacology</i> , 2002, 89, 184-187.	1.2	28
82	Specific expression of spinal Fos after PAR-2 stimulation in mast cell-depleted rats. <i>NeuroReport</i> , 2002, 13, 511-514.	0.6	24
83	Protease-activated receptor-2 (PAR-2) in the pancreas and parotid gland: Immunolocalization and involvement of nitric oxide in the evoked amylase secretion. <i>Life Sciences</i> , 2002, 71, 2435-2446.	2.0	64
84	The PAR-1-activating peptide attenuates carrageenan-induced hyperalgesia in rats. <i>Peptides</i> , 2002, 23, 1181-1183.	1.2	36
85	Role of N-methyl-d-aspartate receptors and the nitric oxide pathway in nociception/hyperalgesia elicited by protease-activated receptor-2 activation in mice and rats. <i>Neuroscience Letters</i> , 2002, 329, 349-353.	1.0	25
86	Protease-activated receptor-2 (PAR-2) in the rat gastric mucosa: immunolocalization and facilitation of pepsin/pepsinogen secretion. <i>British Journal of Pharmacology</i> , 2002, 135, 1292-1296.	2.7	51
87	Factor Xa-Evoked Relaxation in Rat Aorta: Involvement of PAR-2. <i>Biochemical and Biophysical Research Communications</i> , 2001, 282, 432-435.	1.0	48
88	Ex Vivo Evidence That the Phosphodiesterase Inhibitor IBMX Attenuates the Up-Regulation of PAR-2 in the Endotoxemic Rat Aorta. <i>Thrombosis Research</i> , 2001, 101, 513-515.	0.8	7
89	Peripheral PAR-2 triggers thermal hyperalgesia and nociceptive responses in rats. <i>NeuroReport</i> , 2001, 12, 715-719.	0.6	94
90	Lipopolysaccharide-induced subsensitivity of protease-activated receptor-2 in the mouse salivary glands in vivo. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 2001, 364, 281-284.	1.4	14

#	ARTICLE	IF	CITATIONS
91	In vivo evidence that protease-activated receptors 1 and 2 modulate gastrointestinal transit in the mouse. <i>British Journal of Pharmacology</i> , 2001, 133, 1213-1218.	2.7	71
92	Secondary somatosensory cortex stimulation facilitates the antinociceptive effect of the NO synthase inhibitor through suppression of spinal nociceptive neurons in the rat. <i>Brain Research</i> , 2001, 903, 110-116.	1.1	20
93	Somatosensory cortex stimulation-evoked analgesia in rats: Potentiation by no synthase inhibition. <i>Life Sciences</i> , 2000, 66, PL271-PL276.	2.0	21