

# Miki Imanishi

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4393345/publications.pdf>

Version: 2024-02-01

65  
papers

1,213  
citations

394421

19  
h-index

434195

31  
g-index

73  
all docs

73  
docs citations

73  
times ranked

1353  
citing authors

#	ARTICLE	IF	CITATIONS
1	Artificial Nanocage Formed via Self-Assembly of $\beta^2$ -Annulus Peptide for Delivering Biofunctional Proteins into Cell Interiors. <i>Bioconjugate Chemistry</i> , 2022, 33, 311-320.	3.6	9
2	L17ER4: A cell-permeable attenuated cationic amphiphilic lytic peptide. <i>Bioorganic and Medicinal Chemistry</i> , 2022, 61, 116728.	3.0	3
3	Grafting Hydrophobic Amino Acids Critical for Inhibition of Protein-Protein Interactions on a Cell-Penetrating Peptide Scaffold. <i>Molecular Pharmaceutics</i> , 2022, 19, 558-567.	4.6	3
4	Recognition of G-quadruplex RNA by a crucial RNA methyltransferase component, METTL14. <i>Nucleic Acids Research</i> , 2022, 50, 449-457.	14.5	21
5	Mechanisms and Strategies for Determining m <sup>6</sup> A RNA Modification Sites by Natural and Engineered m <sup>6</sup> A Effector Proteins. <i>Chemistry - an Asian Journal</i> , 2022, 17, .	3.3	3
6	Split luciferase-based estimation of cytosolic cargo concentration delivered intracellularly via attenuated cationic amphiphilic lytic peptides. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2022, 72, 128875.	2.2	0
7	Discovery of a Macropinocytosis-Inducing Peptide Potentiated by Medium-Mediated Intramolecular Disulfide Formation. <i>Angewandte Chemie - International Edition</i> , 2021, 60, 11928-11936.	13.8	11
8	Discovery of a Macropinocytosis-Inducing Peptide Potentiated by Medium-Mediated Intramolecular Disulfide Formation. <i>Angewandte Chemie</i> , 2021, 133, 12035-12043.	2.0	2
9	Use of homoarginine to obtain attenuated cationic membrane lytic peptides. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2021, 40, 127925.	2.2	7
10	Liquid Droplet Formation and Facile Cytosolic Translocation of IgG in the Presence of Attenuated Cationic Amphiphilic Lytic Peptides. <i>Angewandte Chemie</i> , 2021, 133, 19957-19965.	2.0	2
11	Titelbild: Liquid Droplet Formation and Facile Cytosolic Translocation of IgG in the Presence of Attenuated Cationic Amphiphilic Lytic Peptides ( <i>Angew. Chem.</i> 36/2021). <i>Angewandte Chemie</i> , 2021, 133, 19645-19645.	2.0	0
12	Liquid Droplet Formation and Facile Cytosolic Translocation of IgG in the Presence of Attenuated Cationic Amphiphilic Lytic Peptides. <i>Angewandte Chemie - International Edition</i> , 2021, 60, 19804-19812.	13.8	21
13	Programmable RNA methylation and demethylation using PUF RNA binding proteins. <i>Chemical Communications</i> , 2020, 56, 1365-1368.	4.1	23
14	Conversion of cationic amphiphilic lytic peptides to cell-penetration peptides. <i>Peptide Science</i> , 2020, 112, e24144.	1.8	11
15	Identification of synthetic inhibitors for the DNA binding of intrinsically disordered circadian clock transcription factors. <i>Chemical Communications</i> , 2020, 56, 11203-11206.	4.1	5
16	Effective RNA Regulation by Combination of Multiple Programmable RNA-Binding Proteins. <i>Applied Sciences (Switzerland)</i> , 2020, 10, 6803.	2.5	3
17	Optimizing Charge Switching in Membrane Lytic Peptides for Endosomal Release of Biomacromolecules. <i>Angewandte Chemie</i> , 2020, 132, 20165-20173.	2.0	6
18	Optimizing Charge Switching in Membrane Lytic Peptides for Endosomal Release of Biomacromolecules. <i>Angewandte Chemie - International Edition</i> , 2020, 59, 19990-19998.	13.8	36

#	ARTICLE	IF	CITATIONS
19	Stimulating Macropinocytosis for Intracellular Nucleic Acid and Protein Delivery: A Combined Strategy with Membrane-Lytic Peptides To Facilitate Endosomal Escape. <i>Bioconjugate Chemistry</i> , 2020, 31, 547-553.	3.6	31
20	Enhancing the activity of membrane remodeling epsin-peptide by trimerization. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2020, 30, 127190.	2.2	12
21	Modified nucleobase-specific gene regulation using engineered transcription activator-like effectors. <i>Advanced Drug Delivery Reviews</i> , 2019, 147, 59-65.	13.7	6
22	Loosening of Lipid Packing by Cell Surface Recruitment of Amphiphilic Peptides by Coiled-Coil Tethering. <i>ChemBioChem</i> , 2019, 20, 2151-2159.	2.6	5
23	Nested PUF Proteins: Extending Target RNA Elements for Gene Regulation. <i>ChemBioChem</i> , 2018, 19, 171-176.	2.6	6
24	Sequence-specific 5mC detection in live cells based on the TALE-split luciferase complementation system. <i>Analyst</i> , 2018, 143, 3793-3797.	3.5	2
25	Detection of N <sup>6</sup> -methyladenosine based on the methyl-sensitivity of MazF RNA endonuclease. <i>Chemical Communications</i> , 2017, 53, 12930-12933.	4.1	113
26	Calmodulin EF-hand peptides as Ca <sup>2+</sup> -switchable recognition tags. <i>Biopolymers</i> , 2017, 108, e22937.	2.4	1
27	Preparation of peptide thioesters from naturally occurring sequences using reaction sequence consisting of regioselective S-cyanylation and hydrazinolysis. <i>Biopolymers</i> , 2016, 106, 531-546.	2.4	16
28	Sequence-specific recognition of methylated DNA by an engineered transcription activator-like effector protein. <i>Chemical Communications</i> , 2016, 52, 14238-14241.	4.1	13
29	Controlling leucine-zipper partner recognition in cells through modification of a <sup>g</sup> interactions. <i>Chemical Communications</i> , 2014, 50, 6364-6367.	4.1	8
30	Creating a TALE protein with unbiased 5 <sup>2</sup> -T binding. <i>Biochemical and Biophysical Research Communications</i> , 2013, 441, 262-265.	2.1	21
31	Identification of cellular proteins interacting with octaarginine (R8) cell-penetrating peptide by photo-crosslinking. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2013, 23, 3738-3740.	2.2	20
32	Construction of a Ca <sup>2+</sup> -Gated Artificial Channel by Fusing Alamethicin with a Calmodulin-Derived Extramembrane Segment. <i>Bioconjugate Chemistry</i> , 2013, 24, 188-195.	3.6	5
33	Dipicolylamine as a unique structural switching element for helical peptides. <i>Organic and Biomolecular Chemistry</i> , 2012, 10, 6062.	2.8	10
34	Construction of a Rhythm Transfer System That Mimics the Cellular Clock. <i>ACS Chemical Biology</i> , 2012, 7, 1817-1821.	3.4	5
35	Zn(II) Binding and DNA Binding Properties of Ligand-Substituted CXHH-Type Zinc Finger Proteins. <i>Biochemistry</i> , 2012, 51, 3342-3348.	2.5	21
36	Signal Transduction Using an Artificial Receptor System that Undergoes Dimerization Upon Addition of a Bivalent Leucine-Zipper Ligand. <i>Angewandte Chemie - International Edition</i> , 2012, 51, 7464-7467.	13.8	39

#	ARTICLE	IF	CITATIONS
37	Rational design of DNA sequence-specific zinc fingers. <i>FEBS Letters</i> , 2012, 586, 918-923.	2.8	5
38	An Arginine Residue Instead of a Conserved Leucine Residue in the Recognition Helix of the Finger 3 of Zif268 Stabilizes the Domain Structure and Mediates DNA Binding. <i>Biochemistry</i> , 2011, 50, 6266-6272.	2.5	8
39	Control of Circadian Phase by an Artificial Zinc Finger Transcription Regulator. <i>Angewandte Chemie - International Edition</i> , 2011, 50, 9396-9399.	13.8	3
40	Octa-Arginine Mediated Delivery of Wild-Type Lnk Protein Inhibits TPO-Induced M-MOK Megakaryoblastic Leukemic Cell Growth by Promoting Apoptosis. <i>PLoS ONE</i> , 2011, 6, e23640.	2.5	31
41	Metal-stimulated Regulation of Transcription by an Artificial Zinc-Finger Protein. <i>ChemBioChem</i> , 2010, 11, 1653-1655.	2.6	10
42	Expressed protein ligation for the preparation of fusion proteins with cell penetrating peptides for endotoxin removal and intracellular delivery. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2010, 1798, 2249-2257.	2.6	15
43	Zinc finger-zinc finger interaction between the transcription factors, GATA-1 and Sp1. <i>Biochemical and Biophysical Research Communications</i> , 2010, 400, 625-630.	2.1	7
44	Non-FokI-Based Zinc Finger Nucleases. <i>Methods in Molecular Biology</i> , 2010, 649, 337-349.	0.9	7
45	Cobalt(II)-Responsive DNA Binding of a GCN4-bZIP Protein Containing Cysteine Residues Functionalized with Iminodiacetic Acid. <i>Angewandte Chemie - International Edition</i> , 2009, 48, 6853-6856.	13.8	20
46	Cytosolic Targeting of Macromolecules Using a pH-Dependent Fusogenic Peptide in Combination with Cationic Liposomes. <i>Bioconjugate Chemistry</i> , 2009, 20, 953-959.	3.6	81
47	Positive and negative cooperativity of modularly assembled zinc fingers. <i>Biochemical and Biophysical Research Communications</i> , 2009, 387, 440-443.	2.1	13
48	New Redesigned Zinc-Finger Proteins: Design Strategy and Its Application. <i>Chemistry - A European Journal</i> , 2008, 14, 3236-3249.	3.3	64
49	Effects of Bulkiness and Hydrophobicity of an Aliphatic Amino Acid in the Recognition Helix of the GAGA Zinc Finger on the Stability of the Hydrophobic Core and DNA Binding Affinity. <i>Biochemistry</i> , 2008, 47, 11717-11724.	2.5	6
50	Rapid Transcriptional Activity <i>in Vivo</i> and Slow DNA Binding <i>in Vitro</i> by an Artificial Multi-Zinc Finger Protein. <i>Biochemistry</i> , 2008, 47, 10171-10177.	2.5	12
51	DNA-Binding Ability of GAGA Zinc Finger Depends on the Nature of Amino Acids Present in the $\hat{1}^2$ -Hairpin. <i>Biochemistry</i> , 2007, 46, 7506-7513.	2.5	10
52	$\hat{1}^{\pm}$ -Helical Linker of an Artificial 6-Zinc Finger Peptide Contributes to Selective DNA Binding to a Discontinuous Recognition Sequence. <i>Biochemistry</i> , 2007, 46, 8517-8524.	2.5	24
53	Autoinhibition regulates the motility of the <i>C. elegans</i> intraflagellar transport motor OSM-3. <i>Journal of Cell Biology</i> , 2006, 174, 931-937.	5.2	105
54	Effects of linking 15-zinc finger domains on DNA binding specificity and multiple DNA binding modes. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2005, 15, 2197-2201.	2.2	10

#	ARTICLE	IF	CITATIONS
55	Site-specific DNA cleavage by artificial zinc finger-type nuclease with cerium-binding peptide. <i>Biochemical and Biophysical Research Communications</i> , 2005, 330, 247-252.	2.1	30
56	An artificial six-zinc finger peptide with polyarginine linker: Selective binding to the discontinuous DNA sequences. <i>Biochemical and Biophysical Research Communications</i> , 2005, 333, 167-173.	2.1	9
57	Swapping of the $\hat{I}^2$ -Hairpin Region between Sp1 and GLI Zinc Fingers: A Significant Role of the $\hat{I}^2$ -Hairpin Region in DNA Binding Properties of C2H2-type Zinc Finger Peptides. <i>Biochemistry</i> , 2005, 44, 2523-2528.	2.5	12
58	Exchange of Histidine Spacing between Sp1 and GLI Zinc Fingers: A Distinct Effect of Histidine Spacing-Linker Region on DNA Binding. <i>Biochemistry</i> , 2004, 43, 6352-6359.	2.5	7
59	Creation and characteristics of unnatural CysHis3-type zinc finger protein. <i>Biochemical and Biophysical Research Communications</i> , 2004, 325, 421-425.	2.1	13
60	DNA cleavage characteristics of non-protein enediyne antibiotic N1999A2. <i>Biochemical and Biophysical Research Communications</i> , 2003, 306, 87-92.	2.1	17
61	Artificial DNA-Bending Six-Zinc Finger Peptides with Different Charged Linkers: Distinct Kinetic Properties of DNA Bindings. <i>Biochemistry</i> , 2002, 41, 1328-1334.	2.5	31
62	Multiconnection of Identical Zinc Finger: Implication for DNA Binding Affinity and Unit Modulation of the Three Zinc Finger Domain. <i>Biochemistry</i> , 2001, 40, 2932-2941.	2.5	17
63	Design of novel zinc finger proteins: towards artificial control of specific gene expression. <i>European Journal of Pharmaceutical Sciences</i> , 2001, 13, 91-97.	4.0	25
64	DNA-Bending Finger: Artificial Design of 6-Zinc Finger Peptides with Polyglycine Linker and Induction of DNA Bending. <i>Biochemistry</i> , 2000, 39, 4383-4390.	2.5	37
65	Artificial Nine Zinc-Finger Peptide with 30 Base Pair Binding Sites. <i>Biochemistry</i> , 1998, 37, 13827-13834.	2.5	69