

Carlotta AbbÃ

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4394116/publications.pdf>

Version: 2024-02-01

13
papers

73
citations

1684188

5
h-index

1474206

9
g-index

13
all docs

13
docs citations

13
times ranked

94
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Clonal Megakaryocyte Dysplasia with Normal Blood Values Is a Distinct Myeloproliferative Neoplasm. <i>Acta Haematologica</i> , 2022, 145, 30-37. | 1.4 | 7 |
| 2 | Reduced CXCR4-expression on CD34-positive blood cells predicts outcomes of persons with primary myelofibrosis. <i>Leukemia</i> , 2021, 35, 468-475. | 7.2 | 7 |
| 3 | Clinical Relevance of VEGFA (rs3025039) +936 C>T Polymorphism in Primary Myelofibrosis: Susceptibility, Clinical Co-Variates, and Outcomes. <i>Genes</i> , 2021, 12, 1271. | 2.4 | 4 |
| 4 | Bone Marrow Microenvironment in Light-Chain Amyloidosis: In Vitro Expansion and Characterization of Mesenchymal Stromal Cells. <i>Biomedicines</i> , 2021, 9, 1523. | 3.2 | 0 |
| 5 | Primary myelofibrosis: rs2010963 VEGFA polymorphism favors a prefibrotic phenotype and is associated with higher risk of thrombosis. <i>Leukemia Research</i> , 2021, 111, 106730. | 0.8 | 3 |
| 6 | VEGFA rs3025020 Polymorphism Contributes to CALR-Mutation Susceptibility and Is Associated with Low Risk of Deep Vein Thrombosis in Primary Myelofibrosis. <i>TH Open</i> , 2021, 05, e513-e520. | 1.4 | 1 |
| 7 | Clinical Applications of Mesenchymal Stem/Stromal Cell Derived Extracellular Vesicles: Therapeutic Potential of an Acellular Product. <i>Diagnostics</i> , 2020, 10, 999. | 2.6 | 34 |
| 8 | Plasma sIL-2R β levels are associated with disease progression in myelofibrosis with JAK2V617F but not CALR mutation. <i>Leukemia Research</i> , 2020, 90, 106319. | 0.8 | 7 |
| 9 | Constitutive STAT5 phosphorylation in CD34+ cells of patients with primary myelofibrosis: Correlation with driver mutation status and disease severity. <i>PLoS ONE</i> , 2019, 14, e0220189. | 2.5 | 3 |
| 10 | Kinetic and Angiogenic Activity of Circulating Endothelial Colony Forming Cells in Patients with Infantile Haemangioma Receiving Propranolol. <i>Thrombosis and Haemostasis</i> , 2019, 119, 274-284. | 3.4 | 7 |
| 11 | Elevated Plasma sIL-2R β Levels in Primary Myelofibrosis Play a Distinct Role on Disease Progression in JAK2V617F and Calr Mutants. <i>Blood</i> , 2019, 134, 1678-1678. | 1.4 | 0 |
| 12 | Increased STAT5/STAT3 Intracellular Signaling in Circulating CD34+ Cells of Patients with PMF Correlates with Disease Severity. <i>Blood</i> , 2018, 132, 4337-4337. | 1.4 | 0 |
| 13 | Possible Role of Impaired Erk1,2 Phosphorylation and Increased sIL2r Alpha Plasma Levels in the Reduced Frequency of Circulating T Regulatory Cells of Patients with Primary Myelofibrosis. <i>Blood</i> , 2015, 126, 1639-1639. | 1.4 | 0 |