

Jinling Liao

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4395083/publications.pdf>

Version: 2024-02-01

24

papers

736

citations

623734

14

h-index

677142

22

g-index

24

all docs

24

docs citations

24

times ranked

534

citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | A novel nematode effector suppresses plant immunity by activating host reactive oxygen speciesâ€scavenging system. <i>New Phytologist</i> , 2016, 209, 1159-1173. | 7.3 | 148 |
| 2 | A novel <i>Meloidogyne graminicola</i> effector, MgGPP, is secreted into host cells and undergoes glycosylation in concert with proteolysis to suppress plant defenses and promote parasitism. <i>PLoS Pathogens</i> , 2017, 13, e1006301. | 4.7 | 90 |
| 3 | A novel <i>Meloidogyne enterolobii</i> effector MeTCTP promotes parasitism by suppressing programmed cell death in host plants. <i>Molecular Plant Pathology</i> , 2017, 18, 45-54. | 4.2 | 76 |
| 4 | A novel <i>Meloidogyne graminicola</i> effector, MgMO237, interacts with multiple host defenceâ€related proteins to manipulate plant basal immunity and promote parasitism. <i>Molecular Plant Pathology</i> , 2018, 19, 1942-1955. | 4.2 | 70 |
| 5 | A Novel Effector Protein, MJ-NULG1a, Targeted to Giant Cell Nuclei Plays a Role in<i>Meloidogyne javanica</i> Parasitism. <i>Molecular Plant-Microbe Interactions</i> , 2013, 26, 55-66. | 2.6 | 68 |
| 6 | Morphological and molecular characterization of <i>Aphelenchoides fujianensis</i> n. sp. (Nematoda:) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 54 | 0.5 | 38 |
| 7 | The Complete Mitochondrial Genome of <i>Meloidogyne graminicola</i> (Tylenchina): A Unique Gene Arrangement and Its Phylogenetic Implications. <i>PLoS ONE</i> , 2014, 9, e98558. | 2.5 | 31 |
| 8 | Molecular cloning and characterization of a calreticulin cDNA from the pinewood nematode <i>Bursaphelenchus xylophilus</i> . <i>Experimental Parasitology</i> , 2011, 128, 121-126. | 1.2 | 30 |
| 9 | Morphological and molecular charaterisation of <i>Pratylenchus parazeae</i> n. sp. (Nematoda:) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 50 | 1.7 | 26 |
| 10 | Loop-mediated isothermal amplification based on the mitochondrial COI region to detect <i>Pratylenchus zeae</i> . <i>European Journal of Plant Pathology</i> , 2017, 148, 435-446. | 1.7 | 22 |
| 11 | The <i>Meloidogyne javanica</i> effector Mj2G02 interferes with jasmonic acid signalling to suppress cell death and promote parasitism in <i>Arabidopsis</i>. <i>Molecular Plant Pathology</i> , 2021, 22, 1288-1301. | 4.2 | 22 |
| 12 | Improved dominant selection markers and co-culturing conditions for efficient Agrobacterium tumefaciens-mediated transformation of <i>Ustilago scitaminea</i> . <i>Biotechnology Letters</i> , 2014, 36, 1309-1314. | 2.2 | 20 |
| 13 | Molecular and biochemical characterization of the ß-1,4-endoglucanase gene Mj-eng-3 in the root-knot nematode <i>Meloidogyne javanica</i> . <i>Experimental Parasitology</i> , 2013, 135, 15-23. | 1.2 | 19 |
| 14 | Loop-Mediated Isothermal Amplification for the Detection of <i>Tylenchulus semipenetrans</i> in Soil. <i>Plant Disease</i> , 2016, 100, 877-883. | 1.4 | 18 |
| 15 | The <i>Meloidogyne graminicola</i> effector MgMO289 targets a novel copper metallochaperone to suppress immunity in rice. <i>Journal of Experimental Botany</i> , 2021, 72, 5638-5655. | 4.8 | 17 |
| 16 | <i>Meloidogyne aberrans</i> sp. nov. (Nematoda: Meloidogynidae), a new root-knot nematode parasitizing kiwifruit in China. <i>PLoS ONE</i> , 2017, 12, e0182627. | 2.5 | 16 |
| 17 | <i>Heterodera fengi</i> n. sp. (Nematoda: Heteroderinae) from bamboo in Guangdong Province, Chinaâ€”a new cyst nematode in the Cyperi group. <i>Zootaxa</i> , 2013, 3652, 179. | 0.5 | 9 |
| 18 | <i>Arabidopsis thaliana</i> as a model plant to study host- <i>Meloidogyne graminicola</i> interactions. <i>Nematology</i> , 2020, 22, 1015-1024. | 0.6 | 7 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Heterodera guangdongensis n. sp. (Nematoda: Heteroderinae) from bamboo in Guangdong Province, China—a new cyst nematode in the Cyperi group. Zootaxa, 2014, 3881, 488-500. | 0.5 | 5 |
| 20 | Duplex real-time quantitative PCR for simultaneous detection and quantification of Pratylenchus neglectus and P. thornei. European Journal of Plant Pathology, 2020, 157, 185-196. | 1.7 | 2 |
| 21 | <p class="HeadingRunIn">Morphological and molecular characterization of Zygotylenchus gansuensis n. sp. (Nematoda: Pratylenchinae) from China</p>. Zootaxa, 2014, 3821, 465. | 0.5 | 1 |
| 22 | Morphological and molecular characterisation of Paralongidorus sacchari (Nematoda:) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 627 Td (Longidoridae). Zootaxa, 2014, 3821, 465, 155, 615-625. | 1.7 | 1 |
| 23 | Transcriptome analysis of nematode-responsive genes in two susceptible Indica rice cultivars. Nematology, 2021, -1, 1-16. | 0.6 | 0 |
| 24 | The effector MJ-10A08 of Meloidogyne javanica is required for parasitism that suppressed programmed cell death in Nicotiana benthamiana. Nematology, 2022, 24, 1-14. | 0.6 | 0 |