

Joost C M Meijers

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4406003/publications.pdf>

Version: 2024-02-01

342
papers

15,234
citations

17776

65
h-index

31191

106
g-index

349
all docs

349
docs citations

349
times ranked

17227
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|------|-----------|
| 1 | Iron-Driven Alterations on Red Blood Cell-Derived Microvesicles Amplify Coagulation during Hemolysis via the Intrinsic Tenase Complex. <i>Thrombosis and Haemostasis</i> , 2022, 122, 080-091. | 1.8 | 3 |
| 2 | Major bleeding during oral anticoagulant therapy associated with factor V activation by factor Xa. <i>Journal of Thrombosis and Haemostasis</i> , 2022, 20, 328-338. | 1.9 | 4 |
| 3 | Coagulation in gout: is there a link with disease activity?. <i>Clinical Rheumatology</i> , 2022, 41, 1809-1815. | 1.0 | 1 |
| 4 | VhH anti- ϵ -thrombomodulin clone 1 inhibits TAFI activation and enhances fibrinolysis in human whole blood under flow. <i>Journal of Thrombosis and Haemostasis</i> , 2022, 20, 1213-1222. | 1.9 | 2 |
| 5 | Basic science research opportunities in thrombosis and hemostasis: Communication from the SSC of the ISTH. <i>Journal of Thrombosis and Haemostasis</i> , 2022, 20, 1496-1506. | 1.9 | 5 |
| 6 | A FRET-based assay for the quantitation of the thrombin-factor XI interaction. <i>Thrombosis Research</i> , 2022, 214, 23-28. | 0.8 | 3 |
| 7 | Reversing direct factor Xa or thrombin inhibitors: Factor V addition to prothrombin complex concentrate is beneficial in vitro. <i>Research and Practice in Thrombosis and Haemostasis</i> , 2022, 6, e12699. | 1.0 | 4 |
| 8 | Anti- ϵ -2- α -glycoprotein I and anti- ϵ -prothrombin antibodies cause lupus anticoagulant through different mechanisms of action. <i>Journal of Thrombosis and Haemostasis</i> , 2021, 19, 1018-1028. | 1.9 | 22 |
| 9 | Structure- ϵ -function of anticoagulant TIX-5, the inhibitor of factor Xa-mediated FV activation. <i>Journal of Thrombosis and Haemostasis</i> , 2021, 19, 1697-1708. | 1.9 | 1 |
| 10 | Induced normothermia ameliorates the procoagulant host response in human endotoxaemia. <i>British Journal of Anaesthesia</i> , 2021, 126, 1111-1118. | 1.5 | 3 |
| 11 | Iron-Driven Alterations on Red Blood Cell-Derived Microvesicles Amplify Coagulation during Hemolysis via the Intrinsic Tenase Complex. <i>Thrombosis and Haemostasis</i> , 2021, . . | 1.8 | 3 |
| 12 | Aprotinin Inhibits Thrombin Generation by Inhibition of the Intrinsic Pathway, but is not a Direct Thrombin Inhibitor. <i>TH Open</i> , 2021, 05, e363-e375. | 0.7 | 2 |
| 13 | Transient ST-elevation myocardial infarction versus persistent ST-elevation myocardial infarction. An appraisal of patient characteristics and functional outcome. <i>International Journal of Cardiology</i> , 2021, 336, 22-28. | 0.8 | 4 |
| 14 | Development of a Plasma-Based Assay to Measure the Susceptibility of Factor V to Inhibition by the C-Terminus of TFP1 \pm . <i>Thrombosis and Haemostasis</i> , 2020, 120, 055-064. | 1.8 | 4 |
| 15 | Complement C5 Contributes to Brain Injury After Subarachnoid Hemorrhage. <i>Translational Stroke Research</i> , 2020, 11, 678-688. | 2.3 | 24 |
| 16 | Linkage analysis combined with whole-exome sequencing identifies a novel prothrombin (<i>F2</i>) gene mutation in a Dutch Caucasian family with unexplained thrombosis. <i>Haematologica</i> , 2020, 105, e370-e372. | 1.7 | 6 |
| 17 | Antisense Inhibition of Prekallikrein to Control Hereditary Angioedema. <i>New England Journal of Medicine</i> , 2020, 383, 1242-1247. | 13.9 | 28 |
| 18 | MiR-223-3p and miR-122-5p as circulating biomarkers for plaque instability. <i>Open Heart</i> , 2020, 7, e001223. | 0.9 | 45 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Prediction of cerebral venous thrombosis with a new clinical score and D-dimer levels. <i>Neurology</i> , 2020, 95, e898-e909. | 1.5 | 18 |
| 20 | Nomenclature of factor XI and the contact system. <i>Journal of Thrombosis and Haemostasis</i> , 2019, 17, 2216-2219. | 1.9 | 11 |
| 21 | Coagulation Factor XIII in Cerebral Venous Thrombosis. <i>TH Open</i> , 2019, 03, e227-e229. | 0.7 | 3 |
| 22 | Hydrogenâ€deuterium exchange mass spectrometry highlights conformational changes induced by factor XI activation and binding of factor IX to factor XIa. <i>Journal of Thrombosis and Haemostasis</i> , 2019, 17, 2047-2055. | 1.9 | 16 |
| 23 | In vitro evaluation of factor IX as novel treatment for factor XI deficiency. <i>Blood</i> , 2019, 134, 573-575. | 0.6 | 7 |
| 24 | Zika Virus Infection Induces Elevation of Tissue Factor Production and Apoptosis on Human Umbilical Vein Endothelial Cells. <i>Frontiers in Microbiology</i> , 2019, 10, 817. | 1.5 | 22 |
| 25 | Design and characterization of Î±1-antitrypsin variants for treatment of contact systemâ€driven thromboinflammation. <i>Blood</i> , 2019, 134, 1658-1669. | 0.6 | 20 |
| 26 | An anticoagulant that does not cause bleeding â€“ an abrupt stop on the road to the Holy Grail. <i>Journal of Thrombosis and Haemostasis</i> , 2019, 17, 2019-2021. | 1.9 | 2 |
| 27 | Albumin plasma exchange for life-threatening angioedema with normal C1-inhibitor. <i>Journal of Allergy and Clinical Immunology: in Practice</i> , 2019, 7, 1360-1361. | 2.0 | 2 |
| 28 | Transfusion of autologous extracellular vesicles from stored red blood cells does not affect coagulation in a model of human endotoxemia. <i>Transfusion</i> , 2018, 58, 1486-1493. | 0.8 | 7 |
| 29 | Chemical Footprinting Reveals Conformational Changes Following Activation of Factor XI. <i>Thrombosis and Haemostasis</i> , 2018, 118, 340-350. | 1.8 | 8 |
| 30 | Prochemerin processing by factor XIa. <i>Blood</i> , 2018, 131, 275-276. | 0.6 | 0 |
| 31 | Coagulation and fibrinolysis in hyperparathyroidism secondary to vitamin D deficiency. <i>Endocrine Connections</i> , 2018, 7, 325-333. | 0.8 | 10 |
| 32 | Coagulopathy in Zellweger spectrum disorders: a role for vitamin K. <i>Journal of Inherited Metabolic Disease</i> , 2018, 41, 249-255. | 1.7 | 7 |
| 33 | Hemostatic profile under fluid resuscitation during rivaroxaban anticoagulation: an in vitro survey. <i>Transfusion</i> , 2018, 58, 3014-3026. | 0.8 | 5 |
| 34 | Activation of coagulation and endothelium with concurrent impairment of anticoagulant mechanisms in patients with typhoid fever. <i>Journal of Infection</i> , 2018, 77, 60-67. | 1.7 | 9 |
| 35 | Abnormal coagulation and enhanced fibrinolysis due to lysinuric protein intolerance associates with bleeds and renal impairment. <i>Haemophilia</i> , 2018, 24, e312-e321. | 1.0 | 13 |
| 36 | Factor XI promotes hemostasis in factor XI-deficient mice. <i>Journal of Thrombosis and Haemostasis</i> , 2018, 16, 2044-2049. | 1.9 | 16 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Defective TAFI activation in hemophilia A mice is a major contributor to joint bleeding. <i>Blood</i> , 2018, 132, 1593-1603. | 0.6 | 31 |
| 38 | Role of von Willebrand factor and ADAMTS-13 in early brain injury after experimental subarachnoid hemorrhage. <i>Journal of Thrombosis and Haemostasis</i> , 2018, 16, 1413-1422. | 1.9 | 17 |
| 39 | Elevated factor VIII increases the risk of cerebral venous thrombosis: a case-control study. <i>Journal of Neurology</i> , 2018, 265, 1612-1617. | 1.8 | 15 |
| 40 | Use of DOAC Stop for elimination of anticoagulants in the thrombin generation assay. <i>Thrombosis Research</i> , 2018, 170, 97-101. | 0.8 | 34 |
| 41 | Inhibition of the extrinsic or intrinsic coagulation pathway during pneumonia-derived sepsis. <i>American Journal of Physiology - Lung Cellular and Molecular Physiology</i> , 2018, 315, L799-L809. | 1.3 | 16 |
| 42 | Hemostatic changes associate with mortality in hospitalized patients with HIV-associated tuberculosis: a prospective cohort study. <i>Journal of Infectious Diseases</i> , 2017, 215, jiw532. | 1.9 | 19 |
| 43 | Circulating activated protein C in thrombophilia carriers. <i>Journal of Thrombosis and Thrombolysis</i> , 2017, 43, 361-364. | 1.0 | 1 |
| 44 | High miR-124-3p expression identifies smoking individuals susceptible to atherosclerosis. <i>Atherosclerosis</i> , 2017, 263, 377-384. | 0.4 | 33 |
| 45 | Dynamics of von Willebrand factor reactivity in sickle cell disease during vaso-occlusive crisis and steady state. <i>Journal of Thrombosis and Haemostasis</i> , 2017, 15, 1392-1402. | 1.9 | 20 |
| 46 | Association between protein C levels and mortality in patients with advanced prostate, lung and pancreatic cancer. <i>Thrombosis Research</i> , 2017, 154, 1-6. | 0.8 | 10 |
| 47 | Contact ignition by single-chain XIIa. <i>Blood</i> , 2017, 129, 1411-1412. | 0.6 | 1 |
| 48 | Coagulation factor XIII-A subunit and activation peptide levels in individuals with established symptomatic acute deep vein thrombosis. <i>Thrombosis Research</i> , 2017, 159, 96-99. | 0.8 | 11 |
| 49 | Factor V Leiden is associated with increased sperm count. <i>Human Reproduction</i> , 2017, 32, 2332-2339. | 0.4 | 2 |
| 50 | Novel targets for anticoagulants lacking bleeding risk. <i>Current Opinion in Hematology</i> , 2017, 24, 419-426. | 1.2 | 17 |
| 51 | Increased Circulating and Urinary Levels of Soluble TAM Receptors in Diabetic Nephropathy. <i>American Journal of Pathology</i> , 2017, 187, 1971-1983. | 1.9 | 16 |
| 52 | Editorial: Introduction to the Presentations at the Factor XI and the Contact System SSC Session of the ISTH, Montpellier, France, May 27, 2016. <i>Frontiers in Medicine</i> , 2017, 4, 140. | 1.2 | 0 |
| 53 | Whole exome sequencing in thrombophilic pedigrees to identify genetic risk factors for venous thromboembolism. <i>PLoS ONE</i> , 2017, 12, e0187699. | 1.1 | 14 |
| 54 | Coagulation factor XI improves host defence during murine pneumonia-derived sepsis independent of factor XII activation. <i>Thrombosis and Haemostasis</i> , 2017, 117, 1601-1614. | 1.8 | 40 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | Increased Von Willebrand factor, decreased ADAMTS13 and thrombocytopenia in melioidosis. PLoS Neglected Tropical Diseases, 2017, 11, e0005468. | 1.3 | 7 |
| 56 | Activated protein C protects against renal ischaemia/reperfusion injury, independent of its anticoagulant properties. Thrombosis and Haemostasis, 2016, 116, 124-133. | 1.8 | 9 |
| 57 | Structure-function relationships in thrombin-activatable fibrinolysis inhibitor. Journal of Thrombosis and Haemostasis, 2016, 14, 633-644. | 1.9 | 46 |
| 58 | Impact of HIV infection on the haemostatic response during sepsis and malaria. British Journal of Haematology, 2016, 173, 918-926. | 1.2 | 8 |
| 59 | Loss of asthma control and activation of coagulation and fibrinolysis. Clinical and Experimental Allergy, 2016, 46, 422-427. | 1.4 | 8 |
| 60 | In vivo reversal of the anticoagulant effect of rivaroxaban with four-factor prothrombin complex concentrate. British Journal of Haematology, 2016, 172, 255-261. | 1.2 | 48 |
| 61 | The influence of corticosteroids on hemostasis in healthy subjects. Journal of Thrombosis and Haemostasis, 2016, 14, 716-723. | 1.9 | 45 |
| 62 | Prothrombotic state in patients with severe and prednisolone-dependent asthma. Journal of Allergy and Clinical Immunology, 2016, 137, 1727-1732. | 1.5 | 36 |
| 63 | PO-28 - Protein C levels are associated with mortality in patients with advanced cancer. Thrombosis Research, 2016, 140, S186-S187. | 0.8 | 2 |
| 64 | Stimulation of thrombin- and plasmin-mediated activation of thrombin-activatable fibrinolysis inhibitor by anionic molecules. Thrombosis Research, 2016, 146, 7-14. | 0.8 | 6 |
| 65 | Use of an absorbable embolization material for reversible portal vein embolization in an experimental model. British Journal of Surgery, 2016, 103, 1306-1315. | 0.1 | 9 |
| 66 | The role of ADAMTS13 in acute myocardial infarction: cause or consequence?. Cardiovascular Research, 2016, 111, 194-203. | 1.8 | 24 |
| 67 | Kinetics of coagulation in ST-elevation myocardial infarction following successful primary percutaneous coronary intervention. Thrombosis Research, 2016, 137, 64-71. | 0.8 | 8 |
| 68 | The hypercoagulable state in hyperthyroidism is mediated via the thyroid hormone β 2 receptor pathway. European Journal of Endocrinology, 2016, 174, 755-762. | 1.9 | 21 |
| 69 | Heparin supplement counteracts the prohemostatic effect of prothrombin complex concentrate and factor IX concentrate: An in vitro evaluation. Thrombosis Research, 2016, 139, 102-110. | 0.8 | 7 |
| 70 | A sustained decrease in plasma fibrinolytic potential following partial liver resection or pancreas resection. Thrombosis Research, 2016, 140, 36-40. | 0.8 | 9 |
| 71 | Low miR-19b-1-5p expression in isolated platelets after aspirin use is related to aspirin insensitivity. International Journal of Cardiology, 2016, 203, 262-263. | 0.8 | 32 |
| 72 | Coagulation activation during air travel is not initiated via the extrinsic pathway. British Journal of Haematology, 2015, 169, 903-905. | 1.2 | 4 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 73 | New clues regarding the mysterious mechanism of activated thrombin-activatable fibrinolysis inhibitor self-destruction. <i>Journal of Thrombosis and Haemostasis</i> , 2015, 13, 1081-1083. | 1.9 | 3 |
| 74 | Fresh frozen plasma transfusion fails to influence the hemostatic balance in critically ill patients with a coagulopathy. <i>Journal of Thrombosis and Haemostasis</i> , 2015, 13, 989-997. | 1.9 | 58 |
| 75 | No association between <i>rs11209938</i> genetic variants and deep venous thrombosis. <i>British Journal of Haematology</i> , 2015, 169, 301-304. | 1.2 | 4 |
| 76 | A novel mutation in the F5 gene (factor V Amsterdam) associated with bleeding independent of factor V procoagulant function. <i>Blood</i> , 2015, 125, 1822-1825. | 0.6 | 58 |
| 77 | Factor XIIIa-dependent retention of red blood cells in clots is mediated by fibrin α -chain crosslinking. <i>Blood</i> , 2015, 126, 1940-1948. | 0.6 | 121 |
| 78 | The plasma levels of protein Z-dependent protease inhibitor increase after gynecological surgery independently of estrogen. <i>Thrombosis Research</i> , 2015, 136, 980-986. | 0.8 | 5 |
| 79 | Decreased plasma levels of activated factor VII in patients with deep vein thrombosis. <i>Journal of Thrombosis and Haemostasis</i> , 2015, 13, 1320-1324. | 1.9 | 4 |
| 80 | In vivo increase in thrombin generation by four-factor prothrombin complex concentrate in apixaban-treated healthy volunteers. <i>Journal of Thrombosis and Haemostasis</i> , 2015, 13, 1799-1805. | 1.9 | 66 |
| 81 | Thrombin-activatable fibrinolysis inhibitor influences disease severity in humans and mice with pneumococcal meningitis. <i>Journal of Thrombosis and Haemostasis</i> , 2015, 13, 2076-2086. | 1.9 | 12 |
| 82 | Introduction to the analysis of next generation sequencing data and its application to venous thromboembolism. <i>Thrombosis and Haemostasis</i> , 2015, 114, 920-932. | 1.8 | 10 |
| 83 | Selective modulation of thrombin-activatable fibrinolysis inhibitor (TAFI) activation by thrombin or the thrombin-thrombomodulin complex using TAFI-derived peptides. <i>Journal of Thrombosis and Haemostasis</i> , 2015, 13, 2093-2101. | 1.9 | 4 |
| 84 | Active but inoperable thrombin is accumulated in a plasma protein layer surrounding <i>Streptococcus pyogenes</i> . <i>Thrombosis and Haemostasis</i> , 2015, 114, 717-726. | 1.8 | 4 |
| 85 | Endogenous tissue factor pathway inhibitor has a limited effect on host defence in murine pneumococcal pneumonia. <i>Thrombosis and Haemostasis</i> , 2015, 114, 115-122. | 1.8 | 5 |
| 86 | Normalization panels for the reliable quantification of circulating microRNAs by RT-qPCR. <i>FASEB Journal</i> , 2015, 29, 3853-3862. | 0.2 | 70 |
| 87 | Effect of transfusion of fresh frozen plasma on parameters of endothelial condition and inflammatory status in non-bleeding critically ill patients: a prospective substudy of a randomized trial. <i>Critical Care</i> , 2015, 19, 163. | 2.5 | 71 |
| 88 | Lowering blood glucose during hip surgery does not influence coagulation activation. <i>BBA Clinical</i> , 2015, 3, 227-232. | 4.1 | 6 |
| 89 | Hemostasis and Fibrinolysis in Delayed Cerebral Ischemia after Aneurysmal Subarachnoid Hemorrhage: A Systematic Review. <i>Journal of Cerebral Blood Flow and Metabolism</i> , 2015, 35, 724-733. | 2.4 | 55 |
| 90 | The cytoprotective effects of endogenous activated protein C reduce activation of coagulation during murine pneumococcal pneumonia and sepsis. <i>Thrombosis Research</i> , 2015, 135, 537-543. | 0.8 | 12 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 91 | Markers of endothelial cell activation and immune activation are increased in patients with severe leptospirosis and associated with disease severity. <i>Journal of Infection</i> , 2015, 71, 437-446. | 1.7 | 17 |
| 92 | Effect of Puumala hantavirus infection on human umbilical vein endothelial cell hemostatic function: platelet interactions, increased tissue factor expression and fibrinolysis regulator release. <i>Frontiers in Microbiology</i> , 2015, 6, 220. | 1.5 | 28 |
| 93 | Development of a Hyperactive Primary Hemostatic System During Off-Pump Lung Transplantation Resulting From an Unbalance Between von Willebrand Factor and Its Cleaving Protease ADAMTS13. <i>American Journal of Transplantation</i> , 2015, 15, 1958-1966. | 2.6 | 3 |
| 94 | Lack of TAFI increases brain damage and microparticle generation after thrombolytic therapy in ischemic stroke. <i>Thrombosis Research</i> , 2015, 136, 445-450. | 0.8 | 15 |
| 95 | Effect of the oral thrombin inhibitor dabigatran on allergic lung inflammation induced by repeated house dust mite administration in mice. <i>American Journal of Physiology - Lung Cellular and Molecular Physiology</i> , 2015, 309, L768-L775. | 1.3 | 8 |
| 96 | Pulmonary tuberculosis induces a systemic hypercoagulable state. <i>Journal of Infection</i> , 2015, 70, 324-334. | 1.7 | 48 |
| 97 | Activated protein C inhibits neutrophil migration in allergic asthma: a randomised trial. <i>European Respiratory Journal</i> , 2015, 46, 1636-1644. | 3.1 | 16 |
| 98 | Development of a Hypercoagulable Status in Patients Undergoing Off-Pump Lung Transplantation despite Prolonged Conventional Coagulation Tests. <i>American Journal of Respiratory and Critical Care Medicine</i> , 2015, 191, 230-233. | 2.5 | 10 |
| 99 | Differential In Vitro Inhibition of Thrombin Generation by Anticoagulant Drugs in Plasma from Patients with Cirrhosis. <i>PLoS ONE</i> , 2014, 9, e88390. | 1.1 | 79 |
| 100 | Individual with Subclinical Atherosclerosis Have Impaired Proliferation of Blood Outgrowth Endothelial Cells, Which Can Be Restored by Statin Therapy. <i>PLoS ONE</i> , 2014, 9, e99890. | 1.1 | 11 |
| 101 | Individuals with coronary artery disease at a young age and features of the metabolic syndrome have an increased prothrombotic potential. <i>Thrombosis and Haemostasis</i> , 2014, 111, 458-464. | 1.8 | 6 |
| 102 | The endothelial protein C receptor impairs the antibacterial response in murine pneumococcal pneumonia and sepsis. <i>Thrombosis and Haemostasis</i> , 2014, 111, 970-980. | 1.8 | 9 |
| 103 | Overexpression of activated protein C hampers bacterial dissemination during pneumococcal pneumonia. <i>BMC Infectious Diseases</i> , 2014, 14, 559. | 1.3 | 9 |
| 104 | Effect of recombinant ADAMTS-13 on microthrombosis and brain injury after experimental subarachnoid hemorrhage. <i>Journal of Thrombosis and Haemostasis</i> , 2014, 12, 943-947. | 1.9 | 27 |
| 105 | Altered platelet contents in survivors of early ischemic ventricular fibrillation: Preliminary findings. <i>Platelets</i> , 2014, 25, 71-74. | 1.1 | 0 |
| 106 | Recent insights into the role of the contact pathway in thrombo-inflammatory disorders. <i>Hematology American Society of Hematology Education Program</i> , 2014, 2014, 60-65. | 0.9 | 39 |
| 107 | Hyperferritinaemia in Dengue Virus Infected Patients Is Associated with Immune Activation and Coagulation Disturbances. <i>PLoS Neglected Tropical Diseases</i> , 2014, 8, e3214. | 1.3 | 46 |
| 108 | Utility of thromboelastography and/or thromboelastometry in adults with sepsis: a systematic review. <i>Critical Care</i> , 2014, 18, R30. | 2.5 | 185 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 109 | Mutations in <i>STAP1</i> Are Associated With Autosomal Dominant Hypercholesterolemia. <i>Circulation Research</i> , 2014, 115, 552-555. | 2.0 | 146 |
| 110 | A role for arginine ¹² in thrombin ¹² -thrombomodulin ¹² -mediated activation of thrombin ¹² -activatable fibrinolysis inhibitor. <i>Journal of Thrombosis and Haemostasis</i> , 2014, 12, 1717-1725. | 1.9 | 14 |
| 111 | Acquired von <i>Willebrand</i> syndrome in patients with overt hypothyroidism: a prospective cohort study. <i>Haemophilia</i> , 2014, 20, 326-332. | 1.0 | 28 |
| 112 | Factor XI Regulates Pathological Thrombus Formation on Acutely Ruptured Atherosclerotic Plaques. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2014, 34, 1668-1673. | 1.1 | 47 |
| 113 | Plasminogen activator inhibitor-1 influences cerebrovascular complications and death in pneumococcal meningitis. <i>Acta Neuropathologica</i> , 2014, 127, 553-564. | 3.9 | 17 |
| 114 | Evaluation of coagulation activation after Rhinovirus infection in patients with asthma and healthy control subjects: an observational study. <i>Respiratory Research</i> , 2014, 15, 14. | 1.4 | 21 |
| 115 | Factor XII Regulates the Pathological Process of Thrombus Formation on Ruptured Plaques. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2014, 34, 1674-1680. | 1.1 | 108 |
| 116 | Evaluating the Optimal Timing of Revascularisation in Patients with Transient ST-Segment Elevation Myocardial Infarction: Rationale and Design of the TRANSIENT Trial. <i>Journal of Cardiovascular Translational Research</i> , 2014, 7, 590-596. | 1.1 | 10 |
| 117 | Activation of coagulation and tissue fibrin deposition in experimental influenza in ferrets. <i>BMC Microbiology</i> , 2014, 14, 134. | 1.3 | 30 |
| 118 | The influence of HIV infection on coagulation activation during sepsis. <i>International Journal of Infectious Diseases</i> , 2014, 21, 128. | 1.5 | 0 |
| 119 | Clot stability and fibrin deposition is strongly reduced in mice in which mouse TAFI is replaced by human TAFI. <i>Thrombosis Research</i> , 2014, 133, 1166-1168. | 0.8 | 0 |
| 120 | No contact, no thrombosis?. <i>Blood</i> , 2014, 123, 1629-1629. | 0.6 | 6 |
| 121 | Sustained pro-haemostatic activity of rFVIIa in plasma and platelets in non-bleeding pigs may explain the efficacy of a once-daily prophylaxis in humans. <i>Thrombosis and Haemostasis</i> , 2014, 112, 304-310. | 1.8 | 6 |
| 122 | Optimisation of lupus anticoagulant tests: should test samples always be mixed with normal plasma?. <i>Thrombosis and Haemostasis</i> , 2014, 112, 736-742. | 1.8 | 26 |
| 123 | C0288: Lack of TAFI Has Deleterious Effect on Experimental Ischemic Stroke: Potential Role of Microparticles. <i>Thrombosis Research</i> , 2014, 133, S5. | 0.8 | 1 |
| 124 | Treatment with everolimus is associated with a procoagulant state. <i>Thrombosis Research</i> , 2013, 132, 307-311. | 0.8 | 42 |
| 125 | Endogenous protein C has a protective role during Gram-negative pneumosepsis (melioidosis). <i>Journal of Thrombosis and Haemostasis</i> , 2013, 11, 282-292. | 1.9 | 20 |
| 126 | Urinary prothrombin fragment 1+2 in patients with venous thrombosis and myocardial infarction. <i>Journal of Thrombosis and Thrombolysis</i> , 2013, 36, 47-49. | 1.0 | 0 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 127 | The Aggregation Potential of the 1â€“15- and 1â€“16-Fragments of the Amyloid Î² Peptide and Their Influence on the Aggregation of AÎ²40. <i>Applied Magnetic Resonance</i> , 2013, 44, 1167-1179. | 0.6 | 3 |
| 128 | Circulating Nucleosomes and Neutrophil Activation as Risk Factors for Deep Vein Thrombosis. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2013, 33, 147-151. | 1.1 | 193 |
| 129 | Endogenous Î±2-Antiplasmin Is Protective during Severe Gram-Negative Sepsis (Melioidosis). <i>American Journal of Respiratory and Critical Care Medicine</i> , 2013, 188, 967-975. | 2.5 | 33 |
| 130 | Decreased tissue factor pathway inhibitor (<sc>TFPI</sc>)â€“dependent anticoagulant capacity in patients with cirrhosis who have decreased protein <sc>S</sc> but normal <sc>TFPI</sc> plasma levels. <i>British Journal of Haematology</i> , 2013, 162, 819-826. | 1.2 | 14 |
| 131 | VEGF Levels in Plasma in Relation to Platelet Activation, Glycemic Control, and Microvascular Complications in Type 1 Diabetes. <i>Diabetes Care</i> , 2013, 36, 1629-1634. | 4.3 | 40 |
| 132 | Lamin A/C mutation is independently associated with an increased risk of arterial and venous thromboembolic complications. <i>International Journal of Cardiology</i> , 2013, 168, 472-477. | 0.8 | 33 |
| 133 | Thrombin-activatable fibrinolysis inhibitor in hypothyroidism and hyperthyroxinaemia. <i>Thrombosis and Haemostasis</i> , 2013, 109, 214-220. | 1.8 | 16 |
| 134 | An unbalance between von Willebrand factor and ADAMTS13 in acute liver failure: Implications for hemostasis and clinical outcome. <i>Hepatology</i> , 2013, 58, 752-761. | 3.6 | 153 |
| 135 | Overexpression of the Endothelial Protein C Receptor Is Detrimental during Pneumonia-Derived Gram-negative Sepsis (Melioidosis). <i>PLoS Neglected Tropical Diseases</i> , 2013, 7, e2306. | 1.3 | 14 |
| 136 | The lectin-like domain of thrombomodulin hampers host defence in pneumococcal pneumonia. <i>European Respiratory Journal</i> , 2013, 41, 935-942. | 3.1 | 8 |
| 137 | Factor Xa Activation of Factor V Is of Paramount Importance in Initiating the Coagulation System. <i>Circulation</i> , 2013, 128, 254-266. | 1.6 | 84 |
| 138 | Differential Effects of Nonselective Versus Selective Î²-Blockers on Cardiac Sympathetic Activity and Hemostasis in Patients with Heart Failure. <i>Journal of Nuclear Medicine</i> , 2013, 54, 1733-1739. | 2.8 | 11 |
| 139 | Clot lysis phenotype and response to recombinant factor <sc>VII</sc>a in plasma of haemophilia A inhibitor patients. <i>British Journal of Haematology</i> , 2013, 162, 827-835. | 1.2 | 10 |
| 140 | Intrabronchial activated protein C enhances lipopolysaccharide-induced pulmonary responses. <i>European Respiratory Journal</i> , 2013, 42, 188-197. | 3.1 | 10 |
| 141 | Obesity in haemophilia patients: effect on bleeding frequency, clotting factor concentrate usage, and haemostatic and fibrinolytic parameters. <i>Haemophilia</i> , 2013, 19, 744-752. | 1.0 | 22 |
| 142 | Statin therapy and levels of hemostatic factors in a healthy population: the Multi-Ethnic Study of Atherosclerosis: a rebuttal. <i>Journal of Thrombosis and Haemostasis</i> , 2013, 11, 1787-1788. | 1.9 | 2 |
| 143 | Two novel inhibitory anti-human factor XI antibodies prevent cessation of blood flow in a murine venous thrombosis model. <i>Thrombosis and Haemostasis</i> , 2013, 110, 1065-1073. | 1.8 | 30 |
| 144 | Differences in platelet microRNA profiles after aspirin use are associated with differences in whole blood aggregation and might identify aspirin resistance. <i>European Heart Journal</i> , 2013, 34, P4884-P4884. | 1.0 | 0 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 145 | Overexpression of Activated Protein C is Detrimental During Severe Experimental Gram-Negative Sepsis (Melioidosis)*. <i>Critical Care Medicine</i> , 2013, 41, e266-e274. | 0.4 | 8 |
| 146 | TAFI deficiency promotes liver damage in murine models of liver failure through defective down-regulation of hepatic inflammation. <i>Thrombosis and Haemostasis</i> , 2013, 109, 948-955. | 1.8 | 19 |
| 147 | The endothelial protein C receptor and activated protein C play a limited role in host defense during experimental tuberculosis. <i>Thrombosis and Haemostasis</i> , 2013, 109, 726-737. | 1.8 | 7 |
| 148 | Venous thromboembolism in adults treated for acute lymphoblastic leukaemia: Effect of fresh frozen plasma supplementation. <i>Thrombosis and Haemostasis</i> , 2013, 109, 633-642. | 1.8 | 33 |
| 149 | Anticoagulation beyond direct thrombin and factor Xa inhibitors: indications for targeting the intrinsic pathway?. <i>Thrombosis and Haemostasis</i> , 2013, 110, 223-232. | 1.8 | 33 |
| 150 | Endogenous tissue-type plasminogen activator impairs host defense during severe experimental gram-negative sepsis (melioidosis)*. <i>Critical Care Medicine</i> , 2012, 40, 2168-2175. | 0.4 | 19 |
| 151 | BH3-only protein Noxa regulates apoptosis in activated B cells and controls high-affinity antibody formation. <i>Blood</i> , 2012, 119, 1440-1449. | 0.6 | 33 |
| 152 | Intact thrombin generation and decreased fibrinolytic capacity in patients with acute liver injury or acute liver failure. <i>Journal of Thrombosis and Haemostasis</i> , 2012, 10, 1312-1319. | 1.9 | 101 |
| 153 | Microcirculation and atherothrombotic parameters in prolactinoma patients: a pilot study. <i>Pituitary</i> , 2012, 15, 472-481. | 1.6 | 13 |
| 154 | Recombinant C1-Inhibitor. <i>BioDrugs</i> , 2012, 26, 43-52. | 2.2 | 28 |
| 155 | Chronic pulmonary embolism in Klippel-Trenaunay syndrome. <i>Journal of the American Academy of Dermatology</i> , 2012, 66, 71-77. | 0.6 | 31 |
| 156 | Review: Viral infections and mechanisms of thrombosis and bleeding. <i>Journal of Medical Virology</i> , 2012, 84, 1680-1696. | 2.5 | 252 |
| 157 | Reply from the authors. <i>Hepatology Research</i> , 2012, 42, 1154-1155. | 1.8 | 0 |
| 158 | The hyperfibrinolytic state of mice with combined thrombin-activatable fibrinolysis inhibitor (TAFI) and plasminogen activator inhibitor-1 gene deficiency is critically dependent on TAFI deficiency. <i>Journal of Thrombosis and Haemostasis</i> , 2012, 10, 2555-2562. | 1.9 | 22 |
| 159 | Effects of the etonogestrel-releasing contraceptive implant inserted immediately postpartum on maternal hemostasis: A randomized controlled trial. <i>Thrombosis Research</i> , 2012, 130, 355-360. | 0.8 | 17 |
| 160 | Increase in the plasma levels of protein Z-dependent protease inhibitor in normal pregnancies but not in non-pregnant patients with unexplained recurrent miscarriage. <i>Thrombosis and Haemostasis</i> , 2012, 107, 507-512. | 1.8 | 5 |
| 161 | Recent developments in our understanding of the antiphospholipid syndrome. <i>International Journal of Laboratory Hematology</i> , 2012, 34, 223-231. | 0.7 | 26 |
| 162 | <i>Factor XIII Val34Leu</i> mutation accelerates the development of fibrosis in patients with chronic hepatitis B and C. <i>Hepatology Research</i> , 2012, 42, 668-676. | 1.8 | 11 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 163 | Clinical practice. <i>European Journal of Pediatrics</i> , 2012, 171, 207-214. | 1.3 | 21 |
| 164 | Blood transfusion during cardiac surgery is associated with inflammation and coagulation in the lung: a case control study. <i>Critical Care</i> , 2011, 15, R59. | 2.5 | 50 |
| 165 | The Hemostatic System in Patients With Type 2 Diabetes With and Without Cardiovascular Disease. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2011, 17, E57-E63. | 0.7 | 23 |
| 166 | Plasma levels of pentraxin-3, an acute phase protein, are increased during sickle cell painful crisis. <i>Blood Cells, Molecules, and Diseases</i> , 2011, 46, 189-194. | 0.6 | 18 |
| 167 | Activated human platelet products induce proarrhythmic effects in ventricular myocytes. <i>Journal of Molecular and Cellular Cardiology</i> , 2011, 51, 347-356. | 0.9 | 5 |
| 168 | Acute Stress Elicited by Bungee Jumping Suppresses Human Innate Immunity. <i>Molecular Medicine</i> , 2011, 17, 180-188. | 1.9 | 36 |
| 169 | Quantitative trait locus for protein C in a family with thrombophilia. <i>Thrombosis and Haemostasis</i> , 2011, 105, 199-201. | 1.8 | 4 |
| 170 | Binding characteristics of thrombin-activatable fibrinolysis inhibitor to streptococcal surface collagen-like proteins A and B. <i>Thrombosis and Haemostasis</i> , 2011, 106, 609-616. | 1.8 | 6 |
| 171 | Diabetes does not influence activation of coagulation, fibrinolysis or anticoagulant pathways in Gram-negative sepsis (melioidosis). <i>Thrombosis and Haemostasis</i> , 2011, 106, 1139-1148. | 1.8 | 13 |
| 172 | Evolutionary conservation of the lipopolysaccharide binding site of β_2 -glycoprotein I. <i>Thrombosis and Haemostasis</i> , 2011, 106, 1069-1075. | 1.8 | 34 |
| 173 | Platelets in Patients with Premature Coronary Artery Disease Exhibit Upregulation of miRNA340* and miRNA624*. <i>PLoS ONE</i> , 2011, 6, e25946. | 1.1 | 86 |
| 174 | Identification and Characterization of Ixodes scapularis Antigens That Elicit Tick Immunity Using Yeast Surface Display. <i>PLoS ONE</i> , 2011, 6, e15926. | 1.1 | 72 |
| 175 | Vascular Endothelial Growth Factor in the Circulation in Cancer Patients May Not Be a Relevant Biomarker. <i>PLoS ONE</i> , 2011, 6, e19873. | 1.1 | 43 |
| 176 | Endothelial dysfunction, platelet activation, thrombogenesis and fibrinolysis in patients with hypertensive crisis. <i>Journal of Hypertension</i> , 2011, 29, 922-927. | 0.3 | 48 |
| 177 | The hemostatic status of pediatric recipients of adult liver grafts suggests that plasma levels of hemostatic proteins are not regulated by the liver. <i>Blood</i> , 2011, 117, 2070-2072. | 0.6 | 22 |
| 178 | β_2 -Glycoprotein I: a novel component of innate immunity. <i>Blood</i> , 2011, 117, 6939-6947. | 0.6 | 101 |
| 179 | Increased coagulation factor VIII activity in patients with familial hypercholesterolemia. <i>Blood</i> , 2011, 118, 6990-6991. | 0.6 | 11 |
| 180 | Decreased active von Willebrand factor level owing to shear stress in aortic stenosis patients. <i>Journal of Thrombosis and Haemostasis</i> , 2011, 9, 953-958. | 1.9 | 21 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 181 | High levels of protein C are determined by PROCR haplotype 3. <i>Journal of Thrombosis and Haemostasis</i> , 2011, 9, 969-976. | 1.9 | 17 |
| 182 | β ₂ -Glycoprotein I: evolution, structure and function. <i>Journal of Thrombosis and Haemostasis</i> , 2011, 9, 1275-1284. | 1.9 | 180 |
| 183 | Acute respiratory tract infection leads to procoagulant changes in human subjects. <i>Journal of Thrombosis and Haemostasis</i> , 2011, 9, 1432-1434. | 1.9 | 23 |
| 184 | Detection of lupus anticoagulant in the presence of rivaroxaban using Taipan snake venom time. <i>Journal of Thrombosis and Haemostasis</i> , 2011, 9, 1657-1659. | 1.9 | 82 |
| 185 | Influenza vaccination and hemostasis: no sustainable procoagulant effects from 2009 H1N1 influenza vaccine in healthy healthcare workers. <i>Journal of Thrombosis and Haemostasis</i> , 2011, 9, 1659-1661. | 1.9 | 1 |
| 186 | Alterations in coagulation and fibrinolysis after levothyroxine exposure in healthy volunteers: a controlled randomized crossover study. <i>Journal of Thrombosis and Haemostasis</i> , 2011, 9, 1816-1824. | 1.9 | 40 |
| 187 | Plasminogen activator inhibitor type I contributes to protective immunity during experimental Gram-negative sepsis (melioidosis). <i>Journal of Thrombosis and Haemostasis</i> , 2011, 9, 2020-2028. | 1.9 | 34 |
| 188 | Induction of anti-β ₂ -glycoprotein I autoantibodies in mice by protein H of <i>Streptococcus pyogenes</i> . <i>Journal of Thrombosis and Haemostasis</i> , 2011, 9, 2447-2456. | 1.9 | 35 |
| 189 | Protein C Inhibitor. <i>Seminars in Thrombosis and Hemostasis</i> , 2011, 37, 349-354. | 1.5 | 34 |
| 190 | Forgotten Factors in Hemostasis and Thrombosis. <i>Seminars in Thrombosis and Hemostasis</i> , 2011, 37, 337-338. | 1.5 | 0 |
| 191 | β ₂ -Glycoprotein I can exist in 2 conformations: implications for our understanding of the antiphospholipid syndrome. <i>Blood</i> , 2010, 116, 1336-1343. | 0.6 | 247 |
| 192 | Venous thrombosis risk associated with plasma hypofibrinolysis is explained by elevated plasma levels of TAFI and PAI-1. <i>Blood</i> , 2010, 116, 113-121. | 0.6 | 309 |
| 193 | The effects of an extremely high dose of levothyroxine on coagulation and fibrinolysis. <i>Journal of Thrombosis and Haemostasis</i> , 2010, 8, 1427-1428. | 1.9 | 13 |
| 194 | Expression patterns of protein C inhibitor in mouse development. <i>Journal of Molecular Histology</i> , 2010, 41, 27-37. | 1.0 | 7 |
| 195 | Markers of inflammation and coagulation indicate a prothrombotic state in HIV-infected patients with long-term use of antiretroviral therapy with or without abacavir. <i>AIDS Research and Therapy</i> , 2010, 7, 9. | 0.7 | 40 |
| 196 | Increased sperm count may account for high population frequency of factor V Leiden. <i>Journal of Thrombosis and Haemostasis</i> , 2010, 8, 513-516. | 1.9 | 11 |
| 197 | Explanations for coagulation activation after air travel. <i>Journal of Thrombosis and Haemostasis</i> , 2010, 8, 971-978. | 1.9 | 25 |
| 198 | Intact thrombomodulin-mediated regulation of fibrinolysis during and after liver transplantation, despite a profoundly defective thrombomodulin-mediated regulation of coagulation. <i>Journal of Thrombosis and Haemostasis</i> , 2010, 8, 1646-1649. | 1.9 | 12 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 199 | Hyperglycemia: a prothrombotic factor?. <i>Journal of Thrombosis and Haemostasis</i> , 2010, 8, 1663-1669. | 1.9 | 227 |
| 200 | Thrombin-activatable fibrinolysis inhibitor is degraded by <i>Salmonella enterica</i> and <i>Yersinia pestis</i> . <i>Journal of Thrombosis and Haemostasis</i> , 2010, 8, 2232-2240. | 1.9 | 21 |
| 201 | Coagulation factor XI as a novel target for antithrombotic treatment. <i>Journal of Thrombosis and Haemostasis</i> , 2010, 8, 2349-2357. | 1.9 | 67 |
| 202 | Coagulation disorders in patients with severe leptospirosis are associated with severe bleeding and mortality. <i>Tropical Medicine and International Health</i> , 2010, 15, 152-159. | 1.0 | 61 |
| 203 | The effect of initiating combined antiretroviral therapy on endothelial cell activation and coagulation markers in South African HIV-infected individuals. <i>Thrombosis and Haemostasis</i> , 2010, 104, 1228-1234. | 1.8 | 58 |
| 204 | Severe malaria is associated with a deficiency of von Willebrand factor cleaving protease, ADAMTS13. <i>Thrombosis and Haemostasis</i> , 2010, 103, 181-187. | 1.8 | 70 |
| 205 | Inhibition of the intrinsic coagulation pathway factor XI by antisense oligonucleotides: a novel antithrombotic strategy with lowered bleeding risk. <i>Blood</i> , 2010, 116, 4684-4692. | 0.6 | 172 |
| 206 | Urokinase Receptor Is Necessary for Bacterial Defense against Pneumonia-Derived Septic Melioidosis by Facilitating Phagocytosis. <i>Journal of Immunology</i> , 2010, 184, 3079-3086. | 0.4 | 49 |
| 207 | Activated factor V is a cofactor for the activation of factor XI by thrombin in plasma. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010, 107, 9083-9087. | 3.3 | 30 |
| 208 | Systemic Inflammatory Response Syndrome and Compensatory Anti-Inflammatory Response Syndrome in Sepsis. <i>Journal of Innate Immunity</i> , 2010, 2, 379-380. | 1.8 | 33 |
| 209 | Normal to increased thrombin generation in patients undergoing liver transplantation despite prolonged conventional coagulation tests. <i>Journal of Hepatology</i> , 2010, 52, 355-361. | 1.8 | 191 |
| 210 | Clinical usefulness of prothrombin fragment 1+2 in patients with suspected pulmonary embolism. <i>Thrombosis Research</i> , 2010, 125, 97-99. | 0.8 | 6 |
| 211 | The role of thrombin-activatable fibrinolysis inhibitor in diabetic wound healing. <i>Thrombosis Research</i> , 2010, 126, 442-446. | 0.8 | 15 |
| 212 | Thrombin-Activatable Fibrinolysis Inhibitor (TAFI) Deficient Mice Are Susceptible to Intracerebral Thrombosis and Ischemic Stroke. <i>PLoS ONE</i> , 2010, 5, e11658. | 1.1 | 32 |
| 213 | Assessment of coagulation and fibrinolysis in families with unexplained thrombophilia. <i>Thrombosis and Haemostasis</i> , 2009, 101, 465-470. | 1.8 | 30 |
| 214 | The effects of hyperglycaemia on thrombin-activatable fibrinolysis inhibitor. <i>Thrombosis and Haemostasis</i> , 2009, 102, 460-468. | 1.8 | 8 |
| 215 | The Hemostatic Balance in HIV-Infected Patients with and without Antiretroviral Therapy: Partial Restoration with Antiretroviral Therapy. <i>AIDS Patient Care and STDs</i> , 2009, 23, 1001-1007. | 1.1 | 47 |
| 216 | Activation of TAFI on the Surface of <i>Streptococcus pyogenes</i> Evokes Inflammatory Reactions by Modulating the Kallikrein/Kinin System. <i>Journal of Innate Immunity</i> , 2009, 1, 18-28. | 1.8 | 33 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 217 | Plasminogen activator inhibitor-1 regulates neutrophil influx during acute pyelonephritis. <i>Kidney International</i> , 2009, 75, 52-59. | 2.6 | 35 |
| 218 | Recent Developments in Thrombin-Activatable Fibrinolysis Inhibitor Research. <i>Mini-Reviews in Medicinal Chemistry</i> , 2009, 9, 1165-1173. | 1.1 | 14 |
| 219 | Protein C Inhibitor—A Novel Antimicrobial Agent. <i>PLoS Pathogens</i> , 2009, 5, e1000698. | 2.1 | 34 |
| 220 | The Receptor for Advanced Glycation End Products Impairs Host Defense in Pneumococcal Pneumonia. <i>Journal of Immunology</i> , 2009, 182, 4349-4356. | 0.4 | 99 |
| 221 | ROLE OF TOLL-LIKE RECEPTORS 2 AND 4, AND THE RECEPTOR FOR ADVANCED GLYCATION END PRODUCTS IN HIGH-MOBILITY GROUP BOX 1-INDUCED INFLAMMATION IN VIVO. <i>Shock</i> , 2009, 31, 280-284. | 1.0 | 237 |
| 222 | Low thrombin activatable fibrinolysis inhibitor activity levels are associated with an increased risk of a first myocardial infarction in men. <i>Haematologica</i> , 2009, 94, 811-818. | 1.7 | 42 |
| 223 | Biologic Effects of Simvastatin in Patients with Aneurysmal Subarachnoid Hemorrhage: A Double-Blind, Placebo-Controlled Randomized Trial. <i>Journal of Cerebral Blood Flow and Metabolism</i> , 2009, 29, 1444-1453. | 2.4 | 118 |
| 224 | Reduced ADAMTS13 Activity in Delayed Cerebral Ischemia after Aneurysmal Subarachnoid Hemorrhage. <i>Journal of Cerebral Blood Flow and Metabolism</i> , 2009, 29, 1734-1741. | 2.4 | 42 |
| 225 | ApoA1-phosphatidylcholine infusion neutralizes the atherothrombotic effects of C-reactive protein in humans. <i>Journal of Thrombosis and Haemostasis</i> , 2009, 7, 347-354. | 1.9 | 20 |
| 226 | Î2-Glycoprotein I is incorrectly named apolipoprotein H. <i>Journal of Thrombosis and Haemostasis</i> , 2009, 7, 235-236. | 1.9 | 54 |
| 227 | The activation peptide of thrombin-activatable fibrinolysis inhibitor: a role in activity and stability of the enzyme?. <i>Journal of Thrombosis and Haemostasis</i> , 2009, 7, 445-452. | 1.9 | 20 |
| 228 | Venous thrombosis is associated with hyperglycemia at diagnosis: a case-control study. <i>Journal of Thrombosis and Haemostasis</i> , 2009, 7, 945-949. | 1.9 | 40 |
| 229 | Circulating erythrocyte-derived microparticles are associated with coagulation activation in sickle cell disease. <i>Haematologica</i> , 2009, 94, 1513-1519. | 1.7 | 241 |
| 230 | Tumor necrosis factor-Î± inhibition protects against endotoxin-induced endothelial glycocalyx perturbation. <i>Atherosclerosis</i> , 2009, 202, 296-303. | 0.4 | 136 |
| 231 | Urokinase Plasminogen Activator Receptor-Deficient Mice Demonstrate Reduced Hyperoxia-Induced Lung Injury. <i>American Journal of Pathology</i> , 2009, 174, 2182-2189. | 1.9 | 12 |
| 232 | Feedback controversy stops here. <i>Blood</i> , 2009, 114, 235-235. | 0.6 | 10 |
| 233 | Assessment of coagulation and fibrinolysis in families with unexplained thrombophilia. <i>Thrombosis and Haemostasis</i> , 2009, 101, 465-70. | 1.8 | 6 |
| 234 | Development of sandwich-type ELISAs for the quantification of rat and murine thrombin activatable fibrinolysis inhibitor in plasma. <i>Journal of Thrombosis and Haemostasis</i> , 2008, 6, 132-138. | 1.9 | 9 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 235 | Activation of coagulation with concurrent impairment of anticoagulant mechanisms correlates with a poor outcome in severe melioidosis. <i>Journal of Thrombosis and Haemostasis</i> , 2008, 6, 32-39. | 1.9 | 38 |
| 236 | Effect of the selective serotonin reuptake inhibitor paroxetine on platelet function is modified by a SLC6A4 serotonin transporter polymorphism. <i>Journal of Thrombosis and Haemostasis</i> , 2008, 6, 2168-2174. | 1.9 | 31 |
| 237 | The antibacterial activity of peptides derived from human beta α 2 glycoprotein I is inhibited by protein H and M1 protein from <i>Streptococcus pyogenes</i> . <i>Molecular Microbiology</i> , 2008, 67, 482-492. | 1.2 | 32 |
| 238 | Nur77 modulates hepatic lipid metabolism through suppression of SREBP1c activity. <i>Biochemical and Biophysical Research Communications</i> , 2008, 366, 910-916. | 1.0 | 55 |
| 239 | Effects of short-term and long-term danazol treatment on lipoproteins, coagulation, and progression of atherosclerosis: Two clinical trials in healthy volunteers and patients with hereditary angioedema. <i>Clinical Therapeutics</i> , 2008, 30, 2314-2323. | 1.1 | 46 |
| 240 | Association Between Thrombotic Microangiopathy and Reduced ADAMTS13 Activity in Malignant Hypertension. <i>Hypertension</i> , 2008, 51, 862-866. | 1.3 | 80 |
| 241 | Platelet-Activating Factor Receptor Contributes to Host Defense against <i>Pseudomonas aeruginosa</i> Pneumonia but Is Not Essential for the Accompanying Inflammatory and Procoagulant Response. <i>Journal of Immunology</i> , 2008, 180, 3357-3365. | 0.4 | 16 |
| 242 | Perioperative hyperinsulinaemic normoglycaemic clamp causes hypolipidaemia after coronary artery surgery. <i>British Journal of Anaesthesia</i> , 2008, 100, 442-450. | 1.5 | 26 |
| 243 | Endogenous β 2-Adrenergic Receptors Inhibit Lipopolysaccharide-Induced Pulmonary Cytokine Release and Coagulation. <i>American Journal of Respiratory Cell and Molecular Biology</i> , 2008, 39, 373-379. | 1.4 | 13 |
| 244 | Crystal structures of TAFI elucidate the inactivation mechanism of activated TAFI: a novel mechanism for enzyme autoregulation. <i>Blood</i> , 2008, 112, 2803-2809. | 0.6 | 69 |
| 245 | Hyperglycemia enhances coagulation and reduces neutrophil degranulation, whereas hyperinsulinemia inhibits fibrinolysis during human endotoxemia. <i>Blood</i> , 2008, 112, 82-89. | 0.6 | 221 |
| 246 | Fluid loss does not explain coagulation activation during air travel. <i>Thrombosis and Haemostasis</i> , 2008, 99, 1053-1059. | 1.8 | 31 |
| 247 | Circulating Erythrocyte-Derived Microparticles Are Associated with Coagulation Activation in Sickle Cell Disease. <i>Blood</i> , 2008, 112, 126-126. | 0.6 | 7 |
| 248 | Severe Malaria and Leptospirosis Are Associated with a Deficiency of the Von Willebrand Factor Cleaving Protease, ADAMTS13. <i>Blood</i> , 2008, 112, 3912-3912. | 0.6 | 2 |
| 249 | Thrombin-activatable Fibrinolysis Inhibitor Binds to <i>Streptococcus pyogenes</i> by Interacting with Collagen-like Proteins A and B. <i>Journal of Biological Chemistry</i> , 2007, 282, 24873-24881. | 1.6 | 36 |
| 250 | Prednisolone Dose-Dependently Influences Inflammation and Coagulation during Human Endotoxemia. <i>Journal of Immunology</i> , 2007, 178, 1845-1851. | 0.4 | 94 |
| 251 | Effects of CRP infusion on endothelial function and coagulation in normocholesterolemic and hypercholesterolemic subjects. <i>Journal of Lipid Research</i> , 2007, 48, 952-960. | 2.0 | 65 |
| 252 | Pharmacokinetics and pharmacodynamics of danaparoid during continuous venovenous hemofiltration: a pilot study. <i>Critical Care</i> , 2007, 11, R102. | 2.5 | 20 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 253 | EFFECTS OF INTRA-ABDOMINAL ADMINISTRATION OF RECOMBINANT TISSUE PLASMINOGEN ACTIVATOR ON COAGULATION, FIBRINOLYSIS AND INFLAMMATORY RESPONSES IN EXPERIMENTAL POLYMICROBIAL PERITONITIS. <i>Shock</i> , 2007, 27, 534-541. | 1.0 | 9 |
| 254 | High-Density Lipoprotein Attenuates Inflammation and Coagulation Response on Endotoxin Challenge in Humans. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2007, 27, 1153-1158. | 1.1 | 102 |
| 255 | Alterations in lipoprotein homeostasis during human experimental endotoxemia and clinical sepsis. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2007, 1771, 1429-1438. | 1.2 | 76 |
| 256 | Tissue factor-dependent blood coagulation is enhanced following delivery irrespective of the mode of delivery. <i>Journal of Thrombosis and Haemostasis</i> , 2007, 5, 2415-2420. | 1.9 | 36 |
| 257 | Activation of coagulation system during air travel: a crossover study. <i>Lancet, The</i> , 2006, 367, 832-838. | 6.3 | 162 |
| 258 | Tu-P10:482 Possible mechanism underlying the association between C-reactive protein and cardiovascular risk. <i>Atherosclerosis Supplements</i> , 2006, 7, 291. | 1.2 | 0 |
| 259 | The effects of continuous venovenous hemofiltration on coagulation activation. <i>Critical Care</i> , 2006, 10, R150. | 2.5 | 26 |
| 260 | Levels of intrinsic coagulation factors and the risk of myocardial infarction among men: opposite and synergistic effects of factors XI and XII. <i>Blood</i> , 2006, 108, 4045-4051. | 0.6 | 214 |
| 261 | Effect of Hypobaric Hypoxia, Simulating Conditions During Long-Haul Air Travel, on Coagulation, Fibrinolysis, Platelet Function, and Endothelial Activation. <i>JAMA - Journal of the American Medical Association</i> , 2006, 295, 2251. | 3.8 | 117 |
| 262 | Predilution Versus Postdilution During Continuous Venovenous Hemofiltration: A Comparison of Circuit Thrombogenesis. <i>ASAIO Journal</i> , 2006, 52, 416-422. | 0.9 | 27 |
| 263 | Proteomic analysis of high-density lipoprotein. <i>Proteomics</i> , 2006, 6, 721-730. | 1.3 | 169 |
| 264 | Discovery of thrombin activatable fibrinolysis inhibitor (TAFI). <i>Journal of Thrombosis and Haemostasis</i> , 2006, 4, 257-258. | 1.9 | 1 |
| 265 | The relationship between ABO blood group and the risk of bleeding during vitamin K antagonist treatment. <i>Journal of Thrombosis and Haemostasis</i> , 2006, 4, 1418-1420. | 1.9 | 24 |
| 266 | The differential association of conjugated equine estrogen and esterified estrogen with activated protein C resistance in postmenopausal women. <i>Journal of Thrombosis and Haemostasis</i> , 2006, 4, 1701-1706. | 1.9 | 12 |
| 267 | Vagus nerve stimulation inhibits activation of coagulation and fibrinolysis during endotoxemia in rats. <i>Journal of Thrombosis and Haemostasis</i> , 2006, 4, 1997-2002. | 1.9 | 166 |
| 268 | Anticoagulant and anti-inflammatory effects after peritoneal lavage with antithrombin in experimental polymicrobial peritonitis. <i>Journal of Thrombosis and Haemostasis</i> , 2006, 4, 2343-2351. | 1.9 | 13 |
| 269 | Endotoxaemia induces resistance to activated protein C in healthy humans. <i>British Journal of Haematology</i> , 2006, 134, 213-219. | 1.2 | 24 |
| 270 | Effects on Coagulation and Fibrinolysis Induced by Influenza in Mice With a Reduced Capacity to Generate Activated Protein C and a Deficiency in Plasminogen Activator Inhibitor Type 1. <i>Circulation Research</i> , 2006, 99, 1261-1269. | 2.0 | 77 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 271 | Mechanical ventilation affects alveolar fibrinolysis in LPS-induced lung injury. <i>European Respiratory Journal</i> , 2006, 28, 992-998. | 3.1 | 28 |
| 272 | Loss of Endothelial Glycocalyx During Acute Hyperglycemia Coincides With Endothelial Dysfunction and Coagulation Activation In Vivo. <i>Diabetes</i> , 2006, 55, 480-486. | 0.3 | 469 |
| 273 | Reduced plasma fibrinolytic potential is a risk factor for venous thrombosis. <i>Blood</i> , 2005, 105, 1102-1105. | 0.6 | 246 |
| 274 | Role of isoleucine residues 182 and 183 in thrombin-activatable fibrinolysis inhibitor. <i>Journal of Thrombosis and Haemostasis</i> , 2005, 3, 1293-1300. | 1.9 | 10 |
| 275 | The prognostic value of the D-dimer test in cancer patients treated with and without low-molecular-weight heparin. <i>Journal of Thrombosis and Haemostasis</i> , 2005, 3, 1531-1533. | 1.9 | 17 |
| 276 | Lupus anticoagulants and the risk of a first episode of deep venous thrombosis. <i>Journal of Thrombosis and Haemostasis</i> , 2005, 3, 1993-1997. | 1.9 | 183 |
| 277 | Effects of high plasma lipoprotein (a) levels on tissue factor pathway inhibitor and the protein C pathway. <i>Journal of Thrombosis and Haemostasis</i> , 2005, 3, 2123-2125. | 1.9 | 11 |
| 278 | Alveolar fibrinolytic capacity suppressed by injurious mechanical ventilation. <i>Intensive Care Medicine</i> , 2005, 31, 724-732. | 3.9 | 39 |
| 279 | Analysis of blood coagulation in mice: pre-analytical conditions and evaluation of a home-made assay for thrombin-antithrombin complexes. <i>Thrombosis Journal</i> , 2005, 3, 12. | 0.9 | 31 |
| 280 | The value of haemostatic markers in the triage of patients with chest pain presenting with a normal or non-diagnostic ECG. <i>Heart</i> , 2005, 91, 1215-1216. | 1.2 | 1 |
| 281 | Activation of Inflammation and Coagulation After Infusion of C-Reactive Protein in Humans. <i>Circulation Research</i> , 2005, 96, 714-716. | 2.0 | 208 |
| 282 | Absence of Thrombin-Activatable Fibrinolysis Inhibitor Protects against Sepsis-Induced Liver Injury in Mice. <i>Journal of Immunology</i> , 2005, 175, 6764-6771. | 0.4 | 56 |
| 283 | Lipopolysaccharide Is Transferred from High-Density to Low-Density Lipoproteins by Lipopolysaccharide-Binding Protein and Phospholipid Transfer Protein. <i>Infection and Immunity</i> , 2005, 73, 2321-2326. | 1.0 | 129 |
| 284 | Recombinant human activated protein C resets thrombin generation in patients with severe sepsis - a case control study. <i>Critical Care</i> , 2005, 9, R490. | 2.5 | 28 |
| 285 | Hypercoagulability in the metabolic syndrome. <i>Current Opinion in Pharmacology</i> , 2005, 5, 155-159. | 1.7 | 157 |
| 286 | Effects of Clopidogrel on the Rebound Hypercoagulable State After Heparin Discontinuation in Patients With Acute Coronary Syndromes. <i>Journal of the American College of Cardiology</i> , 2005, 46, 1582-1583. | 1.2 | 17 |
| 287 | Pentoxifylline reduces fibrin deposition and prolongs survival in neonatal hyperoxic lung injury. <i>Journal of Applied Physiology</i> , 2004, 97, 2014-2019. | 1.2 | 33 |
| 288 | Anti-Inflammatory and Anticoagulant Effects of Pravastatin in Patients With Type 2 Diabetes. <i>Diabetes Care</i> , 2004, 27, 468-473. | 4.3 | 39 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 289 | Generation and Characterization of a Highly Stable Form of Activated Thrombin-activable Fibrinolysis Inhibitor. <i>Journal of Biological Chemistry</i> , 2004, 279, 6620-6628. | 1.6 | 21 |
| 290 | Absence of mutations in the PCI gene in subfertile men. <i>Molecular Human Reproduction</i> , 2004, 10, 807-813. | 1.3 | 10 |
| 291 | Recombinant factor VIIa reverses the anticoagulant effect of the long-acting pentasaccharide idraparinux in healthy volunteers. <i>British Journal of Haematology</i> , 2004, 124, 653-658. | 1.2 | 99 |
| 292 | Recombinant nematode anticoagulant protein c2, an inhibitor of tissue factor/factor VIIa, attenuates coagulation and the interleukin-10 response in human endotoxemia. <i>Journal of Thrombosis and Haemostasis</i> , 2004, 2, 65-70. | 1.9 | 36 |
| 293 | Tumor growth and metastasis are not affected in thrombin-activatable fibrinolysis inhibitor-deficient mice. <i>Journal of Thrombosis and Haemostasis</i> , 2004, 2, 769-779. | 1.9 | 19 |
| 294 | Mechanism of action of carboxypeptidase U: staying above the threshold. <i>Journal of Thrombosis and Haemostasis</i> , 2004, 2, 414-415. | 1.9 | 1 |
| 295 | Protein C and S and inflammation in sickle cell disease. <i>American Journal of Hematology</i> , 2004, 76, 26-32. | 2.0 | 31 |
| 296 | New insights into factors affecting clot stability: a role for thrombin activatable fibrinolysis inhibitor (TAFI; plasma procarboxypeptidase B, plasma procarboxypeptidase U, procarboxypeptidase R). <i>Seminars in Hematology</i> , 2004, 41, 13-19. | 1.8 | 37 |
| 297 | Prospective validation of the International Society of Thrombosis and Haemostasis scoring system for disseminated intravascular coagulation*. <i>Critical Care Medicine</i> , 2004, 32, 2416-2421. | 0.4 | 419 |
| 298 | Thrombomodulin mutant mice with a strongly reduced capacity to generate activated protein C have an unaltered pulmonary immune response to respiratory pathogens and lipopolysaccharide. <i>Blood</i> , 2004, 103, 1702-1709. | 0.6 | 111 |
| 299 | Effect of second- and third-generation oral contraceptives on the protein C system in the absence or presence of the factor VLeiden mutation: a randomized trial. <i>Blood</i> , 2004, 103, 927-933. | 0.6 | 135 |
| 300 | Dominant factor XI deficiency caused by mutations in the factor XI catalytic domain. <i>Blood</i> , 2004, 104, 128-134. | 0.6 | 92 |
| 301 | The Role of the Fibrinolytic System in Corneal Angiogenesis. <i>Angiogenesis</i> , 2003, 6, 311-316. | 3.7 | 15 |
| 302 | Procoagulant protein levels are differentially increased during human endotoxemia. <i>Journal of Thrombosis and Haemostasis</i> , 2003, 1, 1019-1023. | 1.9 | 43 |
| 303 | Thrombin-activatable fibrinolysis inhibitor (TAFI, plasma procarboxypeptidase B, procarboxypeptidase R,) Tj ETQq1 1 0.784314 rgBT /Ove 155 | 1.9 | 155 |
| 304 | Impaired healing of cutaneous wounds and colonic anastomoses in mice lacking thrombin-activatable fibrinolysis inhibitor. <i>Journal of Thrombosis and Haemostasis</i> , 2003, 1, 2087-2096. | 1.9 | 82 |
| 305 | Recombinant factor VIIa reverses the in vitro and ex vivo anticoagulant and profibrinolytic effects of fondaparinux. <i>Journal of Thrombosis and Haemostasis</i> , 2003, 1, 2368-2373. | 1.9 | 70 |
| 306 | The impact on coagulation of an intravenous loading dose in addition to a subcutaneous regimen of low-molecular-weight heparin in the initial treatment of acute coronary syndromes. <i>Journal of the American College of Cardiology</i> , 2003, 42, 424-427. | 1.2 | 14 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 307 | Potential Role of Platelets in Endothelial Damage Observed During Treatment With Cisplatin, Gemcitabine, and the Angiogenesis Inhibitor SU5416. <i>Journal of Clinical Oncology</i> , 2003, 21, 2192-2198. | 0.8 | 113 |
| 308 | Effects of IC14, an Anti-CD14 Antibody, on Coagulation and Fibrinolysis during Low-Grade Endotoxemia in Humans. <i>Journal of Infectious Diseases</i> , 2003, 187, 55-61. | 1.9 | 39 |
| 309 | Distribution and Kinetics of Lipoprotein-Bound Lipoteichoic Acid. <i>Infection and Immunity</i> , 2003, 71, 3280-3284. | 1.0 | 86 |
| 310 | Identification of thrombin activatable fibrinolysis inhibitor (TAFI) in human platelets. <i>Blood</i> , 2003, 101, 4844-4846. | 0.6 | 102 |
| 311 | Amyloid endostatin induces endothelial cell detachment by stimulation of the plasminogen activation system. <i>Molecular Cancer Research</i> , 2003, 1, 561-8. | 1.5 | 30 |
| 312 | Characterization of the H-kininogen-binding Site on Factor XI. <i>Journal of Biological Chemistry</i> , 2002, 277, 4892-4899. | 1.6 | 64 |
| 313 | Analysis of Coagulation Cascade and Endothelial Cell Activation During Inhibition of Vascular Endothelial Growth Factor/Vascular Endothelial Growth Factor Receptor Pathway in Cancer Patients. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2002, 22, 1500-1505. | 1.1 | 141 |
| 314 | Ability of Recombinant Factor VIIa to Reverse the Anticoagulant Effect of the Pentasaccharide Fondaparinux in Healthy Volunteers. <i>Circulation</i> , 2002, 106, 2550-2554. | 1.6 | 311 |
| 315 | Effect of second- and third-generation oral contraceptives on fibrinolysis in the absence or presence of the factor V Leiden mutation. <i>Blood Coagulation and Fibrinolysis</i> , 2002, 13, 373-381. | 0.5 | 15 |
| 316 | Inhibition of fibrinolysis by recombinant factor VIIa in plasma from patients with severe hemophilia A. <i>Blood</i> , 2002, 99, 175-179. | 0.6 | 159 |
| 317 | Effects of Second and Third Generation Oral Contraceptives and their Respective Progestagens on the Coagulation System in the Absence or Presence of the Factor V Leiden Mutation. <i>Thrombosis and Haemostasis</i> , 2002, 87, 199-205. | 1.8 | 58 |
| 318 | Plasmin-Mediated Activation and Inactivation of Thrombin-Activatable Fibrinolysis Inhibitor. <i>Biochemistry</i> , 2002, 41, 6688-6696. | 1.2 | 71 |
| 319 | Role of Zinc Ions in Activation and Inactivation of Thrombin-Activatable Fibrinolysis Inhibitor. <i>Biochemistry</i> , 2002, 41, 1211-1216. | 1.2 | 20 |
| 320 | Rebound thrombin generation after heparin therapy in unstable angina. <i>Journal of the American College of Cardiology</i> , 2002, 39, 811-817. | 1.2 | 54 |
| 321 | No effect of acenocoumarol therapy on levels of endothelial activation markers in sickle cell disease. <i>American Journal of Hematology</i> , 2002, 71, 53-55. | 2.0 | 10 |
| 322 | Protein C inhibitor (plasminogen activator inhibitor-3) and the risk of venous thrombosis. <i>British Journal of Haematology</i> , 2002, 118, 604-609. | 1.2 | 15 |
| 323 | Activation of coagulation factor XI, without detectable contact activation in dengue haemorrhagic fever. <i>British Journal of Haematology</i> , 2001, 113, 94-99. | 1.2 | 38 |
| 324 | Complexes of anti-prothrombin antibodies and prothrombin cause lupus anticoagulant activity by competing with the binding of clotting factors for catalytic phospholipid surfaces. <i>British Journal of Haematology</i> , 2001, 113, 621-629. | 1.2 | 66 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 325 | Reduced activity of TAFI (thrombin-activatable fibrinolysis inhibitor) in acute promyelocytic leukaemia. British Journal of Haematology, 2000, 108, 518-523. | 1.2 | 66 |
| 326 | Protein C inhibitor in platelets?. Blood, 2000, 95, 1880-1881. | 0.6 | 0 |
| 327 | C4b-Binding Protein Protects Coagulation Factor Va from Inactivation by Activated Protein C. Biochemistry, 2000, 39, 14543-14548. | 1.2 | 13 |
| 328 | R-205. Protein C inhibitor/plasminogen activator inhibitor-3 is present in the mouse ovary and is down-regulated by human chorionic gonadotrophin treatment. Human Reproduction, 1999, 14, 370-371. | 0.4 | 0 |
| 329 | Binding of amyloid β precursor protein to coagulation factor XIa in vivo may favour haemorrhagic stroke. Journal of Neurology, 1998, 245, 111-115. | 1.8 | 6 |
| 330 | Inhibition of serine proteases by reactive site mutants of protein C inhibitor (plasminogen activator) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 5 | 1.1 | 7 |
| 331 | Factor XI Activation by Meizothrombin: Stimulation by Phospholipid Vesicles Containing both Phosphatidylserine and Phosphatidylethanolamine. Thrombosis and Haemostasis, 1997, 78, 834-839. | 1.8 | 27 |
| 332 | Role of the A+ Helix in Heparin Binding to Protein C Inhibitor. Thrombosis and Haemostasis, 1996, 75, 760-766. | 1.8 | 32 |
| 333 | Effect of Heparin on the Activation of Factor XI by Fibrin-bound Thrombin. Thrombosis and Haemostasis, 1996, 76, 347-353. | 1.8 | 15 |
| 334 | The Interaction of Protein S with the Phospholipid Surface Is Essential for the Activated Protein C-independent Activity of Protein S. Thrombosis and Haemostasis, 1996, 76, 397-403. | 1.8 | 63 |
| 335 | Apple four in human blood coagulation factor XI mediates dimer formation. Biochemistry, 1992, 31, 4680-4684. | 1.2 | 87 |
| 336 | Inhibition of human blood coagulation factor XIa by C.hivin.1 inhibitor. Biochemistry, 1988, 27, 959-963. | 1.2 | 34 |
| 337 | Inactivation of human plasma kallikrein and factor XIa by protein C inhibitor. Biochemistry, 1988, 27, 4231-4237. | 1.2 | 122 |
| 338 | Oral contraceptives reduce total protein S, but not free protein S.. Thrombosis Research, 1987, 45, 109-114. | 0.8 | 45 |
| 339 | Inhibition of human blood coagulation factor Xa by .alpha.2-macroglobulin. Biochemistry, 1987, 26, 5932-5937. | 1.2 | 37 |
| 340 | Characterization of human protein S. Thrombosis Research, 1986, 41, 135. | 0.8 | 1 |
| 341 | Analysis of tumour-localizing haematoporphyrin derivative by high-performance liquid chromatography and fast-atom bombardment mass spectrometry. Journal of Chromatography A, 1986, 352, 231-239. | 1.8 | 13 |
| 342 | Nanobodies against factor α XI and apple 3 domain inhibit binding of factor α IX and reveal a novel binding site for high molecular weight kininogen. Journal of Thrombosis and Haemostasis, 0, , . | 1.9 | 5 |