

Guillaume Louis

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4410879/publications.pdf>

Version: 2024-02-01

35
papers

2,621
citations

516561

16
h-index

345118

36
g-index

40
all docs

40
docs citations

40
times ranked

3979
citing authors

#	ARTICLE	IF	CITATIONS
1	Comparison of SARS-CoV-2 Variants of Concern Alpha (B.1.1.7) vs. Beta (B.1.351) in Critically Ill Patients: A Multicenter Cohort Study. <i>Frontiers in Medicine</i> , 2022, 9, 828402.	1.2	5
2	Acute kidney injury in severe SARS-CoV-2 infection: An experience report in Eastern France. <i>Anaesthesia, Critical Care & Pain Medicine</i> , 2021, 40, 100788.	0.6	5
3	Clinical characteristics and day-90 outcomes of 4244 critically ill adults with COVID-19: a prospective cohort study. <i>Intensive Care Medicine</i> , 2021, 47, 60-73.	3.9	597
4	SARS-Cov-2 fulminant myocarditis: an autopsy and histopathological case study. <i>International Journal of Legal Medicine</i> , 2021, 135, 577-581.	1.2	63
5	Performance of the ROX index to predict intubation in immunocompromised patients receiving high-flow nasal cannula for acute respiratory failure. <i>Annals of Intensive Care</i> , 2021, 11, 17.	2.2	26
6	Prognostic factors associated with six month mortality of critically ill elderly patients admitted to the intensive care unit with severe acute cholangitis. <i>Hpb</i> , 2021, 23, 459-467.	0.1	3
7	Comparison of two delayed strategies for renal replacement therapy initiation for severe acute kidney injury (AKIKI 2): a multicentre, open-label, randomised, controlled trial. <i>Lancet, The</i> , 2021, 397, 1293-1300.	6.3	106
8	Chronic use of renin-angiotensin-aldosterone system blockers and mortality in COVID-19: A multicenter prospective cohort and literature review. <i>Fundamental and Clinical Pharmacology</i> , 2021, 35, 1141-1158.	1.0	4
9	Infection related catheter complications in patients undergoing prone positioning for acute respiratory distress syndrome: an exposed/unexposed study. <i>BMC Infectious Diseases</i> , 2021, 21, 534.	1.3	4
10	Mental health and stress among ICU healthcare professionals in France according to intensity of the COVID-19 epidemic. <i>Annals of Intensive Care</i> , 2021, 11, 90.	2.2	29
11	Preliminary data on severe SARS-CoV-2 infection caused by the 501Y.V2 variant. <i>Anaesthesia, Critical Care & Pain Medicine</i> , 2021, 40, 100890.	0.6	4
12	The AKIKI 2 trial: a case for strategy of initiation instead of timing – Authors' reply. <i>Lancet, The</i> , 2021, 398, 1215-1216.	6.3	2
13	A Prospective Cohort Study to Identify Clinical, Biological, and Imaging Features That Predict the Etiology of Acute Encephalitis. <i>Clinical Infectious Diseases</i> , 2021, 73, 264-270.	2.9	14
14	End of life in the critically ill patient: evaluation of experience of end of life by caregivers (EOLE) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 22	2.2	5
15	Liberal or Conservative Oxygen Therapy for Acute Respiratory Distress Syndrome. <i>New England Journal of Medicine</i> , 2020, 382, 999-1008.	13.9	290
16	Epidemiology and outcome of patients admitted to intensive care after anaphylaxis in France: a retrospective multicentre study. <i>British Journal of Anaesthesia</i> , 2020, 125, 1025-1033.	1.5	15
17	Influence of deprivation on initial severity and prognosis of patients admitted to the ICU: the prospective, multicentre, observational IVOIRE cohort study. <i>Annals of Intensive Care</i> , 2020, 10, 20.	2.2	16
18	Disseminated intravascular coagulation in pneumococemia. <i>Intensive Care Medicine</i> , 2019, 45, 698-699.	3.9	0

#	ARTICLE	IF	CITATIONS
19	Effect of fluid balance control in critically ill patients: Design of the stepped wedge trial POINCARE-2. Contemporary Clinical Trials, 2019, 83, 109-116.	0.8	6
20	Center effect in intubation risk in critically ill immunocompromised patients with acute hypoxemic respiratory failure. Critical Care, 2019, 23, 306.	2.5	11
21	The Artificial Kidney Initiation in Kidney Injury 2 (AKIKI2): study protocol for a randomized controlled trial. Trials, 2019, 20, 726.	0.7	15
22	Outcomes of Stenotrophomonas maltophilia hospital-acquired pneumonia in intensive care unit: a nationwide retrospective study. Critical Care, 2019, 23, 371.	2.5	41
23	Timing of Renal-Replacement Therapy in Patients with Acute Kidney Injury and Sepsis. New England Journal of Medicine, 2018, 379, 1431-1442.	13.9	417
24	Effect of High-Flow Nasal Oxygen vs Standard Oxygen on 28-Day Mortality in Immunocompromised Patients With Acute Respiratory Failure. JAMA - Journal of the American Medical Association, 2018, 320, 2099.	3.8	202
25	Epinephrine Versus Norepinephrine for Cardiogenic Shock After Acute Myocardial Infarction. Journal of the American College of Cardiology, 2018, 72, 173-182.	1.2	282
26	High-flow nasal oxygen vs. standard oxygen therapy in immunocompromised patients with acute respiratory failure: study protocol for a randomized controlled trial. Trials, 2018, 19, 157.	0.7	11
27	Functional outcome of patients with prolonged hypoglycemic encephalopathy. Annals of Intensive Care, 2017, 7, 54.	2.2	17
28	Grey Turner's sign associated with hemorrhagic pancreatitis. Intensive Care Medicine, 2017, 43, 265-266.	3.9	2
29	Limitation et arrêt de traitement: collaboration entre une équipe mobile de soins palliatifs et un service de réanimation. Médecine Palliative, 2016, 15, 78-85.	0.0	5
30	Functional outcome of prolonged refractory status epilepticus. Critical Care, 2015, 19, 199.	2.5	43
31	Outcomes of patients admitted to intensive care units for acute manifestation of small-vessel vasculitis: a multicenter, retrospective study. Critical Care, 2015, 20, 27.	2.5	28
32	Angioedema induced by angiotensin converting enzyme inhibitors, potentiated by m-TOR inhibitors: successful treatment with icatibant. Intensive Care Medicine, 2014, 40, 893-894.	3.9	18
33	The epidemiology of septic shock in French intensive care units: the prospective multicenter cohort EPISS study. Critical Care, 2013, 17, R65.	2.5	156
34	Profile of the Risk of Death After Septic Shock in the Present Era. Critical Care Medicine, 2013, 41, 2600-2609.	0.4	84
35	Long-term outcome of patients hospitalized in intensive care units with central or extrapontine myelinolysis*. Critical Care Medicine, 2012, 40, 970-972.	0.4	75